

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 February 2003 (27.02.2003)

PCT

(10) International Publication Number
WO 03/015774 A1

(51) International Patent Classification: A61K 31/425,
C07D 417/12, 233/48, 285/12, 213/75, 241/20, 231/40,
237/20, 239/42, 307/66, A61P 03/10

(21) International Application Number: PCT/GB02/03745

(22) International Filing Date: 15 August 2002 (15.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0102764-8 17 August 2001 (17.08.2001) SE

(71) Applicant (for all designated States except MG, US): AS-
TRAZENECA AB [SE/SE]; Sodertalje, S-151 85 (SE).

(71) Applicant (for MG only): ASTRAZENECA UK LIM-
ITED [GB/GB]; 15 Stanhope Gate, London, Greater Lon-
don W1Y 6LN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BOYD, Scott
[GB/GB]; Alderley Park, Macclesfield, Cheshire SK10
4TG (GB). CAULKETT, Peter, William, Rodney
[GB/GB]; Alderley Park, Macclesfield, Cheshire SK10
4TG (GB). HARGREAVES, Rodney, Brian [GB/GB];
Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
BOWKER, Suzanne, Saxon [GB/GB]; Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB). JAMES, Roger
[GB/GB]; Alderley Park, Macclesfield, Cheshire SK10
4TG (GB). JOHNSTONE, Craig [GB/GB]; Alderley

Park, Macclesfield, Cheshire SK10 4TG (GB). JONES,
Clifford, David [GB/GB]; Alderley Park, Macclesfield,
Cheshire SK10 4TG (GB). MCKERRECHER, Darren
[GB/GB]; Alderley Park, Macclesfield, Cheshire SK10
4TG (GB). BLOCK, Michael, Howard [GB/US]; 35
Gatehouse Drive, Waltham, MA 02451 (US).

(74) Agent: BRYANT, Tracey, et al; Astrazeneca, Global In-
tellectual Property, Mereside, Alderley Park, Macclesfield,
Cheshire SK10 4TG (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: COMPOUNDS EFFECTING GLUCOKINASE



(57) Abstract: The invention relates to the use of a compound of Formula (I) or a salt, solvate or prodrug thereof, wherein R¹, R², R³, n and m are as described in the specification, in the preparation of a medicament for the treatment or prevention of a disease condition mediated through glucokinase (GLK), such as type 2 diabetes. The invention also relates to a novel group of compounds of Formula (I) and to methods for preparing compounds of Formula (I).

WO 03/015774 A1

COMPOUNDS EFFECTING GLUCOKINASE

The present invention relates to the use of a group of benzamide compounds in the preparation of a medicament for use in the treatment or prevention of a disease or medical condition mediated through glucokinase (GLK), leading to a decreased glucose threshold for insulin secretion. In addition the compounds are predicted to lower blood glucose by increasing hepatic glucose uptake. Such compounds may have utility in the treatment of Type 2 diabetes and obesity. The invention also relates to pharmaceutical compositions comprising said benzamide compound, a sub-group of novel compounds of said benzamide compounds, and the use of such a compound in the conditions described above.

In the pancreatic β -cell and liver parenchymal cells the main plasma membrane glucose transporter is GLUT2. Under physiological glucose concentrations the rate at which GLUT2 transports glucose across the membrane is not rate limiting to the overall rate of glucose uptake in these cells. The rate of glucose uptake is limited by the rate of phosphorylation of glucose to glucose-6-phosphate (G-6-P) which is catalysed by glucokinase (GLK) [1]. GLK has a high (6-10mM) K_m for glucose and is not inhibited by physiological concentrations of G-6-P [1]. GLK expression is limited to a few tissues and cell types, most notably pancreatic β -cells and liver cells (hepatocytes) [1]. In these cells GLK activity is rate limiting for glucose utilisation and therefore regulates the extent of glucose induced insulin secretion and hepatic glycogen synthesis. These processes are critical in the maintenance of whole body glucose homeostasis and both are dysfunctional in diabetes [2].

In one sub-type of diabetes, Type 2 maturity-onset diabetes of the young (MODY-2), the diabetes is caused by GLK loss of function mutations [3, 4]. Hyperglycaemia in MODY-2 patients results from defective glucose utilisation in both the pancreas and liver [5]. Defective glucose utilisation in the pancreas of MODY-2 patients results in a raised threshold for glucose stimulated insulin secretion. Conversely, rare activating mutations of GLK reduce this threshold resulting in familial hyperinsulinism [6, 7]. In addition to the reduced GLK activity observed in MODY-2 diabetics, hepatic glucokinase activity is also decreased in type 2 diabetics [8]. Importantly, global or liver selective overexpression of GLK prevents or reverses the development of the diabetic phenotype in both dietary and genetic models of the disease [9-12]. Moreover, acute treatment of type 2 diabetics with fructose improves glucose tolerance through stimulation of hepatic glucose utilisation [13]. This effect is believed to be

- 2 -

mediated through a fructose induced increase in cytosolic GLK activity in the hepatocyte by the mechanism described below [13].

Hepatic GLK activity is inhibited through association with GLK regulatory protein (GLKRP). The GLK/GLKRP complex is stabilised by fructose-6-phosphate (F6P) binding to the GLKRP and destabilised by displacement of this sugar phosphate by fructose-1-phosphate (F1P). F1P is generated by fructokinase mediated phosphorylation of dietary fructose. Consequently, GLK/GLKRP complex integrity and hepatic GLK activity is regulated in a nutritionally dependent manner as F6P is elevated in the post-absorptive state whereas F1P predominates in the post-prandial state. In contrast to the hepatocyte, the pancreatic β -cell expresses GLK in the absence of GLKRP. Therefore, β -cell GLK activity is regulated exclusively by the availability of its substrate, glucose. Small molecules may activate GLK either directly or through destabilising the GLK/GLKRP complex. The former class of compounds are predicted to stimulate glucose utilisation in both the liver and the pancreas whereas the latter are predicted to act exclusively in the liver. However, compounds with either profile are predicted to be of therapeutic benefit in treating Type 2 diabetes as this disease is characterised by defective glucose utilisation in both tissues.

GLK and GLKRP and the K_{ATP} channel are expressed in neurones of the hypothalamus, a region of the brain that is important in the regulation of energy balance and the control of food intake [14-18]. These neurones have been shown to express orectic and anorectic neuropeptides [15, 19, 20] and have been assumed to be the glucose-sensing neurones within the hypothalamus that are either inhibited or excited by changes in ambient glucose concentrations [17, 19, 21, 22]. The ability of these neurones to sense changes in glucose levels is defective in a variety of genetic and experimentally induced models of obesity [23-28]. Intracerebroventricular (icv) infusion of glucose analogues, that are competitive inhibitors of glucokinase, stimulate food intake in lean rats [29, 30]. In contrast, icv infusion of glucose suppresses feeding [31]. Thus, small molecule activators of GLK may decrease food intake and weight gain through central effects on GLK. Therefore, GLK activators may be of therapeutic use in treating eating disorders, including obesity, in addition to diabetes. The hypothalamic effects will be additive or synergistic to the effects of the same compounds acting in the liver and/or pancreas in normalising glucose homeostasis, for the treatment of Type 2 diabetes. Thus the GLK/GLKRP system can be described as a potential "Diabesity" target (of benefit in both Diabetes and Obesity).

In WO0058293 and WO01/44216 (Roche), a series of benzylcarbamoyl compounds are described as glucokinase activators. The mechanism by which such compounds activate GLK is assessed by measuring the direct effect of such compounds in an assay in which GLK activity is linked to NADH production, which in turn is measured optically - see details of the
5 *in vitro* assay described in Example A. Compounds of the present invention may activate GLK directly or may activate GLK by inhibiting the interaction of GLKRP with GLK. The latter mechanism offers an important advantage over direct activators of GLK in that they will not cause the severe hypoglycaemic episodes predicted after direct stimulation. Many compounds of the present invention may show favourable selectivity compared to known GLK activators.
10 WO9622282, WO9622293, WO9622294, WO9622295, WO9749707 and WO9749708 disclose a number of intermediates used in the preparation of compounds useful as vasopressin agents which are structurally similar to those disclosed in the present invention. Structurally similar compounds are also disclosed in WO9641795 and JP8143565 (vasopressin antagonism), in JP8301760 (skin damage prevention) and in EP619116
15 (osetopathy).


WO01/12621 describes the preparation of as isoxazolympyrimidines and related compounds as inhibitors of cJUN N-terminal kinases, and pharmaceutical compositions containing such compounds.

Cushman *et al* [Bioorg Med Chem Lett (1991) 1(4), 211-14] describe the synthesis of
20 pyridine-containing stilbenes and amides and their evaluation as protein-tyrosine kinase inhibitors. Rogers *et al* [J Med Chem (1981) 24(11) 1284-7] describe mesoionic purinone analogs as inhibitors of cyclic-AMP phosphodiesterase.

WO00/26202 describes the preparation of 2-amino-thiazole derivatives as antitumour agents. GB 2331748 describes the preparation of insecticidal thiazole derivatives.
25 WO96/36619 describes the preparation of aminothiazole derivatives as ameliorating agents for digestive tract movements. US 5466715 and US 5258407 describe the preparation of 3,4-disubstituted phenol immunostimulants. JP 58069812 describes hypoglycemic pharmaceuticals containing benzamide derivatives. US 3950351 describes 2-benzamido-5-nitrothiazoles and Cavier *et al* [Eur J Med Chem – Chim Ther (1978) 13(6), 539-43] discuss
30 the biological interest of these compounds.

- 4 -

We present as a feature of the invention the use of a compound of Formula (I) or a salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the treatment or prevention of a disease or medical condition mediated through GLK:



Formula (I)

wherein

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

and $n + m > 0$;

- each R^1 is independently selected from OH, $-(CH_2)_{1-4}OH$, $-CH_3F_a$, $-(CH_2)_{1-4}CH_3F_a$, $-OCH_3F_a$, halo, $C_{1-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, NH_2 , $-NH-C_{1-4}alkyl$, $-N-di-(C_{1-4}alkyl)$, CN, formyl, phenyl or heterocyclyl optionally substituted by $C_{1-6}alkyl$;

each R^2 is the group $Y-X-$

- wherein each X is a linker independently selected from:

$-O-Z-$, $-O-Z-O-Z-$, $-C(O)O-Z-$, $-OC(O)-Z-$, $-S-Z-$, $-SO-Z-$, $-SO_2-Z-$, $-N(R^6)-Z-$, $-N(R^6)SO_2-Z-$, $-SO_2N(R^6)-Z-$, $-CH=CH-Z-$, $-C\equiv C-Z-$, $-N(R^6)CO-Z-$, $-CON(R^6)-Z-$, $-C(O)N(R^6)S(O)_2-Z-$, $-S(O)_2N(R^6)C(O)-Z-$, $-C(O)-Z-$, $-Z-$, $-C(O)-Z-O-Z-$, $-N(R^6)-C(O)-Z-O-Z-$, $-O-Z-N(R^6)-Z-$, $-O-C(O)-Z-O-Z-$ or a direct bond;

each Z is independently a direct bond, $C_{2-6}alkenylene$ or a group of the formula

$-(CH_2)_p-C(R^{6a})_2-(CH_2)_q-$;

each Y is independently selected from aryl- Z^1- , heterocyclyl- Z^1- , $C_{3-7}cycloalkyl-Z^1-$,

$C_{1-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, $-(CH_2)_{1-4}CH_3F_a$ or $-CH(OH)CH_3F_a$; wherein

- each Y is independently optionally substituted by up to 3 R^4 groups;

each R^4 is independently selected from halo, $-CH_3F_a$, CN, NH_2 , $C_{1-6}alkyl$,

$-OC_{1-6}alkyl$, $-COOH$, $-C(O)OC_{1-6}alkyl$, OH or phenyl optionally substituted by $C_{1-6}alkyl$ or $-C(O)OC_{1-6}alkyl$,

- 5 -

or R^5-X^1 -, where X^1 is independently as defined in X above and R^5 is selected from hydrogen, C_{1-6} alkyl, $-CH_3-aF_a$, phenyl, naphthyl, heterocyclyl or C_{3-7} cycloalkyl; and R^5 is optionally substituted by one or more substituents independently selected from: halo, C_{1-6} alkyl, $-OC_{1-6}$ alkyl, $-CH_3-aF_a$, CN, OH, NH_2 , COOH, or $-C(O)OC_{1-6}$ alkyl,

5 each Z^1 is independently a direct bond, C_{2-6} alkenylene or a group of the formula $-(CH_2)_p-C(R^{6a})_2-(CH_2)_q$;

R^3 is selected from phenyl or a heterocyclyl, and R^3 is optionally substituted by one or more R^7 groups;

10 R^6 is independently selected from hydrogen, C_{1-6} alkyl or $-C_{2-4}$ alkyl-O- C_{1-4} alkyl;
 R^{6a} is independently selected from hydrogen, halo, C_{1-6} alkyl or $-C_{2-4}$ alkyl-O- C_{1-4} alkyl;
each R^7 is independently selected from:

C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, $(CH_2)_{0-3}$ aryl, $(CH_2)_{0-3}$ heterocyclyl, $(CH_2)_{0-3}C_{3-7}$ cycloalkyl, OH, C_{1-6} alkyl-OH, halo, C_{1-6} alkyl-halo, OC_{1-6} alkyl, $(CH_2)_{0-3}S(O)_{0-2}R^8$, SH, SO_3 , thioxo, NH_2 , CN, $(CH_2)_{0-3}NHSO_2R^8$, $(CH_2)_{0-3}COOH$, $(CH_2)_{0-3}-O-(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)OR^8$, $(CH_2)_{0-3}C(O)NH_2$, $(CH_2)_{0-3}C(O)NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NHC(O)(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)NHSO_2-R^8$ and $(CH_2)_{0-3}SO_2NHC(O)-R^8$ wherein an alkyl chain, cycloalkyl ring or heterocyclyl ring within R^7 is optionally substituted by one of more substituents independently selected from: C_{1-4} alkyl, OH, halo, CN, NH_2 , $N-C_{1-4}$ alkylamino, N,N -di- C_{1-4} alkylamino and OC_{1-4} alkyl;

15 $(CH_2)_{0-3}S(O)_{0-2}R^8$, SH, SO_3 , thioxo, NH_2 , CN, $(CH_2)_{0-3}NHSO_2R^8$, $(CH_2)_{0-3}COOH$, $(CH_2)_{0-3}-O-(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)OR^8$, $(CH_2)_{0-3}C(O)NH_2$, $(CH_2)_{0-3}C(O)NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NHC(O)(CH_2)_{0-3}R^8$; $(CH_2)_{0-3}C(O)NHSO_2-R^8$ and $(CH_2)_{0-3}SO_2NHC(O)-R^8$ wherein an alkyl chain, cycloalkyl ring or heterocyclyl ring within R^7 is optionally substituted by one of more substituents independently selected from: C_{1-4} alkyl, OH, halo, CN, NH_2 , $N-C_{1-4}$ alkylamino, N,N -di- C_{1-4} alkylamino and OC_{1-4} alkyl;

20

R^8 is selected from hydrogen, C_{1-6} alkyl, aryl, heterocyclyl, C_{3-7} cycloalkyl, OH, C_{1-6} alkyl-OH, COOH, $C(O)OC_{1-6}$ alkyl, $N(R^6)C_{1-6}$ alkyl, OC_{1-6} alkyl, C_{0-6} alkylOC(O) C_{1-6} alkyl, $C(OH)(C_{1-6}alkyl)C_{1-6}alkyl$; wherein an alkyl chain or aryl, heterocyclyl or cycloalkyl ring within R^8 is optionally substituted by one of more substituents independently selected from: C_{1-4} alkyl, OH, halo, CN, NH_2 , $-NH-C_{1-4}$ alkyl, $-N$ -di- $(C_{1-4}alkyl)$ and $OC_{1-4}alkyl$;

25

each a is independently 1, 2 or 3;

30 p is an integer between 0 and 3;


q is an integer between 0 and 3;

and $p + q < 4$.

- 6 -


provided that when R^3 is 2-pyridyl and X is other than -Z-, -C(O)-Z-O-Z-,
 -N(R^6)-C(O)-Z-O-Z- or -O-Z-N(R^6)-Z-, then R^3 cannot be mono-substituted at the 5-position
 with an R^7 group selected from COOH or C(O)OC₁₋₆alkyl.

For the avoidance of doubt the numbering in the above proviso is relative to the amide
 5 bond attached to the pyridine ring, thus R^3 in the proviso relates to a group of the following
 structure:



wherein  represents the point of attachment to the amide group in Formula (I).

According to a further feature of the invention there is provided the use of a compound
 10 of Formula (Ia) or a salt thereof, in the preparation of a medicament for use in the treatment or
 prevention of a disease or medical condition mediated through GLK:



Formula (Ia)

wherein

15 m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

and $n + m > 0$;

each R^1 is independently selected from OH, (CH₂)₁₋₄OH, CH₃₋₃F_a, (CH₂)₁₋₄CH₃₋₃F_a,

OCH₃₋₃F_a, halo, C₁₋₆ alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, NH₂, N(C₁₋₆alkyl)C₁₋₆alkyl,

20 CN, phenyl or a heterocyclyl optionally substituted by C₁₋₆alkyl;

each R^2 is the group Y-X-

wherein each X is a linker independently selected from

-O(CH₂)₀₋₃-, -(CH₂)₀₋₃O-, -C(O)O(CH₂)₀₋₃-, -S(CH₂)₀₋₃-, -SO(CH₂)₀₋₃-,


-SO₂(CH₂)₀₋₃-, -NHSO₂-, -SO₂NH-, -N(CH₂)₀₋₃-, -N(CH₂)₁₋₃O(CH₂)₀₋₃-, -(CH₂)₁₋₄-,

25 -CH=CH(CH₂)₀₋₂-, -C≡C(CH₂)₀₋₂-, -NHCO-, -CONH-;

- 7 -

- each Y is independently selected from phenyl(CH₂)₀₋₂, naphthyl(CH₂)₀₋₂, heterocyclyl(CH₂)₀₋₂, C₃₋₇ cycloalkyl(CH₂)₀₋₂, C₁₋₆ alkyl, OC₁₋₆alkyl, C₂₋₆ alkenyl, C₂₋₆alkynyl, or CH(OH)CH_{3-a}F_a;
- each Y is independently optionally substituted by one or more R⁴ groups;
- 5 each R⁴ is independently selected from halo, CH_{3-a}F_a, OCH_{3-a}F_a, CN, NH₂, C₁₋₆alkyl, OC₁₋₆alkyl, COOH, (CH₂)₀₋₃COOH, O(CH₂)₀₋₃COOH, C(O)OC₁₋₆alkyl, C₁₋₆alkylC(O)OC₁₋₆alkyl, CO-phenyl, CONH₂, CONH-phenyl, SO₂NH₂, SO₂C₁₋₆alkyl, OH, or phenyl optionally substituted by one or more R⁵ groups, or R^{6b}-X-;
- 10 R⁵ is selected from hydrogen, C₁₋₆alkyl or C(O)OC₁₋₆alkyl, R^{6b} is selected from hydrogen, C₁₋₆alkyl, CH_{3-a}F_a phenyl, naphthyl, heterocyclyl or C₃₋₇cycloalkyl; and R^{6b} is optionally substituted by halo, C₁₋₆alkyl, CH_{3-a}F_a, CN, NH₂, COOH and COOC₁₋₆alkyl;
- each a is independently 1, 2 or 3;
- 15 R³ is selected from phenyl or a heterocyclyl, and R³ is optionally substituted by one or more R⁷ groups;
- each R⁷ is independently selected from:
- C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, heterocyclyl, (CH₂)₀₋₃C₃₋₇cycloalkyl, OH, C₁₋₆alkyl-OH, halo, C₁₋₆alkyl-halo, OC₁₋₆alkyl, SC₁₋₆alkyl, SH, SO₃, NH₂, CN, NHCHO, NSO₂C₁₋₆alkyl, (CH₂)₀₋₃COOH, (CH₂)₀₋₃C(O)OC₁₋₆alkyl,
- 20 (CH₂)₀₋₃CONH₂, (CH₂)₀₋₃CON(CH₂)₀₋₃R⁸, (CH₂)₀₋₃NH(CH₂)₀₋₃R⁸, (CH₂)₀₋₃NHC(O)(CH₂)₀₋₃R⁸;
- R⁸ is selected from hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, OH, C₁₋₆alkyl-OH, COOH, C(O)OC₁₋₆alkyl, N(C₀₋₆ alkyl)C₁₋₆ alkyl, O(C₀₋₆ alkyl)C₁₋₆alkyl,
- 25 C₀₋₆alkylOC(O)C₁₋₆alkyl, C(OH)(C₁₋₆alkyl)C₁₋₆alkyl;
- provided that when R³ is pyridine, then R⁷ is other than COOH or COOC₁₋₆alkyl.

According to a further feature of the invention there is provided a compound of Formula (Ib) or a salt, solvate or pro-drug thereof;



Formula (Ib)

5 wherein

m is 0, 1 or 2;

n is 1, 2 or 3;

and $n + m$ is 2 or 3;

each R^1 is independently selected from OH, $-(CH_2)_{1-4}OH$, $-CH_2F$, $-(CH_2)_{1-4}CH_2F$,

- 10 $-OCH_2F$, halo, OCH_3 , C_2H_5O , $CH_3C(O)O-$, $C_{1-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, $-NH-C_{1-4}alkyl$, $-N-di-(C_{1-4}alkyl)$, CN, formyl, phenyl or heterocyclyl optionally substituted by $C_{1-6}alkyl$;


each R^2 is the group $Y-X-$

with the proviso that $Y-X-$ cannot be CH_3O , C_2H_5O or $CH_3C(O)O-$;

- 15 wherein each X is a linker independently selected from:

$-O-Z-$, $-O-Z-O-Z-$, $-C(O)O-Z-$, $-OC(O)-Z-$, $-S-Z-$, $-SO-Z-$, $-SO_2-Z-$, $-N(R^6)-Z-$,
 $-N(R^6)SO_2-Z-$, $-SO_2N(R^6)-Z-$, $-CH=CH-Z-$, $-C\equiv C-Z-$, $-N(R^6)CO-Z-$,
 $-CON(R^6)-Z-$, $-C(O)N(R^6)S(O)_2-Z-$, $-S(O)_2N(R^6)C(O)-Z-$, $-C(O)-Z-$, $-Z-$,
 $-C(O)-Z-O-Z-$, $-N(R^6)-C(O)-Z-O-Z-$, $-O-Z-N(R^6)-Z-$, $-O-C(O)-Z-O-Z-$ or a direct
 20 bond except where Z is $C_{1-6}alkyl$;

each Z is independently a direct bond, $C_{2-6}alkenylene$ or a group of the formula



each Y is independently selected from $aryl-Z^1-$, $heterocyclyl-Z^1-$, $C_{3-7}cycloalkyl-Z^1-$,

$C_{1-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, $-(CH_2)_{1-4}CH_2F$ or $-CH(OH)CH_2F$; wherein

- 25 each Y is independently optionally substituted by up to 3 R^4 groups;

each R^4 is independently selected from halo, $-CH_2F$, CN, NH_2 , $C_{1-4}alkyl$,

$-OC_{1-6}alkyl$, $-COOH$, $-C(O)OC_{1-6}alkyl$, OH or phenyl optionally substituted by $C_{1-6}alkyl$ or $-C(O)OC_{1-6}alkyl$,

- 9 -

or R^5-X^1 , where X^1 is independently as defined in X above and R^5 is selected from hydrogen, C_{1-6} alkyl, $-CH_2-aF_a$, phenyl, naphthyl, heterocyclyl or C_{3-7} cycloalkyl; and R^5 is optionally substituted by one or more substituents independently selected from: halo, C_{1-6} alkyl, $-OC_{1-6}$ alkyl, $-CH_2-aF_a$, CN, OH, NH_2 , COOH, or $-C(O)OC_{1-6}$ alkyl,

5 each Z^1 is independently a direct bond, C_{2-6} alkenylene or a group of the formula $-(CH_2)_p-C(R^{6a})_2-(CH_2)_q$;

R^3 is heterocyclyl, wherein the atom at the two position of the heterocyclyl ring relative to the amide group, to which R^3 is attached, is a heteroatom and when the atom at the two position of the heterocyclyl ring relative to the amide group is nitrogen, this is an SP^2 hybridised nitrogen, and R^3 is optionally substituted by up to 2 R^7 groups;

10

R^6 is independently selected from hydrogen, C_{1-6} alkyl or $-C_{2-4}$ alkyl-O- C_{1-4} alkyl;

R^{6a} is independently selected from hydrogen, halo, C_{1-6} alkyl or $-C_{2-4}$ alkyl-O- C_{1-4} alkyl;

each R^7 is independently selected from:

15 C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, $(CH_2)_{0-3}$ aryl, $(CH_2)_{0-3}$ heterocyclyl, $(CH_2)_{0-3}C_{3-7}$ cycloalkyl, OH, C_{1-6} alkyl-OH, halo, C_{1-6} alkyl-halo, OC_{1-6} alkyl, $(CH_2)_{0-3}S(O)_{0-2}R^8$, SH, SO_3 , thioxo, NH_2 , CN, $(CH_2)_{0-3}NHSO_2R^8$, $(CH_2)_{0-3}COOH$, $(CH_2)_{0-3}-O-(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)OR^8$, $(CH_2)_{0-3}C(O)NH_2$, $(CH_2)_{0-3}C(O)NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NHC(O)(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)NHSO_2-R^8$ and $(CH_2)_{0-3}SO_2NHC(O)-R^8$ wherein an alkyl chain, cycloalkyl ring or heterocyclyl ring within R^7 is optionally substituted by one of more substituents independently selected from: C_{1-4} alkyl, OH, halo, CN, NH_2 , N- C_{1-4} alkylamino, N,N-di- C_{1-4} alkylamino and OC_{1-4} alkyl;

20

25 R^8 is selected from hydrogen, C_{1-6} alkyl, aryl, heterocyclyl, C_{3-7} cycloalkyl, OH, C_{1-6} alkyl-OH, COOH, $C(O)OC_{1-6}$ alkyl, $N(R^6)C_{1-6}$ alkyl, OC_{1-6} alkyl, C_{0-6} alkylOC(O) C_{1-6} alkyl, $C(OH)(C_{1-6}alkyl)C_{1-6}alkyl$; wherein an alkyl chain or aryl, heterocyclyl or cycloalkyl ring within R^8 is optionally substituted by one of more substituents independently selected from: C_{1-4} alkyl, OH, halo, CN, NH_2 , $-NH-C_{1-4}$ alkyl, $-N-di-(C_{1-4}alkyl)$ and $OC_{1-4}alkyl$;

30

each a is independently 1, 2 or 3;

p is an integer between 0 and 3;

q is an integer between 0 and 3;

and $p + q < 4$.


provided that


- (i) when R^3 is 2-pyridyl and X is other than -Z-, -C(O)-Z-O-Z-, -N((R^6))-C(O)-Z-O-Z- or
 5 -O-Z-N(R^6)-Z-, then R^3 cannot be mono-substituted at the 5-position with an R^7 group
 selected from COOH or C(O)OC₁₋₆alkyl;
- (ii) positions 3,5 on the phenyl ring (to which R^1 and R^2 are attached) relative to the amide
 bond are substituted and at least one of the groups at position 3 and 5 is an R^2 group;
- (iii) an unbranched, unsubstituted C₁₋₆alkyl chain cannot exceed C₆alkyl in length;
- 10 (iv) when n is 2 or 3 then only one X group can be -- NHC(O)-;
- (v) when R^3 is pyridyl and R^7 is halo or methyl then the phenyl ring to which R^2 is attached
 cannot be substituted by an R^2 group at the 2-position relative to the amide bond wherein
 X is -C(O)NH- and Y is optionally substituted phenyl, optionally substituted thienyl or
 optionally substituted pyridyl;
- 15 (vi) when n+m is 2, m is 0 or m is 1 and R^1 is OH, n is 1 and X is -NHC(O)- or n is 2 and X
 is independently selected from -C(O)NH-, -NHC(O)-, -O-, -S(O₂)NH- or a direct bond
 wherein one X group is -NHC(O)-, Y is selected from phenyl, cyclohexyl,
 4,5-dihydro-5-oxo-pyrazolyl, thienyl, 1,3-dihydro-1,3-dioxo-isoindolyl,
 2-oxo-1-benzopyran or pyridyl and Y is optionally substituted by R^4 then R^3 cannot be
 20 unsubstituted thiazole, 4,5-dihydro-5-oxo-pyrazolyl substituted by trichlorophenyl,
 4,5,6,7-tetrahydro-benzo[b]thiophene substituted by ethoxycarbonyl or pyridyl optionally
 independently mono or di-substituted by methyl, ethoxy or propylcarbonylamino; and
- (vii) when n+m is 3, m is 0 or 2, R^1 is independently selected from methyl, methoxy or
 hydroxy, n is 1, 2 or 3, X is independently selected from -O-, -S(O₂)NH-, -C(O)-,
 25 -S(O₂)-, -CH₂- or a direct bond, Y is selected from pyrrolidinyl, morpholino, phenyl,
 tetrazolyl or propyl wherein Y is optionally substituted by R^4 and R^4 is selected from
 di-hydroxy, methoxy, C₁₋₄alkyl then R^3 cannot be unsubstituted tetrazolyl, unsubstituted
 thiazolyl or thiazolyl substituted by ethoxycarbonylmethyl.

For the avoidance of doubt C₆alkyl is -CH₂-CH₂-CH₂-CH₂-CH₂-CH₃.


- 11 -

For the avoidance of doubt examples of R^3 wherein R^3 is heterocyclyl and the atom at the two position of the R^3 heterocyclyl ring, relative to the amide group to which R^3 is attached, is an sp^2 hybridised nitrogen include:



5 wherein  represents the point of attachment to the amide group.

According to a further feature of the invention there is provided a compound of Formula (Ic) or a salt thereof;



Formula (Ic)

10 wherein

m is 0, 1 or 2;

n is 0, 1, 2, 3 or 4;

and $n + m > 0$;

each R^1 is independently selected from OH, $(CH_2)_{1-4}OH$, CH_3-F_a , $(CH_2)_{1-4}CH_3-F_a$,

15 OCH_3-F_a , halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, NH_2 , $N(C_{1-6}alkyl)_{2-6}alkyl$, CN, phenyl or a heterocyclyl optionally substituted by $C_{1-6}alkyl$;

each R^2 is the group $Y-X-$

wherein each X is a linker independently selected from

-O $(CH_2)_{0-3}-$, -(CH₂)₀₋₃O-, -C(O)O(CH₂)₀₋₃-, -S(CH₂)₀₋₃-, -SO(CH₂)₀₋₃-,
 20 -O₂(CH₂)₀₋₃-, -NH₂SO₂-, -SO₂NH-, -N(CH₂)₀₋₃-, -N(CH₂)₁₋₃O(CH₂)₀₋₃-, -(CH₂)₁₋₄-,
 -CH=CH(CH₂)₀₋₂-, -C≡C(CH₂)₀₋₂-, -NHCO-, -CONH-;

each Y is independently selected from phenyl(CH₂)₀₋₂, naphthyl(CH₂)₀₋₂, heterocyclyl(CH₂)₀₋₂, C₃₋₇ cycloalkyl(CH₂)₀₋₂, C₂₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, or CH(OH)CH_{3-a}F_a;

each Y is independently optionally substituted by one or more R⁴ groups;

- 5 each R⁴ is independently selected from halo, CH_{3-a}F_a, OCH_{3-a}F_a, CN, NH₂, C₁₋₆alkyl, OC₁₋₆alkyl, COOH, (CH₂)₀₋₃COOH, O(CH₂)₀₋₃COOH, C(O)OC₁₋₆alkyl, C₁₋₆alkylC(O)OC₁₋₆alkyl, CO-phenyl, CONH₂, CONH-phenyl, SO₂NH₂, SO₂C₁₋₆alkyl, OH, or phenyl optionally substituted by one or more R⁵ groups, or R^{6b}-X-;

- 10 R⁵ is selected from hydrogen, C₁₋₆alkyl or C(O)OC₁₋₆alkyl, R^{6b} is selected from hydrogen, C₁₋₆alkyl, CH_{3-a}F_a phenyl, naphthyl, heterocyclyl or C₃₋₇cycloalkyl; and R^{6b} is optionally substituted by halo, C₁₋₆alkyl, CH_{3-a}F_a, CN, NH₂, COOH and COOC₁₋₆alkyl;

each a is independently 1, 2 or 3;

- 15 R³ is a heterocyclyl, and R³ is optionally substituted by one or more R⁷ groups; each R⁷ is independently selected from:


- C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, heterocyclyl, (CH₂)₀₋₃C₃₋₇cycloalkyl, OH, C₁₋₆alkyl-OH, halo, C₁₋₆alkyl-halo, OC₁₋₆alkyl, SC₁₋₆alkyl, SH, SO₃, NH₂, CN, NHCHO, NSO₂C₁₋₆alkyl, (CH₂)₀₋₃COOH, (CH₂)₀₋₃C(O)OC₁₋₆alkyl, 20 (CH₂)₀₋₃CONH₂, (CH₂)₀₋₃CON(CH₂)₀₋₃R⁸, (CH₂)₀₋₃NH(CH₂)₀₋₃R⁸, (CH₂)₀₋₃NHC(O)(CH₂)₀₋₃R⁸;
- R⁸ is selected from hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, OH, C₁₋₆alkyl-OH, COOH, C(O)OC₁₋₆alkyl, N(C₀₋₆ alkyl)C₁₋₆ alkyl, O(C₀₋₆ alkyl)C₁₋₆ alkyl, C₀₋₆alkylOC(O)C₁₋₆alkyl, C(OH)(C₁₋₆alkyl)C₁₋₆alkyl;

25 provided that

- (i) when R³ is thiazole and R⁷ is nitro, then at least one R² group is other than -O-propene;
- (ii) when R³ is pyrimidine or pyridine, then R¹ is other than OH;
- (iii) when R³ is pyridine, then R⁷ is other than COOH or COOC₁₋₆alkyl.

A further feature of the invention is a compound of Formula (Id) or a salt, solvate of 30 pro-drug thereof;

- 13 -



Formula (Id)

wherein

R^3 is phenyl, optionally substituted by one or more R^7 groups;

5 m , n , R^1 , R^2 , X , Y , R^4 , R^5 , R^6 , R^7 , R^8 , and a are as defined above for a Compound of Formula (I).

Compounds of Formula (I), (Ia), (Ib), (Ic), or (Id) may form salts which are within the ambit of the invention. Pharmaceutically acceptable salts are preferred although other salts may be useful in, for example, isolating or purifying compounds.

10 When m is 2, each R^1 group may be the same or different; preferably both R^1 groups are the same. When n is 2, 3 or 4, each R^2 group may be the same or different to any other R^2 group; preferably at least two R^2 groups are different. The R^1 and/or R^2 group(s) may be attached at the -2, -3, -4, -5 or -6 positions.


The term "aryl" refers to phenyl, naphthyl or a partially saturated bicyclic carbocyclic ring containing between 8 and 12 carbon atoms, preferably between 8 and 10 carbon atoms. Example of partially saturated bicyclic carbocyclic ring include: 1,2,3,4-tetrahydronaphthyl, indanyl, indenyl, 1,2,4a,5,8,8a-hexahydronaphthyl or 1,3a-dihydropentalene.


The term "halo" includes chloro, bromo, fluoro and iodo; preferably chloro, bromo and fluoro; most preferably fluoro.

The expression " $-\text{CH}_3\text{-}_a\text{F}_a$ " wherein a is an integer between 1 and 3 refers to a methyl group in which 1, 2 or all 3 hydrogen are replaced by a fluorine atom. Examples include: trifluoromethyl, difluoromethyl and fluoromethyl. An analogous notation is used with reference to the group $-(\text{CH}_2)_{1-4}\text{CH}_3\text{-}_a\text{F}_a$, examples include: 2,2-difluoroethyl and 3,3,3-trifluoropropyl.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups. For example, " C_{1-4} alkyl" includes propyl, isopropyl and *t*-butyl. For the avoidance of doubt, an alkyl chain can be joined to the rest of the molecule at the end of the alkyl

chain or in the middle of an alkyl chain, i.e. the definition of "alkyl" includes the following structures:



wherein  represents the point of attachment to the rest of the molecule.

- 5 A "heterocyclyl" is a saturated, partially saturated or unsaturated, monocyclic or fused bicyclic ring containing 3-12 atoms of which at least one atom is chosen from nitrogen, sulphur or oxygen, wherein a $\text{-CH}_2\text{-}$ group can optionally be replaced by a -C(O)- or sulphur atoms in a heterocyclic ring may be oxidised to S(O) or S(O)_2 . A 'heterocyclyl ring may, unless otherwise specified, be carbon or nitrogen linked, unless linking via nitrogen leads to a
- 10 charged quaternary nitrogen.

Preferably a "heterocyclyl" is a saturated, partially saturated or unsaturated, monocyclic or fused bicyclic ring wherein each ring contains 5 or 6 atoms of which 1 to 3 atoms are nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon or nitrogen linked, wherein a $\text{-CH}_2\text{-}$ group can optionally be replaced by a -C(O)- or sulphur atoms in a

15 heterocyclic ring may be oxidised to S(O) or S(O)_2 groups.

Examples and suitable values of the term "heterocyclyl" are thiazolidinyl, pyrrolidinyl, pyrrolinyl, 2-pyrrolidonyl, 2,5-dioxopyrrolidinyl, 2-benzoxazolinonyl, 1,1-dioxotetrahydrothienyl, 2,4-dioxoimidazolidinyl, 2-oxo-1,3,4-(4-triazoliny), 2-oxazolidinonyl, 5,6-dihydrouracilyl, 1,3-benzodioxolyl, 1,2,4-oxadiazolyl, 2-

20 azabicyclo[2.2.1]heptyl, 4-thiazolidonyl, morpholino, 2-oxotetrahydrofuranlyl, tetrahydrofuranlyl, 2,3-dihydrobenzofuranlyl, benzothienyl, isoxazolyl, tetrahydropyranlyl, piperidyl, 1-oxo-1,3-dihydroisoindolyl, piperazinyl, thiomorpholino, 1,1-dioxothiomorpholino, tetrahydropyranlyl, 1,3-dioxolanyl, homopiperazinyl, thienyl, isoxazolyl, imidazolyl, pyrrolyl, thiazolyl, thiadiazolyl, isothiazolyl, 1,2,4-triazolyl, 1,2,3-

25 triazolyl, pyranlyl, indolyl, pyrimidyl, thiazolyl, pyrazinyl, pyridazinyl, pyridyl, 4-pyridonyl, quinolyl and 1-isoquinolonyl.


Preferably the term "heterocyclyl" refers to monocyclic heterocyclic rings with 5- or 6-membered systems, such as isoxazolyl, pyrrolidinyl, 2-pyrrolidonyl, 2,5-dioxopyrrolidinyl, morpholino, tetrahydrofuranlyl, piperidyl, piperazinyl, thiomorpholino, tetrahydropyranlyl,

thienyl, imidazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl, indolyl, thiazolyl, thiadiazolyl, pyrazinyl, pyridazinyl and pyridyl.


- Preferred examples of 5/6 and 6/6 bicyclic ring systems include benzofuranyl, benzimidazolyl, benzthiophenyl, benzthiazolyl, benzisothiazolyl, benzoxazolyl, 5 benzisoxazolyl, pyridoimidazolyl, pyrimidoimidazolyl, quinolinyl, isoquinolinyl, quinoxalinyl, quinazolinyl, phthalazinyl, cinnolinyl and naphthyridinyl.

- The term "cycloalkyl" refers to a saturated carbocyclic ring containing between 3 to 12 carbon atoms, preferably between 3 and 7 carbon atoms. Examples of C₃₋₇cycloalkyl include cycloheptyl, cyclohexyl, cyclopentyl, cyclobutyl or cyclopropyl. Preferably cyclopropyl, 10 cyclopentyl or cyclohexyl.

- Examples of C₁₋₆alkyl include methyl, ethyl, propyl, isopropyl, *sec*-butyl, *tert*-butyl and 2-ethyl-butyl; examples of C₁₋₆alkyl-OH include hydroxymethylen and hydroxyethylene; examples of C₁₋₆alkyl-halo include chloromethylene, fluoromethylene, chloroethylene and fluoroethylene; examples of C₂₋₆alkenyl include: ethenyl, 2-propenyl, 2-butenyl, or 15 2-methyl-2-butenyl; examples of C₂₋₆alkynyl include: ethynyl, 2-propynyl, 2-butylnyl, or 2-methyl-2-butylnyl, examples of -OC₁₋₆alkyl include methoxy, ethoxy, propoxy and *tert*-butoxy; examples of -C(O)OC₁₋₆alkyl include methoxycarbonyl, ethoxycarbonyl and *tert*-butyloxycarbonyl; examples of -NH-C₁₋₆alkyl include:



20. examples of -N-di-(C₁₋₆alkyl):



- For the avoidance of doubt, in the definition of linker group 'X', the right hand side of the group is attached to the phenyl ring and the left hand side is bound to 'Y'. The same orientation applies to the linker group 'X¹', thus the right hand side of 'X¹' is attached to Y 25 and the left hand side is attached to 'R⁵'.

It is to be understood that, insofar as certain of the compounds of Formula (I), (Ia), (Ib), (Ic) and (Id) defined above or compounds of Formula (II) to (IIk) defined below may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention includes in its definition any such optically active or racemic form which possesses

- 16 -

the property of stimulating GLK directly or inhibiting the GLK/GLKRP interaction. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form. It is also to be understood that certain
 5 compounds may exist in tautomeric forms and that the invention also relates to any and all tautomeric forms of the compounds of the invention which activate GLK.

Preferred compounds of Formula (I), (Ia), (Ib), (Ic), and (Id) above, and of compounds of Formula (II) to (IIk) below are those wherein any one or more of the following apply:

- 10 (1) m is 0 or 1;
 n is 1 or 2; preferably n is 2;
 most preferably m is 0 and n is 2.
- (2) The R^1 and/or R^2 group(s) are attached at the 2-position and/or the 3-position and/or the 5- position; when $n + m$ is 3, the groups are preferably at the 2-, 3- and 5-
 15 positions; when $n + m$ is 2, the groups are preferably at the 2- and 5- or the 3- and 5- positions; most preferably there are two groups in total, substituted at the 3- and 5- positions.
- (3) each R^1 is independently selected from OH, formyl, CH_3-aF_a (preferably CF_3), OCH_3-aF_a , halo, C_{1-6} alkyl, NH_2 , CN, $(CH_2)_{1-4}OH$ or a heterocyclyl optionally substituted by
 20 C_{1-6} alkyl;
 Preferably R^1 is selected from:
 OH, formyl, CH_3-aF_a (preferably CF_3), OCH_3-aF_a (preferably OCF_3), halo, C_{1-4} alkyl (preferably methyl), NH_2 , CN and $(CH_2)_{1-4}OH$;
 Most preferably R^1 is selected from:
 25 OH, formyl, NH_2 , halo (preferably chloro) or $(CH_2)_{1-4}OH$.
- (4) each R^2 is the group Y-X-
 wherein each X is independently selected from:
 $-Z-$, $-CH=CH-Z-$, $-O-Z-$, $-C(O)-Z-$, $-C(O)O-Z-$, $-OC(O)-Z-$, $-C(O)-Z-O-Z-$,
 $-O-C(O)-Z-O-Z-$, $-S-Z-$, $-SO-Z-$, $-SO_2-Z-$, $-N(R^6)-Z-$, $-N(R^6)CO-Z-$, $-CON(R^6)-Z-$,
 30 $-N(R^6)-C(O)-Z-O-Z-$, $-SO_2N(R^6)-Z-$, $-N(R^6)SO_2-Z-$ or $-O-Z-N(R^6)-Z-$;
 preferably each X is selected from:

- 17 -

-Z-, -CH=CH-Z-, -O-Z-, -C(O)-Z-, -C(O)O-Z-, -C(O)-Z-O-Z-, -O-C(O)-Z-O-Z-,
 -N(R⁶)-Z-, -N(R⁶)CO-Z-, -N(R⁶)-C(O)-Z-O-Z- or -O-Z-N(R⁶)-Z-;

further preferably each X is selected from:

5 -Z-, -CH=CH-Z-, -O-Z-, -C(O)-Z-, -C(O)O-Z-, -C(O)-Z-O-Z-, -N(R⁶)-Z-, or
 -N(R⁶)CO-Z-;

Most preferably each X is selected from:

-CH=CH-Z-, -O-Z- or -C(O)-Z-.

each Z is independently selected from:

10 a direct bond, -(CH₂)₁₋₂, or a group of the formula -(CH₂)_p-C(R^{6a})₂-(CH₂)_q-,
 wherein R^{6a} is independently selected from hydrogen and C₁₋₄alkyl;
 preferably a direct bond, -(CH₂)₁₋₂- or a group of the formula -(CH₂)_p-C(R^{6a})₂-(CH₂)_q-,
 wherein R^{6a} is independently selected from hydrogen and C₁₋₄alkyl and p and q are
 independently 0 or 1;

more preferably a direct bond, -CH₂- or -C(CH₃)₂-.

15 and each Y is independently selected from:

C₁₋₆alkyl, C₂₋₆alkenyl, aryl-Z¹-, heterocyclyl-Z¹-, C₃₋₇ cycloalkyl(CH₂)₀₋₂,
 -(CH₂)₁₋₄CH_{3-a}F_a;

preferably each Y is selected from:

20 C₁₋₆alkyl (preferably a branched chain C₂₋₆ alkyl such as isopropyl, isobutyl, etc) ,
 C₂₋₆alkenyl, phenyl-Z¹- or heterocyclyl-Z¹-,

Most preferably each Y is selected from:

-CH₃, -C₂H₅, prop-2-yl, iso-propyl, 1-methyl-propyl, 2-methyl-propyl, allyl, phenyl,
 2-ethyl-butyl, phenyl-Z¹-, cyclopropyl-Z¹-, cyclopentyl-Z¹-, morpholino-Z¹-,
 piperidiny-Z¹-, piperaziny-Z¹-, pyrrolidiny-Z¹-, tetrahydro-2H-pyranyl-Z¹-,
 25 isoxazolyl-Z¹-, oxazolyl-Z¹-, pyridyl-Z¹-, thiazolyl-Z¹-, thienyl-Z¹- or
 isoindoliny-Z¹-,

each Z¹ is independently selected from:

30 a direct bond, -(CH₂)₁₋₂, or a group of the formula -(CH₂)_p-C(R^{6a})₂-(CH₂)_q-,
 wherein R^{6a} is independently selected from hydrogen and C₁₋₄alkyl;
 preferably a direct bond, -(CH₂)₁₋₂- or a group of the formula
 -(CH₂)_p-C(R^{6a})₂-(CH₂)_q-, wherein R^{6a} is independently selected from hydrogen
 and C₁₋₂alkyl and p and q are independently 0 or 1;

- 18 -

further preferably a direct bond, $-\text{CH}_2-$, $-\text{CH}_2\text{-CH}(\text{CH}_3)-$ or $-(\text{CH}_2)_2-$;

most preferably a direct bond, $-\text{CH}_2-$ or $-(\text{CH}_2)_2-$

wherein in each of the above Y is independently optionally substituted by R^4 .

(5) each R^2 is the group Y-X-, Z within the definition of X is a direct bond and Z^1 within the
5 definition of Y is a group of the formula $-(\text{CH}_2)_p\text{-C}(\text{R}^{6a})_2\text{-(CH}_2)_q-$.

(6) each R^4 is independently selected from

halo, $\text{CH}_3\text{-}_a\text{F}_2$ (ideally CF_3), $\text{OCH}_3\text{-}_a\text{F}_2$ (ideally OCF_3), CN, C_{1-6} alkyl, OC_{1-6} alkyl,
COOH, $\text{C}(\text{O})\text{OC}_{1-6}$ alkyl, $(\text{CH}_2)_{0-3}\text{COOH}$, $\text{O}(\text{CH}_2)_{0-3}\text{COOH}$, CO-phenyl, CONH_2 ,
CONH-phenyl, SO_2NH_2 , $\text{SO}_2\text{C}_{1-6}$ alkyl, OH, or phenyl optionally substituted by one
10 or more R^5 groups where R^5 is selected from hydrogen, C_{1-6} alkyl or
 $\text{C}(\text{O})\text{OC}_{1-6}$ alkyl;

Preferably each R^4 is selected from

halo, CN, C_{1-6} alkyl, OC_{1-6} alkyl or COOH.

(7) each R^5 is selected from:

15 C_{1-6} alkyl, phenyl, heterocyclyl or C_{3-7} cycloalkyl;

Preferably each R^5 is selected from:

C_{1-6} alkyl, tetrahydrofuranyl, imidazolyl, isoxazolyl, pyrazinyl, pyrimidinyl, thienyl,
1,3-benzodioxole, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl;

Most preferably each R^5 is selected from:

20 CH_3 , C_2H_5 , prop-2-yl, tetrahydrofuranyl, imidazolyl, isoxazolyl, pyrazinyl,
pyrimidinyl, thienyl, 1,3-benzodioxolyl or cyclopentyl;

(8) each X^1 is independently selected from:

a direct bond, -Z-, $-\text{O}-\text{C}(\text{O})-\text{Z}-$, $-\text{C}(\text{O})-\text{O}-\text{Z}-$, $-\text{C}(\text{O})-\text{Z}-$, $-\text{N}(\text{R}^6)-\text{C}(\text{O})-\text{Z}-$,
 $-\text{C}(\text{O})-\text{N}(\text{R}^6)-\text{Z}-$, $-\text{S}(\text{O}_2)-\text{Z}-$, $-\text{N}(\text{R}^6)\text{SO}_2-\text{Z}-$ or $-\text{SO}_2\text{N}(\text{R}^6)-\text{Z}-$;

25 Preferably each X^1 is independently selected from:

a direct bond, -Z-, $-\text{O}-\text{C}(\text{O})-\text{Z}-$, $-\text{C}(\text{O})-\text{Z}-$, $\text{N}(\text{R}^6)-\text{C}(\text{O})-\text{Z}-$ or $-\text{S}(\text{O}_2)-\text{Z}-$;

Most preferably each X^1 is independently selected from:

a direct bond, $-\text{CH}_2-$, $-\text{O}-\text{C}(\text{O})-$, $-\text{C}(\text{O})-$, $-\text{N}(\text{CH}_3)-\text{C}(\text{O})-\text{CH}_2-$ or $-\text{S}(\text{O})_2-$;

(9) optional substituents on R^5 are independently selected from:

30 OH, CN, NH_2 , C_{1-6} alkyl, OC_{1-6} alkyl or halo;

Preferably optional substituents on R^5 are independently selected from:

OH, C_{1-6} alkyl, OC_{1-6} alkyl or halo;


- 19 -

Most preferably optional substituents on R^5 are independently selected from:

OH, CH₃, t-butyl, OCH₃, chloro or fluoro;

(10) R^3 is a heterocyclyl (preferably a nitrogen-containing heterocyclyl group), optionally substituted by one or more R^7 groups;

5 Preferably R^3 is a heterocyclyl selected from the following:



More preferably R^3 is selected from:

thiazole, thiadiazole, pyridine, pyrazine, pyridazine, pyrazole, pyrimidine, isoxazole, furan, benzothiazole, benzimidazole and benzoxazole.

10 Further preferably R^3 is selected from:

thiazole, benzothiazole, thiadiazole, pyridine, pyrazine, pyridazine, pyrazole, imidazole, pyrimidine, oxazole and indole.

Most preferably R^3 is selected from:

pyridine, thiazole or thiadiazole.

15 In a further embodiment of the invention, R^3 is selected from:

- 20 -

benzothiazole, thiazole, thiadiazole, pyridine, pyrazine, pyridazine, pyrazole, pyrimidine, isoxazole and furan.

(11) R^3 is not substituted or is substituted by one R^7 group.

(12) each R^7 is independently selected from:

- 5 OH, CN, NH_2 , SO_3 , thioxo, halo, C_{1-6} alkyl, C_{1-6} alkyl-OH, O- C_{1-6} alkyl, C_{1-6} alkyl-halo, $(CH_2)_{0-3}COOH$, $(CH_2)_{0-3}C(O)OR^8$, $(CH_2)_{0-3}NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NHC(O)(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)NH(CH_2)_{0-3}R^8$, $-(CH_2)_{0-3}S(O)_{0-2}R^8$, $-(CH_2)_{0-3}N(R^6)SO_2R^8$, $(CH_2)_{0-3}C(O)N(R^6)S(O)_2R^8$ or $(CH_2)_{0-3}$ heterocyclyl;

preferably R^7 is selected from:

- 10 OH, CN, NH_2 , SO_3 , thioxo, halo, C_{1-4} alkyl, C_{1-4} alkyl-OH, O- C_{1-4} alkyl, C_{1-4} alkyl-halo, $(CH_2)_{0-1}COOH$, $(CH_2)_{0-1}C(O)OR^8$, $(CH_2)_{0-1}NH(CH_2)_{0-2}R^8$, $(CH_2)_{0-1}NHC(O)(CH_2)_{0-2}R^8$, $(CH_2)_{0-1}C(O)NH(CH_2)_{0-2}R^8$, $-(CH_2)_{0-2}S(O)_{0-2}R^8$, $-(CH_2)_{0-1}N(R^6)SO_2R^8$, $(CH_2)_{0-1}C(O)N(R^6)S(O)_2R^8$ or $(CH_2)_{0-1}$ heterocyclyl
(preferably the heterocyclyl is selected from furanyl, morpholino, 5-oxo-oxadiazolyl or tetrazolyl);
- 15 or tetrazolyl);

further preferably R^7 is selected from:

$COOH$, $C(O)OC_{1-6}$ alkyl, $(CH_2)_{0-1}C(O)NH(CH_2)_{0-2}R^8$, $(CH_2)_{0-3}C(O)NHSO_2R^8$ or $(CH_2)_{0-3}SO_2NHC(O)-R^8$;

most preferably R^7 is selected from:

- 20 $COOH$, $C(O)OC_{1-6}$ alkyl or $(CH_2)_{0-1}C(O)NH(CH_2)_{0-2}R^8$,

(13) R^8 is selected from:

hydrogen, OH, $COOH$, C_{1-6} alkyl, O- C_{1-6} alkyl, $-C(O)-O-C_{1-6}$ alkyl, C_{0-6} alkylOC(O) C_{1-6} alkyl, $N(R^6)C_{1-6}$ alkyl, aryl, heterocyclyl or C_{3-7} cycloalkyl;

Preferably R^8 is selected from:

- 25 hydrogen, OH, $COOH$, CH_3 , isopropyl, 2-methyl-butyl, pent-3-yl, $-O-CH_3$, $-C(O)-O-C_2H_5$, $-CH_2-O-C(O)-CH_3$, $-CH_2-O-C(O)-C_2H_5$, $-C(CH_3)_2-O-C(O)-CH_3$, NH -isopropyl, NH -t-butyl, $N(CH_3)-CH_3$, phenyl, isoxazolyl, pyrazolyl, pyridyl, thienyl, cyclopropyl or cyclobutyl;

(14) Preferred optional substituents on R^8 are independently selected from:

- 30 OH, CN, NH_2 , halo or C_{1-6} alkyl;

More preferred optional substituents on R^8 are independently selected from:

OH, halo or C_{1-6} alkyl;

- 21 -

More preferred optional substituents on R^8 are independently selected from:

OH, chloro, fluoro and CH_3 .

For example, particularly preferred compounds of the invention are those wherein:

m is 0 and n is 2, the two R^2 groups are attached at the 2- and 5- or the 3- and 5- positions (ideally the 3- and 5- positions), and X is $-O(CH_2)_{0-2}-$ (ideally $-OCH_2-$); or

m is 0 and n is 2, the two R^2 groups are attached at the 2- and 5- or the 3- and 5- positions (ideally the 3- and 5- positions), X is $-O(CH_2)_{0-2}-$ (ideally $-O-$ or $-OCH_2-$), and Y is benzyl optionally substituted by halo (such as fluoro or chloro, ideally fluoro) or C_{1-6} alkyl; or


m is 0 and n is 2, the two R^2 groups are attached at the 2- and 5- or the 3- and 5- positions (ideally the 3- and 5- positions), X is $-O(CH_2)_{0-2}-$ (ideally $-O-$ or $-OCH_2-$), and R^3 is a heterocyclyl optionally substituted by R^7 ; or

m is 0 and n is 2, the two R^2 groups are attached at the 2- and 5- or the 3- and 5- positions (ideally the 3- and 5- positions), X is $-O-$ or $-O(CH_2)_{0-2}-$ (ideally $-O-$ or $-OCH_2-$), Y is phenyl optionally substituted by halo (such as fluoro or chloro, ideally fluoro) or C_{1-6} alkyl, and R^3 is a heterocyclyl optionally substituted by R^7 ; or

m is 1 and n is 1, the R^1 and R^2 groups are attached at the 2- and 5- or the 3- and 5- positions (ideally the 3- and 5- positions), R^1 is halo (such as fluoro, chloro), and X is $-O(CH_2)_{0-2}-$ (ideally $-O-$ or $-OCH_2-$).

According to a further feature of the invention there is provided the following preferred groups of compounds of the invention:

(I) a compound of Formula (II)




Formula (II)

wherein:

X , Z^1 , R^3 and R^4 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

(II) a compound of Formula (IIa)

- 22 -



Formula (IIa)


wherein:

Het is a monocyclic heterocyclyl, optionally substituted with up to 3 groups selected from R^4 and,

X , Z^1 , R^3 and R^4 are as defined above in a compound of Formula (I);

or a salt, solvate or pro-drug thereof.

(III) a compound of Formula (IIb)



Formula (IIb)

wherein:


the $C_{1-6}alkyl$ group is optionally substituted with up to 3 groups selected from R^4 , preferably unsubstituted;

the $C_{1-6}alkyl$ group optionally contains a double bond, preferably the $C_{1-6}alkyl$ group does not contain a double bond; and

X , Z^1 , R^3 and R^4 are as defined above in a compound of Formula (I);

or a salt, solvate or pro-drug thereof.

(IV) a compound of Formula (IIc)



Formula (IIc)

wherein:


the $C_{3-7}cycloalkyl$ group is optionally substituted with up to 3 groups selected from R^4 ,

and

- 23 -

X , Z^1 , R^3 and R^4 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

(V) a compound of Formula (IIc)



Formula (IIc)


wherein:

the C_{1-6} alkyl groups are independently optionally substituted with up to 3 groups selected from R^4 , preferably one of the C_{1-6} alkyl groups is unsubstituted,

the C_{1-6} alkyl groups independently optionally contain a double bond, preferably only one of the C_{1-6} alkyl groups contain a double bond, preferably neither of the C_{1-6} alkyl group contains a double bond, and

X , R^3 and R^4 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

(VI) a compound of Formula (IId)



Formula (IId)

wherein:


the C_{3-7} cycloalkyl and C_{1-6} alkyl groups are independently optionally substituted with up to 3 groups selected from R^4 , preferably the C_{1-6} alkyl group is unsubstituted;

the C_{1-6} alkyl group optionally contains a double bond, preferably the C_{1-6} alkyl group does not contain a double bond; and

X , Z^1 , R^3 and R^4 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

(VII) a compound of Formula (IIe)

- 24 -



Formula (IIf)

wherein:

Het is a monocyclic heterocyclyl,

- 5 the Het and C₁₋₆alkyl groups are independently optionally substituted with up to 3 groups selected from R⁴, preferably the C₁₋₆alkyl group is unsubstituted;

the C₁₋₆alkyl group optionally contains a double bond, preferably the C₁₋₆alkyl group does not contain a double bond; and

X, Z¹, R³ and R⁴ are as defined above in a compound of Formula (I);

- 10 or a salt, solvate or pro-drug thereof.

A further preferred group of compounds of group (VII) comprise compounds of Formula (IIf) wherein:

Het is a saturated monocyclic heterocyclyl;


X is -Z-, preferably -CH₂-;

- 15 R⁴ is a group of R⁵-X¹-;

X¹ is as defined for a compound of Formula (I);

R⁵ is C₁₋₆alkyl, phenyl, heterocyclyl, each of which is optionally substituted as defined for a compound of Formula (I);

(VIII) a compound of Formula (IIg)



20

Formula (IIg)

wherein:

Het is a monocyclic heterocyclyl,


the Het and C₃₋₇cycloalkyl groups are independently optionally substituted with up to 3 groups selected from R⁴, and

25

X, Z¹, R³ and R⁴ are as defined above in a compound of Formula (I);

or a salt, solvate or pro-drug thereof.

(IX) a compound of Formula (IIIh)




Formula (IIIh)

wherein:

- 5 Y is aryl- Z^1 -, wherein aryl is preferably a partially saturated bicyclic carbocyclic ring;
 Y and the C_{1-6} alkyl group are independently optionally substituted with up to 3 groups
 selected from R^4 , preferably the C_{1-6} alkyl group is unsubstituted,
 the C_{1-6} alkyl group optionally contains a double bond, preferably the C_{1-6} alkyl group does
 not contain a double bond; and
- 10 X, Z^1 , R^3 and R^4 are as defined above in a compound of Formula (I);
 or a salt, solvate or pro-drug thereof.

(X) a compound of Formula (IIj)




Formula (IIj)

- 15 wherein:
- X is selected from $-SO_2N(R^6)-Z-$ or $-N(R^6)SO_2-Z-$, preferably X is $-SO_2N(R^6)-Z-$;
 Z is as described above, preferably Z is propylene, ethylene or methylene, more
 preferably Z is methylene;
- Z^a is selected from a direct bond or a group of the formula $-(CH_2)_p-C(R^{6a})_2-(CH_2)_q-$;
 preferably Z^a is selected from C_{1-2} alkylene or a direct bond; preferably Z^a is a direct
 bond;
- R^{6a} is selected from: C_{1-4} alkyl or hydrogen, preferably methyl or hydrogen;
- Y is selected from aryl- Z^1 - or heterocyclyl- Z^1 -;
- Y and the C_{1-6} alkyl group are independently optionally substituted with up to 3 groups
- 25 selected from R^4 ,
 the C_{1-6} alkyl group optionally contains a double bond, preferably the C_{1-6} alkyl group does
 not contain a double bond, and

- 26 -

Z^1 , R^3 and R^4 are as defined above in a compound of Formula (I);
or a salt, solvate or pro-drug thereof.

(XI) a compound of Formula (IIIk)



Formula (IIIk)

wherein:

the C₃₋₇cycloalkyl groups are independently optionally substituted with up to 3 groups selected from R^4 , and

X , Z^1 , R^3 and R^4 are as defined above in a compound of Formula (I);


or a salt, solvate or pro-drug thereof.

A further preferred groups of compounds of the invention in either of groups (I)-(XI) above is wherein:

X is independently selected from: -O-Z-, SO₂N(R⁶)-Z- or -N(R⁶)-Z-;

Z is a direct bond or -CH₂-;

Z^1 is selected from a direct bond, -CH₂-, -(CH₂)₂- or



R^3 is as defined above in a compound of Formula (I);

or a salt, solvate or pro-drug thereof.

A further preferred groups of compounds of the invention in either of groups (I)-(XI)

above is wherein:

R^3 is substituted by at least one R^7 group (preferably one R^7 group);

R^7 is a group of the formula (CH₂)₀₋₃NH(CH₂)₀₋₃R⁸, (CH₂)₀₋₃N(R⁶)S(O)₂R⁸ or

(CH₂)₀₋₃heterocyclyl (preferably 5-oxo-1,2,4-oxadiazol-3-yl or -tetrazol-5-yl);

R^3 , R^6 and R^8 are as defined above in a compound of Formula (I);

or a salt, solvate or pro-drug thereof.

The compounds of the invention may be administered in the form of a pro-drug. A pro-drug is a bioprecursor or pharmaceutically acceptable compound being degradable in the body to produce a compound of the invention (such as an ester or amide of a

compound of the invention, particularly an *in vivo* hydrolysable ester). Various forms of prodrugs are known in the art. For examples of such prodrug derivatives, see:

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
- 5 b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen;
- c) H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard p. 113-191 (1991);
- d) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- e) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77, 285 (1988); and
- 10 f) N. Kakeya, *et al.*, Chem Pharm Bull, 32, 692 (1984).

The contents of the above cited documents are incorporated herein by reference.

Examples of pro-drugs are as follows. An in-vivo hydrolysable ester of a compound of the invention containing a carboxy or a hydroxy group is, for example, a pharmaceutically-acceptable ester which is hydrolysed in the human or animal body to produce the parent acid
 15 or alcohol. Suitable pharmaceutically-acceptable esters for carboxy include C₁ to C₆alkoxymethyl esters for example methoxymethyl, C₁ to ₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃ to ₈cycloalkoxycarbonyloxyC₁ to ₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2 onylmethyl esters, for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁-₆alkoxycarbonyloxyethyl esters.

- 20 An in-vivo hydrolysable ester of a compound of the invention containing a hydroxy group includes inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and α -acyloxyalkyl ethers and related compounds which as a result of the in-vivo hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy.
- 25 A selection of in-vivo hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and N-(dialkylaminoethyl)-N-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl.

A suitable pharmaceutically-acceptable salt of a compound of the invention is, for
 30 example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In

addition a suitable pharmaceutically-acceptable salt of a benzoxazinone derivative of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

A further feature of the invention is a pharmaceutical composition comprising a compound of Formula (I) to (Id) or (II) to (IIIk) as defined above, or a salt, solvate or prodrug thereof, together with a pharmaceutically-acceptable diluent or carrier.

10 According to another aspect of the invention there is provided a compound of Formula (Ib) to (Id), or (II) to (IIIk) as defined above for use as a medicament; provided that when R^3 is 2-pyridyl and X is other than -Z-, -C(O)-Z-O-Z-, -N(R^6)-C(O)-Z-O-Z- or -O-Z-N(R^6)-Z-, then R^3 cannot be mono-substituted at the 4-position with an R^7 group selected from COOH or C(O)OC₁₋₆alkyl.

15 Further according to the invention there is provided a compound of Formula (Ib) to (Id), or (II) to (IIIk) for use in the preparation of a medicament for treatment of a disease mediated through GLK, in particular type 2 diabetes.

The compound is suitably formulated as a pharmaceutical composition for use in this way.

20 According to another aspect of the present invention there is provided a method of treating GLK mediated diseases, especially diabetes, by administering an effective amount of a compound of Formula (Ib) to (Id), or (II) to (IIIk), or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

Specific disease which may be treated by the compound or composition of the invention include: blood glucose lowering in Diabetes Mellitus type 2 without a serious risk of hypoglycaemia (and potential to treat type 1), dyslipidemia, obesity, insulin resistance, metabolic syndrome X, impaired glucose tolerance.

As discussed above, thus the GLK/GLKRP system can be described as a potential "Diabetes" target (of benefit in both Diabetes and Obesity). Thus, according to another aspect of the invention there is provided the use of a compound of Formula (Ib) to (Id), or (II) to (IIIk), or salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the combined treatment or prevention of diabetes and obesity.

According to another aspect of the invention there is provided the use of a compound of Formula (Ib) to (Id), or (II) to (IIk), or salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the treatment or prevention of obesity.

According to a further aspect of the invention there is provided a method for the combined treatment of obesity and diabetes by administering an effective amount of a compound of Formula (Ib) to (Id), or (II) to (IIk), or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

According to a further aspect of the invention there is provided a method for the treatment of obesity by administering an effective amount of a compound of Formula (Ib) to (Id), or (II) to (IIk), or salt, solvate or pro-drug thereof, to a mammal in need of such treatment.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or alginic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl p-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal tract, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

- 5 Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxethylene stearate), or
10 condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example
15 heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and
20 hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl *p*-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring
25 agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting
30 agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of

oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

For further information on formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in

Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula (I), (Ia), (Ib), (Ic) or (Id) will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine.

In using a compound of the Formula (I), (Ia), (Ib), (Ic) or (Id) for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.5 mg to 75 mg per kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.5 mg to 30 mg per kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.5 mg to 25 mg per kg body weight will be used. Oral administration is however preferred.

The elevation of GLK activity described herein may be applied as a sole therapy or may involve, in addition to the subject of the present invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. Simultaneous treatment may be in a single tablet or in separate tablets. For example in the treatment of diabetes mellitus chemotherapy may include the following main categories of treatment:

- 1) Insulin and insulin analogues;
- 2) Insulin secretagogues including sulphonylureas (for example glibenclamide, glipizide) and prandial glucose regulators (for example repaglinide, nateglinide);
- 3) Insulin sensitising agents including PPAR γ agonists (for example pioglitazone and rosiglitazone);
- 4) Agents that suppress hepatic glucose output (for example metformin).
- 5) Agents designed to reduce the absorption of glucose from the intestine (for example acarbose);
- 6) Agents designed to treat the complications of prolonged hyperglycaemia;
- 7) Anti-obesity agents (for example sibutramine and orlistat);
- 8) Anti-dyslipidaemia agents such as, HMG-CoA reductase inhibitors (statins, eg pravastatin); PPAR α agonists (fibrates, eg gemfibrozil); bile acid sequestrants

(cholestyramine); cholesterol absorption inhibitors (plant stanols, synthetic inhibitors); bile acid absorption inhibitors (IBATi) and nicotinic acid and analogues (niacin and slow release formulations);

- 9) Antihypertensive agents such as, β blockers (eg atenolol, inderal); ACE inhibitors (eg
5 lisinopril); Calcium antagonists (eg. nifedipine); Angiotensin receptor antagonists (eg
candesartan), α antagonists and diuretic agents (eg. furosemide, benzthiazide);
- 10) Haemostasis modulators such as, antithrombotics, activators of fibrinolysis and
antiplatelet agents; thrombin antagonists; factor Xa inhibitors; factor VIIa inhibitors;
antiplatelet agents (eg. aspirin, clopidogrel); anticoagulants (heparin and Low
10 molecular weight analogues, hirudin) and warfarin; and
- 11) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (eg. aspirin)
and steroidal anti-inflammatory agents (eg. cortisone).

According to another aspect of the present invention there is provided individual
compounds produced as end products in the Examples set out below and salts, solvates and
15 pro-drugs thereof.


A compound of the invention, or a salt thereof, may be prepared by any process known
to be applicable to the preparation of such compounds or structurally related compounds.
Such processes are illustrated by the following representative schemes (Routes 1 - 16) in
which variable groups have any of the meanings defined for formula (I) unless stated
20 otherwise. Functional groups may be protected and deprotected using conventional methods.
For examples of protecting groups such as amino and carboxylic acid protecting groups (as
well as means of formation and eventual deprotection), see T.W. Greene and P.G.M. Wuts,
"Protective Groups in Organic Synthesis", Second Edition, John Wiley & Sons, New York,
1991.

25 The condensation of an acid with a heterocyclic amine (Route 1) is used in the
preparation of compounds of the invention or in the preparation of intermediates to the final
products. One or more further reactions (such as ester hydrolysis, Routes 2a and 2b) may then
be performed on these intermediates. The amide-forming reaction (Route 1) is best
accomplished via the acid chloride, which is usually prepared using oxalyl chloride. However,
30 alternative methods for acid chloride formation (such as resin-bound triphenyl phosphine with
carbon tetrachloride and dichloromethane) may also be employed. Additionally, alternative

methods of amide-bond formation (such as a peptide coupling agent such as EDC or HATU, with or without additives such as DIPEA or DMAP) may be used.

The remaining preparative routes (Routes 2 - 18) consist of further manipulation of the compound with the amide bond in place. Further preparative routes are summarise in Routes 5 19 – 29. Examples of routes 1-29 are provided in the examples below. Reagents and conditions given are only for illustration and alternative methods may generally be employed.


Route 1




Other amide forming reactions include:



- 1a: Oxalyl chloride in the presence of a suitable solvent or base;
1b: coupling reagents such as HATU or EDAC in the presence of a suitable solvent or base; and
15 1c: POC13/Pyridine, according to Dirk T.S. Rijkers, Hans P.H.M. Adams, H. Coenraad Hemker, Godefridus I. Tesser; Tetrahedron, 1995, 51(41), pp11235-11250.

Route 2a and 2b




Route 3






Route 4**5 Route 5****Route 6**

10

**Route 7**

15




Route 7b:**5 Route 7c:****Route 8**

10


**Route 9**



15

**Route 10**


20




Route 11

Route 12





5

Route 13


10

Route 14


15

Route 15**5 Route 16****Route 17****Route 18**


15

Route 19:

Route 20:



5 Route 21:




Route 22:


10




Route 23:

15






Route 24:**Route 25:****Route 26:**

10


**Route 27:**

15



Route 28:**5 Route 29:**

- Processes for the synthesis of compounds of Formula (I) are provided as a further
 10 feature of the invention. Thus, according to a further aspect of the invention there is provided
 a process for the preparation of a compound of Formula (I) which comprises:
 (a) reaction of a compound of Formula (IIIa) with a compound of Formula (IIIb),




Formula (IIIa)



Formula (IIIb); or


- 15 wherein X¹ is a leaving group
 (b) for compounds of Formula (I) wherein R³ is substituted with -(CH₂)₀₋₃COOH,
 de-protection of a compound of Formula (IIIc),




Formula (IIIc)

- 20 wherein P¹ is a protecting group;

- (c) for compounds of Formula (I) wherein n is 1, 2, 3 or 4, reaction of a compound of Formula (III_d) with a compound of Formula (III_e),

Formula (III_d)Formula (III_e)

- 5 wherein X' and X'' comprises groups which when reacted together form the group X ;
 (d) for a compound of Formula (I) wherein n is 1, 2, 3 or 4 and X or X^1 is $-SO-Z-$ or $-SO_2-Z-$,
 oxidation of the corresponding compound of Formula (I) wherein X or X^1 respectively is $-S-Z-$;
 (e) reaction of a compound of Formula (III_f) with a compound of Formula (III_g),

Formula (III_f)Formula (III_g); or

10

wherein X^2 is a leaving group

and thereafter, if necessary:

- i) converting a compound of Formula (I) into another compound of Formula (I);
 15 ii) removing any protecting groups;
 iii) forming a salt, pro-drug or solvate thereof.

Specific reaction conditions for the above reactions are as follows:

Process a) – as described above for Route 1);

Process b) – as described above for Route 2);

- 20 *Process c)* – examples of this process are as follows:

- (i) to form a group when X is $-O-Z-$, X' is a group of formula $HO-Z-$ and X'' is a leaving group (alternatively X' is a group of formula L^2-Z- wherein L^2 is a leaving group and X'' is a hydroxyl group), compounds of Formula (III_d) and (III_e) are reacted together in a suitable solvent, such as DMF or THF, with a base such as sodium hydride or potassium
 25 *tert*-butoxide, at a temperature in the range 0 to 100°C, optionally using metal catalysis such as palladium on carbon or cuprous iodide;

- (ii) to form a group when X is $\text{N(R}^6\text{)-Z-}$, X' is a group of formula $\text{H-(R}^6\text{)N-Z-}$ and X'' is a leaving group (alternatively X' is a group of formula $\text{L}^2\text{-Z-}$ wherein L^2 is a leaving group and X'' is a group or formula $\text{-N(R}^6\text{)-H}$), compounds of Formula (III d) and (III e) are reacted together in a suitable solvent such as THF, an alcohol or acetonitrile, using a reducing agent such as sodium cyano borohydride or sodium trisacetoxyborohydride at room temperature;
- (iii) to form a group when X is $\text{-SO}_2\text{N(R}^6\text{)-Z-}$, X' is a group of formula $\text{H-N(R}^6\text{)-Z-}$ wherein L^2 is a leaving group and X'' is an activated sulphonyl group such as a group of formula $\text{-SO}_2\text{-Cl}$, compounds of Formula (III d) and (III e) are reacted together in a suitable solvent such as methylene chloride, THF or pyridine, in the presence of a base such as triethylamine or pyridine at room temperature;
- (iv) to form a group when X is $\text{-N(R}^6\text{)SO}_2\text{-Z-}$, X' is an activated sulphonyl group such as a group of formula $\text{Cl-SO}_2\text{-Z-}$ group and X'' is a group of formula $\text{-N(R}^6\text{)-L}^2$ wherein L^2 is a leaving group, compounds of Formula (III d) and (III e) are reacted together in a suitable solvent such as methylene chloride, THF or pyridine, in the presence of a base such as triethylamine or pyridine at room temperature;
- (v) to form a group when X is $\text{-C(O)N(R}^6\text{)-Z-}$, X' is a group of formula $\text{H-N(R}^6\text{)-Z-}$ wherein L^2 is a leaving group and X'' is an activated carbonyl group such as a group of formula -C(O)-Cl , compounds of Formula (III d) and (III e) are reacted together in a suitable solvent such as THF or methylene chloride, in the presence of a base such as triethylamine or pyridine at room temperature;
- (vi) to form a group when X is $\text{-N(R}^6\text{)C(O)-Z-}$, X' is an activated carbonyl group such as a group of formula Cl-C(O)-Z- group and X'' is a group of formula $\text{-N(R}^6\text{)-L}^2$ wherein L^2 is a leaving group, compounds of Formula (III d) and (III e) are reacted together in a suitable solvent such as THF or methylene chloride, in the presence of a base such as triethylamine or pyridine at room temperature;
- (vii) to form a group when X is -CH=CH-Z- , a Wittig reaction or a Wadsworth-Emmans Horner reaction can be used. For example, X' terminates in an aldehyde group and Y-X'' is a phosphine derivative of the formula $\text{Y-CH-P}^+\text{PH}_3$ which can be reacted together in a strong base such as sodium hydride or potassium *tert*-butoxide, in a suitable solvent such as THF at a temperature between room temperature and 100°C.

Process d) - the oxidization of a compound of Formula (I) wherein X or X¹ is -S-Z- is well known in the art, for example, reaction with metachloroperbenzoic acid (MCPBA) is the presence of a suitable solvent such as dichloromethane at ambient temperature. If an excess of MCPBA is used a compound of Formula (I) wherein X is —S(O₂)— is obtained.

- 5 *Process e*) - reaction of a Formula (III_f) with a compound of Formula (III_g) can be performed in a polar solvent, such as DMF or a non-polar solvent such as THF with a strong base, such as sodium hydride or potassium *tert*-butoxide at a temperature between 0 and 100°C, optionally using metal catalysis, such as palladium on carbon or cuprous iodide.

During the preparation process, it may be advantageous to use a protecting group for a
10 functional group within R². Protecting groups may be removed by any convenient method as described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with minimum disturbance of groups elsewhere in the molecule.

Specific examples of protecting groups are given below for the sake of convenience, in
15 which "lower" signifies that the group to which it is applied preferably has 1-4 carbon atoms. It will be understood that these examples are not exhaustive. Where specific examples of methods for the removal of protecting groups are given below these are similarly not exhaustive. The use of protecting groups and methods of deprotection not specifically mentioned is of course within the scope of the invention.

- 20 A carboxy protecting group may be the residue of an ester-forming aliphatic or araliphatic alcohol or of an ester-forming silanol (the said alcohol or silanol preferably containing 1-20 carbon atoms). Examples of carboxy protecting groups include straight or branched chain (1-12C)alkyl groups (e.g. isopropyl, *t*-butyl); lower alkoxy lower alkyl groups (e.g. methoxymethyl, ethoxymethyl, isobutoxymethyl; lower aliphatic acyloxy lower alkyl groups, (e.g.
25 acetoxymethyl, propionyloxymethyl, butyryloxymethyl, pivaloyloxymethyl); lower alkoxycarbonyloxy lower alkyl groups (e.g. 1-methoxycarbonyloxyethyl, 1-ethoxycarbonyloxyethyl); aryl lower alkyl groups (e.g. *p*-methoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, benzhydryl and phthalidyl); tri(lower alkyl)silyl groups (e.g. trimethylsilyl and *t*-butyldimethylsilyl); tri(lower alkyl)silyl lower alkyl groups (e.g. trimethylsilylethyl); and
30 (2-6C)alkenyl groups (e.g. allyl and vinyllethyl).

Methods particularly appropriate for the removal of carboxyl protecting groups include for example acid-, metal- or enzymically-catalysed hydrolysis.

Examples of hydroxy protecting groups include lower alkenyl groups (e.g. allyl); lower alkanoyl groups (e.g. acetyl); lower alkoxycarbonyl groups (e.g. *t*-butoxycarbonyl); lower alkenyloxycarbonyl groups (e.g. allyloxycarbonyl); aryl lower alkoxycarbonyl groups (e.g. benzoyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, *o*-nitrobenzyloxycarbonyl, 5 *p*-nitrobenzyloxycarbonyl); tri lower alkyl/arylsilyl groups (e.g. trimethylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl); aryl lower alkyl groups (e.g. benzyl) groups; and triaryl lower alkyl groups (e.g. triphenylmethyl).

Examples of amino protecting groups include formyl, aralkyl groups (e.g. benzyl and substituted benzyl, e.g. *p*-methoxybenzyl, nitrobenzyl and 2,4-dimethoxybenzyl, and 10 triphenylmethyl); di-*p*-anisylmethyl and furylmethyl groups; lower alkoxycarbonyl (e.g. *t*-butoxycarbonyl); lower alkenyloxycarbonyl (e.g. allyloxycarbonyl); aryl lower alkoxycarbonyl groups (e.g. benzyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, *o*-nitrobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl); trialkylsilyl (e.g. trimethylsilyl and *t*-butyldimethylsilyl); alkylidene (e.g. methylenidene); benzylidene and substituted benzylidene groups.

15 Methods appropriate for removal of hydroxy and amino protecting groups include, for example, acid-, base, metal- or enzymically-catalysed hydrolysis, or photolytically for groups such as *o*-nitrobenzyloxycarbonyl, or with fluoride ions for silyl groups.

Examples of protecting groups for amide groups include aralkoxymethyl (e.g. benzyloxymethyl and substituted benzyloxymethyl); alkoxymethyl (e.g. methoxymethyl and 20 trimethylsilylethoxymethyl); tri alkyl/arylsilyl (e.g. trimethylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl); tri alkyl/arylsilyloxymethyl (e.g. *t*-butyldimethylsilyloxymethyl, *t*-butyldiphenylsilyloxymethyl); 4-alkoxyphenyl (e.g. 4-methoxyphenyl); 2,4-di(alkoxy)phenyl (e.g. 2,4-dimethoxyphenyl); 4-alkoxybenzyl (e.g. 4-methoxybenzyl); 2,4-di(alkoxy)benzyl (e.g. 2,4-di(methoxy)benzyl); and alk-1-enyl (e.g. allyl, but-1-enyl and 25 substituted vinyl e.g. 2-phenylvinyl).

Aralkoxymethyl, groups may be introduced onto the amide group by reacting the latter group with the appropriate aralkoxymethyl chloride, and removed by catalytic hydrogenation. Alkoxymethyl, tri alkyl/arylsilyl and tri alkyl/silyloxymethyl groups may be introduced by reacting the amide with the appropriate chloride and removing with acid; or in the case of the silyl 30 containing groups, fluoride ions. The alkoxyphenyl and alkoxybenzyl groups are conveniently introduced by arylation or alkylation with an appropriate halide and removed by oxidation with ceric ammonium nitrate. Finally alk-1-enyl groups may be introduced by reacting the amide with the appropriate aldehyde and removed with acid.

The following examples are for illustration purposes and are not intended to limit the scope of this application. Each exemplified compound represents a particular and independent aspect of the invention. In the following non-limiting Examples, unless otherwise stated:

- 5 (i) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids such as drying agents by filtration;
- (ii) operations were carried out at room temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as argon or nitrogen;
- (iii) yields are given for illustration only and are not necessarily the maximum
- 10 attainable;
- (iv) the structures of the end-products of the Formula (I) were confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic resonance chemical shift values were measured on the delta scale and peak multiplicities are shown as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad; q, quartet, quin,
- 15 quintet;
- (v) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), high-performance liquid chromatography (HPLC), infra-red (IR) or NMR analysis;
- (vi) chromatography was performed on silica (Merck Silica gel 60, 0.040 - 0.063
- 20 mm, 230 - 400 mesh); and
- (vi) Biotage cartridges refer to pre-packed silica cartridges (from 40g up to 400g), eluted using a biotage pump and fraction collector system; Biotage UK Ltd, Hertford, Herts, UK.

25 Abbreviations



ADDP	azodicarbonyl)dipiperidine;
DCM	dichloromethane;
DEAD	diethyldiazocarboxylate;
DIAD	di-i-propyl azodicarboxylate;
30 DIPEA	di-isopropylethylamine
DMSO	dimethyl sulphoxide;
DMF	dimethylformamide;

- 48 -

DtAD	di-t-butyl azodicarboxylate;
EDAC/EDC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride;
HATU	O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium 5 hexafluorophosphate;
LCMS	liquid chromatography / mass spectroscopy;
MPLC	medium pressure liquid chromatography;
RT	room temperature; and
THF	tetrahydrofuran.

Generic Methods for alkylation of mono- and di-hydroxy benzoate esters:


The following generic alkylation methods are referred to in the Examples below.

**Generic Method A – synthesis of symmetrical diethers (R1 = R2)**

Compound (a)


Methyl 3,5-dihydroxybenzoate (74.1g, 0.44M) was dissolved in dimethylformamide (400ml), potassium carbonate (152g, 1.10M) added, stirred for 15mins then 2-chlorobenzylchloride (117ml, 0.92M) added and heated at 100°C under an argon atmosphere. After 3hrs the reaction mixture was cooled to ambient temperature, concentrated *in vacuo*, diluted with
 20 water (800ml), extracted with ethyl acetate (2x600ml). The organic extracts were washed with brine (300ml), dried (MgSO₄), filtered, concentrated *in vacuo* to yield a brown oil which was triturated with diethyl ether/ isohexane to give compound (a) as an off-white solid (195g, 100%); ¹H nmr (d₆-DMSO, δ values): 3.81 (3H, s); 5.18 (4H, s); 6.98 (1H, m); 7.16 (1H, d); 7.36 (4H, m); 7.50 (2H, m); 7.58 (2H, m).

25

Generic Method B – synthesis of unsymmetrical diethers (R1 \neq R2)


Compound (b)

- Methyl 3,5-dihydroxybenzoate (16.8g, 0.1mol) was dissolved in dimethylformamide (180ml), powdered potassium carbonate (27.6g, 0.2mol) added, followed by 2-iodopropane (10ml, 0.1mol), and the resulting suspension stirred overnight at ambient temperature under an argon atmosphere. The reaction mixture was diluted with water (1l) and extracted with diethyl ether (2x200ml). The organic extracts were washed sequentially with water and brine, dried (MgSO₄), filtered and concentrated *in vacuo* to yield a pale golden oil which was triturated with toluene and filtered to remove unreacted ether starting material. The filtrate was
- 10 concentrated *in vacuo* and the residue chromatographed (2x90g Biotage cartridges, eluting with isohexane containing ethyl acetate (10% v/v increasing to 15% v/v) to give methyl 3-hydroxy 5-isopropoxy benzoate as a colourless solid (5.3g, 25%); ¹H nmr (d₆-DMSO, δ values): 1.2 (6H, d); 3.8 (3H, s); 4.6 (1H, hept); 6.55 (1H, m); 6.85 (1H, m); 6.95 (1H, m); 9.8 (1H, s).
- 15 Methyl 3-hydroxy 5-isopropoxy benzoate (1.5g, 7.2mmol) was dissolved in dimethylformamide (10ml), potassium carbonate (2.5g, 18mmol) added, followed by 2-bromobutane (1.2ml, 11mmol), and the resulting suspension stirred for 7 hours at 80 deg C under an argon atmosphere. The reaction mixture was cooled to ambient temperature, diluted with hexane / ethyl acetate (1:1 v/v) and washed sequentially with water and brine, dried
- 20 (MgSO₄), filtered and concentrated *in vacuo* to yield a colourless oil which was chromatographed (flash column on silica (20g), eluting with isohexane containing ethyl acetate (5 % v/v) to give methyl 3-(2-butyloxy) 5-isopropoxy benzoate as a colourless oil (1.06g); ¹H nmr (d₆-DMSO, δ values): 0.9 (3H, t); 1.2 (3H, d + 6H, d); 1.6 (2h, m); 3.85 (3H, s); 4.4 (1H, hept); 4.55 (1H, hept); 6.7 (1H, m); 7.0 (2H, m); m/z 267 (M+H)⁺.

Generic Method C – synthesis of unsymmetrical diethers (R1 \neq R2):

Compound (c)

Methyl 3-hydroxy 5-isopropoxy benzoate (0.5g, 2.4mmol) was dissolved in dichloromethane (10ml) and cooled to 0 deg C whilst stirring under an argon atmosphere; the solution was treated sequentially with triphenyl phosphine (Polymer supported, 1.19g, 3.6mmol), furfuryl alcohol (0.23 ml, 2.7 mmol) and di-*t*-butyl azodicarboxylate (DtAD, 0.082g, 3.5 mmol) added dropwise in dichloromethane (4ml), and the resulting solution stirred for 1.5 hours. The reaction was monitored by hplc and further reagents were added until the starting phenol was consumed – total reagents added were triphenyl phosphine (Polymer supported, 2.38g, 3 eq), furfuryl alcohol (0.53 ml, 2.5 eq) and DtAD (1.64g, 3 eq). The reaction mixture was concentrated *in vacuo* and purified by chromatography (flash column on silica, eluting with isohexane containing ethyl acetate (5 % v/v) to give methyl 3-(2-furyl methoxy) 5-isopropoxy benzoate as a colourless oil, (0.225g); ¹H nmr (d₆-DMSO, δ values): 1.25 (6H, d); 3.85 (3H, s); 4.65 (1H, hept); 5.1 (2H, s); 6.45 (1H, m); 6.6 (1H, m); 6.85 (1H, m); 7.05 (1H, m); 7.15 (1H, m) 7.75 (1H, m).

Generic Method D – synthesis of unsymmetrical diethers:

Compound (d)

Di-*i*-propyl azodicarboxylate (DIAD, 0.74ml, 3.7 mM) was added to methyl (5-isopropoxy-3-methanol)-benzoate (0.56g, 2.5 mM), triphenylphosphine (0.98g, 3.7 mM) and 2-fluorophenol (0.24ml, 2.7 mM) in DCM (40ml) under argon at ambient temperature. After 10 mins concentrated, purified on silica gel (10-15%EtOAc/iso-hexane) gave the title compound as a

- 51 -

pale yellow oil, which solidified under high-vacuum (0.71g, 90%); ^1H NMR δ (d_6 -DMSO): 1.26 (d, 6H), 3.82 (s, 3H), 4.64 (m, 1H), 5.21 (s, 2H), 6.92 (m, 1H), 7.09 (m, 1H), 7.16-7.26 (m, 3H), 7.35 (s, 1H), 7.58 (s, 1H).


- 5 The above generic methods are for illustration only; it will be appreciated that alternative conditions that may optionally be used include: use of alternative solvents (such as acetone or tetrahydrofuran), alternative stoichiometries of reagents, alternative reaction temperatures and alternative methods of purification.

All analytical data (NMR and/or MS) were consistent with the proposed structures.

10

EXAMPLE A

Route 1: 2-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-thiazole




- Diisopropylethylamine (DIPEA, 0.34ml, 2.0mM) then N,N -dimethylaminopyridine (DMAP, 12mg, 0.1mM) were added to a solution of 2-aminothiazole (0.10g, 1.0mM) and 3,5-di-(2-chlorobenzoyloxy) benzoic acid chloride (0.42g, 1.0mM) in dichloromethane (10ml) under argon at ambient temperature. After 80mins the reaction mixture was filtered, washed with dichloromethane and dried under high vacuum to give the title compound as a colourless solid (0.20g, 41%); ^1H NMR δ (d_6 -DMSO): 5.24 (4H, s); 6.93 (1H, s); 7.26 (1H, d); 7.36-7.43 (6H, m); 7.50 (2H, m); 7.55 (1H, d); 7.61 (2H, m); 12.60 (1H, br s).
- 20

Alternative conditions that may optionally be used include: use of an alternative solvent, such as tetrahydrofuran; use of pyridine as solvent, with or without the addition of DMAP or DIPEA; dissolving the acid chloride component in the solvent of choice, and adding the amine component to it.

25


The requisite 3,5-Di-(2-chlorobenzoyloxy) benzoic acid chloride starting material, compound (c), was prepared as follows:

- 52 -




Compound (a)

Methyl 3,5-dihydroxybenzoate (74.1g, 0.44M) was dissolved in dimethylformamide (400ml), potassium carbonate (152g, 1.10M) added, stirred for 15mins then 2-chlorobenzylchloride (117ml, 0.92M) added and heated at 100°C under an argon atmosphere. After 3hrs the reaction mixture was cooled to ambient temperature, concentrated *in vacuo*, diluted with water (800ml), extracted with ethyl acetate (2x600ml). The organic extracts were washed with brine (300ml), dried (MgSO₄), filtered, concentrated *in vacuo* to yield a brown oil which was triturated with diethyl ether/ isohexane to give compound (a) as an off-white solid (195g, 100%); ¹H nmr (d₆-DMSO, δ values): 3.81 (3H, s); 5.18 (4H, s); 6.98 (1H, m); 7.16 (1H, d); 7.36 (4H, m); 7.50 (2H, m); 7.58 (2H, m).



Compound (b)


2M Sodium hydroxide (700ml, 1.40M) was added to a solution of compound (a), methyl 3,5-di-(2-chlorobenzoyloxy) benzoate, (195g, 0.45M) in methanol (600ml)/ tetrahydrofuran (150ml) and stirred for 6hrs at 55°C. The organics were then removed *in vacuo*, acidified to pH 3-4 with concentrated hydrochloric acid, the precipitate filtered, washed with water and dried under high-vacuum at 60°C. Compound (b) was obtained as a colourless solid (.2/3NaCl) (199g, 100%); ¹H nmr (d₆-DMSO, δ values): 5.18 (4H, s); 6.93 (1H, m); 7.15 (1H, d); 7.37 (4H, m); 7.49 (2H, m); 7.58 (2H, m).



Compound (c)

- 53 -

Oxalyl chloride (7.91ml, 91mM) was added to a suspension of compound (b), 3,5-di-(2-chlorobenzoyloxy) benzoic acid.2/3NaCl (18.3g, 45.4mM) in dichloromethane (500ml) containing dimethylformamide (4 drops) under argon at ambient temperature. After 16 hrs the reaction mixture was filtered under argon, concentrated *in vacuo* then azeotroped with toluene
 5 (2x) to give the title compound as an off-white solid (17.5g, 100%); ¹H nmr (d₆-DMSO, δ values): 5.18 (4H, s); 6.94 (1H, m); 7.16 (1H, d); 7.35 (4H, m); 7.50 (2H, m); 7.58 (2H, m).


EXAMPLE B**Route 2a: 2-[3,5-di-(2-chlorobenzoyloxy)benzoyl] aminothiazole-5-carboxylic acid**

10

A solution of ethyl 2-[3,5-di-(2-chlorobenzoyloxy)benzoyl] aminothiazole-5-carboxylate (158mg, 0.28 mmol) in THF (2 ml) was treated with sodium hydroxide solution (0.57 ml of 2M, 1.4 mmol), and the reaction stirred at 40 - 50 deg C, until complete hydrolysis was achieved (with tlc monitoring, approximate reaction time 2hrs). The resulting solution was
 15 cooled, diluted with water (5 ml) and acidified to pH1 using c.HCl. The precipitate thus formed was filtered off, washed (water) and dried to give the title compound as a colourless solid, 130mg, ¹H NMR δ (d₆-DMSO): 5.25 (4H, s); 7.0 (1H, s); 7.4 (6H, m); 7.5 (2H, m); 7.6 (2H,m); 8.2 (1H, d).

The requisite starting material was prepared by a route analogous to that given in Example A.

20


EXAMPLE C**Route 2b: [3,5-di-(2-chlorobenzoyloxy)benzoyl] aminobenzene-3-carboxylic acid**

A suspension of methyl [3,5-di-(2-chlorobenzoyloxy)benzoyl] aminobenzene-3-carboxylate (455mg, 1.04 mmol) in THF was treated with sodium hydroxide solution (0.85 ml of 2M, 1.7 mmol), and the reaction stirred at ambient temperature, with tlc monitoring. Methanol (3 drops) and further additions of sodium hydroxide solution (2 x 0.85 ml of 2M, 3.4 mmol) were made, until complete hydrolysis was achieved. The resulting solution was diluted with water (30 ml) and acidified to pH1 (2M HCl); the precipitate thus formed was filtered off, washed (water) and dried to give the title compound as a colourless solid, 328mg, ¹H NMR δ (d₆-DMSO): 5.25 (4H, s); 7.0 (1H, s); 7.4 (6H, m); 7.5 (2H, m); 7.6 (2H, m); 8.2 (1H, d).

- 10 The requisite methyl ester starting material was prepared by a method analogous to that given in Example A.

EXAMPLE D

Route 3: 2-[3,5-Di-(2-chlorobenzoyloxy)benzoyl] amino-4-methyl aminomethyl thiazole




15

2-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-4-chloromethylthiazole (56mg, 0.10mM) was dissolved in 33% methylamine in methylated spirit (4ml) and stirred at ambient temperature for 16hrs. The reaction mixture was concentrated *in vacuo*, triturated with methanol, filtered and dried under high-vacuum to give the title compound as a colourless solid (30mg, 57%); ¹H nmr (d₆-DMSO, δ values): 2.63 (3H, m); 4.16 (2H, m); 5.24 (4H, s); 6.99 (1H, s); 7.38-7.44 (7H, m); 7.52 (2H, m); 7.62 (2H, m); 9.06 (1H, br s); 12.75 (1H, br s).


2-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-4-chloromethylthiazole was prepared from 3,5-di-(2-chlorobenzoyloxy) benzoyl chloride (prepared according to the method described in Example A) and 2-amino 4-chloromethyl-thiazole (JACS, 1946, 68, 2155; prepared by route 1 described in Example A).

25

EXAMPLE E**Route 4: 2-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-6-aminobenzothiazole**

2-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-6-nitrobenzothiazole (235mg, 0.40mM) was dissolved in ethyl acetate (40ml), ethanol (20ml) and dimethylformamide (5ml). 5% Palladium on carbon (46mg) was added under an argon atmosphere then the reaction mixture stirred under a hydrogen atmosphere for 16hrs. The reaction mixture was filtered through celite, concentrated *in vacuo*, triturated with methanol to give the title compound as a pale yellow solid (140mg, 63%); ¹H nmr (d₆-DMSO, δ values): 5.19 (2H, br s); 5.23 (4H, s); 6.72 (1H, dd); 6.93 (1H, m); 7.03 (1H, m); 7.35-7.44 (7H, m); 7.51 (2H, m); 7.61 (2H, m); 12.46 (1H, br s).

2-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-6-nitrobenzothiazole was prepared from 3,5-di-(2-chlorobenzoyloxy) benzoyl chloride (prepared according to the method described in Example A) and 2-amino-6-nitrobenzothiazole (prepared by route 1 described in Example A). ¹H nmr (d₆-DMSO, δ values): 5.27 (4H, s); 7.03 (1H, s); 7.38-7.46 (4H, m); 7.49-7.55 (4H, m); 7.65 (2H, m); 7.93 (1H, d); 8.30 (1H, dd); 9.09 (1H, m); 13.28 (1H, br s).

EXAMPLE F**Route 5: 5-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-[1,3,4]thiadiazole-2-sulfonic acid**

5-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-[1,3,4]thiadiazole-2-thiol (200mg, 0.38mM) was suspended in 2M NaOH (5ml), cooled (ice bath) and 30% aqueous hydrogen peroxide (0.16ml, 1.54mM) added dropwise then allowed to warm to ambient temperature. After 40hrs

the reaction mixture was filtered, washed with water then methanol and dried under high-vacuum to give the title compound as a colourless solid (122mg, 57%); ¹H nmr (d₆-DMSO, δ values): 5.20 (4H, s); 6.68 (1H, m); 7.37 (4H, m); 7.45 (2H, m); 7.50 (2H, m); 7.62 (2H, m). MS (M-H⁺)⁻ 564, 566.


5

5-[3,5-Di-(2-chlorobenzoyloxy)benzoyl]amino]-[1,3,4]thiadiazole-2-thiol was prepared from 3,5-di-(2-chlorobenzoyloxy) benzoyl chloride and 5-amino-[1,3,4]thiadiazole-2-thiol (Maybridge) by route 1 as described in Example A. ¹H nmr (d₆-DMSO, δ values): 5.21 (4H, s); 6.98 (1H, m); 7.34-7.40 (6H, m); 7.50 (2H, m); 7.59 (2H, m). MS (M-H⁺)⁻ 516, 518.

10

EXAMPLE G


Route 6: 2-[(3-isopropoxy-5-(2-chlorobenzylamino)benzoyl)amino]-5-thiazolecarboxylic acid



- 15 2-Chlorobenzaldehyde (0.012ml, 0.11mM) was added to 2-[(3-isopropoxy-5-aminobenzoyl)amino]-5-thiazolecarboxylic acid (29mg 0.09mM) and 4A molecular sieves (90mg) in methanol under an inert atmosphere at room temperature. After 1 hr sodium cyanoborohydride (7mg, 0.11mM) was added and the reaction mixture stirred for 16 hrs. The reaction mixture was filtered, concentrated *in vacuo*, the residue stirred with water then
- 20 extracted with ethyl acetate (3x10ml). The organic extracts were washed with brine (20ml), dried (MgSO₄), filtered and concentrated *in vacuo* to give the title compound as a pale yellow solid (22mg, 55%); ¹H nmr (d₆-DMSO, δ values): 1.22 (6H, d); 4.36 (2H, m); 4.58 (1H, m); 6.24 (1H, s); 6.47 (1H, m); 6.84 (2H, m); 7.26 (3H, m); 7.37 (2H, m); 7.45 (1H, m); 7.76 (1H, br s). MS [M-CO₂H]⁻ 400, 402.


25

2-[(3-isopropoxy-5-aminobenzoyl)amino]-5-thiazolecarboxylic acid was prepared as follows:



Compound (d)

3-Nitro-5-hydroxy benzoic acid (6.1g, 33.3mM) was dissolved in methanol (150ml), concentrated sulfuric acid (2.0ml) was added, and the solution stirred at room temperature for 5 days. The reaction mixture was concentrated *in vacuo*, saturated aqueous sodium hydrogencarbonate (60ml) added cautiously and the aqueous layer extracted with ethyl acetate (200ml). The organic layer was washed with brine (80ml), dried (MgSO₄), filtered and concentrated *in vacuo* to give compound (d) as a pale yellow solid (6.0g, 91%); ¹H nmr (d6-DMSO, δ values): 3.85 (3H, s); 7.67 (1H, m); 7.75 (1H, m); 8.05 (1H, m); 10.88 (1H, br s).




10

Compound (e)

2-Iodopropane (0.54ml, 5.4mM) was added to a solution of methyl 3-nitro-5-hydroxy benzoate (1.06g, 5.4mM) and potassium carbonate (1.12g, 8.1mM) in dimethylformamide (15ml) under an argon atmosphere at room temperature. The reaction mixture was heated at 60°C for 1hr then additional 2-iodopropane (0.32ml, 3.2mM) added and heating continued for a further 1hr. The reaction mixture was then concentrated *in vacuo*, water (50ml) and ethyl acetate (100ml) added. The organic layer was separated and washed with brine (40ml), dried (MgSO₄) filtered, concentrated *in vacuo* to give compound (e) as a mobile brown oil; ¹H nmr (d6-DMSO, δ values): 1.30 (6H, s); 3.90 (3H, s); 4.84 (1H, m); 7.76 (1H, m); 7.89 (1H, m); 8.16 (1H, m).

20




Compound (f)

2M Sodium hydroxide (12.3ml, 24.7mM) was added to a solution of methyl (3-nitro-5-isopropoxy) benzoic acid (1.18g, 4.9mM) in methanol (60ml) and stirred for 5hrs at room temperature. The reaction mixture was then concentrated *in vacuo*, acidified to pH 1-2 with 2M hydrochloric acid, the precipitate filtered, washed with water and dried under high vacuum over silica gel. Compound (f) was obtained as an off-white solid (1.04g, 94%); ¹H

- 58 -


nmr (d6-DMSO, δ values): 1.30 (6H, s); 4.81 (1H, m); 7.74 (1H, m); 7.85 (1H, m); 8.14 (1H, m). MS ($M-H^+$) 224.



Compound (g)


- 5 Oxalyl chloride (0.75ml, 8.6mM) was added to 3-nitro-5-isopropoxy benzoic acid (1.03g, 4.3mM) in dichloromethane (50ml) containing dimethylformamide (2 drops) under an argon atmosphere at room temperature. After 3hrs the reaction mixture was concentrated *in vacuo* and azeotroped with toluene to give an orange oil which was dissolved in dichloromethane (40ml). Ethyl 2-aminothiazole-5-carboxylate (0.89g, 5.1mM), diisopropylethylamine (1.77g, 10.3mM) and N,N-dimethylaminopyridine (50mg, 0.43mM) were added and stirred for 1hr under an argon atmosphere. After which the reaction mixture was concentrated *in vacuo* then the pale brown residue purified on silica gel using 15 to 20% ethyl acetate/isohehexane as eluant. Compound (g) was obtained as a pale yellow solid (1.56g, 92%); 1H nmr (d6-DMSO, δ values): 1.32 (6H, d); 4.88 (1H, m); 7.87 (1H, s); 8.05 (1H, s); 8.14 (1H, s); 8.45 (1H, s).

15



Compound (h)

- 10% Palladium on carbon (20mg) was added under an argon atmosphere to a solution of ethyl 2-[(3-isopropoxy-5-nitro)benzoylamino]-5-thiazolecarboxylate (209mg, 0.53mM) in ethyl acetate (35ml). Hydrogen gas was introduced and the reaction mixture stirred vigorously for 18hrs before filtering through celite and concentration *in vacuo* to give compound (h) as pale yellow solid (160mg, 83%); 1H nmr (d6-DMSO, δ values): 1.25 (6H, d); 1.29 (3H, t); 4.28 (2H, q); 4.58 (1H, m); 5.31 (2H, br s); 6.33 (1H, m); 6.81 (1H, m); 6.87 (1H, s); 8.17 (1H, s).




Compound (k)

- 59 -

2M Sodium hydroxide (0.3ml, 0.57mM) was added to a solution of ethyl 2-[(3-isopropoxy-5-amino)benzoylamino]-5-thiazolecarboxylate (40mg, 0.11mM) in tetrahydrofuran (1.2ml)/methanol (0.5ml) and heated at 50°C for 5hrs then at room temperature overnight. The reaction mixture was then concentrated *in vacuo*, acidified to pH 4-5 with 2M hydrochloric acid, the precipitate filtered, washed with water and dried under high-vacuum over silica gel. Compound (k) was obtained as a red-brown solid (35mg, 100%); ¹H nmr (d₆-DMSO, δ values): 1.27 (6H, d); 4.63 (1H, m); 6.58 (1H, s); 7.05 (1H, s); 7.16 (1H, s); 8.14 (1H, s).


EXAMPLE H

10 Route 7: 2-[(3,5-dibenzyloxybenzoyl)amino]-5-aminopyridine




To a stirred solution of 2-[(3,5-dibenzyloxybenzoyl)amino]-5-nitropyridine (910 mg) in DMF (6 ml) was added Zinc dust (1300 mg) and a solution of ferric chloride hexahydrate (1700 mg) in water (6 ml). The resulting mixture was stirred at 120°C for three hours. Allowed to cool to ambient temperature. The mixture was extracted with ethyl acetate. The extract was washed with water (50 ml), brine (50 ml), dried over MgSO₄, then volatile material was removed by evaporation to leave a solid, which was dried under high vacuum for 24 hours at 100°C to give the title compound (518 mg) as a solid, ¹H NMR δ (d₆-DMSO): 5.17 (m, 6H), 6.80 (s, 1H), 7.00 (d, 1H), 7.26 to 7.46 (m, 12H), 7.71 (s, 1H), 7.78 (d, 1H), 10.28 (br s, 1H). MS ES⁺ 426.52 (M+H)⁺.

The requisite 6-[(3,5-dibenzyloxybenzoyl)amino]-3-nitropyridine starting material was prepared by a method analogous to that given in Example A (route 1), starting from 2-amino-5-nitropyridine; ¹H NMR δ (d₆-DMSO): 5.18 (s, 4H), 6.90 (s, 1H), 7.29-7.50 (m, 12H), 8.42 (d, 1H), 8.64 (d, 1H), 9.23 (s, 1H), 11.46 (brs, 1H). MS ES⁺ 456.12 (M+H)⁺.

EXAMPLE I**Route 8: N-{6-[3,5-dibenzyloxybenzoyl]amino}-pyridin-3-yl}-2-acetoxy-2-methyl-propionamide**

- 5 To a stirred solution of 2-[(3,5-dibenzyloxybenzoyl)amino]-5-aminopyridine (200 mg) in THF (2 ml) and pyridine (2 ml) was added a solution of 2-acetoxyisobutyryl chloride (98 mg) in THF (1 ml). The mixture was stirred at ambient temperature for 16 hours. Volatile material was removed by evaporation. The residue was dissolved in ethyl acetate (50 ml), washed with water (25 ml), brine (25 ml), dried over MgSO_4 . Volatile material was removed by
- 10 evaporation to leave a gum which was triturated under ether to give the title compound (211 mg) as a solid, $^1\text{H NMR } \delta$ (d_6 -DMSO): 1.55 (s, 6H), 2.08 (s, 3H), 5.18 (s, 4H), 6.85 (s, 1H), 7.29 to 7.50 (m, 12H), 7.98 (dd, 1H), 8.13 (d, 1H), 8.61 (s, 1H), 9.70 (s, 1H), 10.72 (s, 1H). MS ES⁻ 552.22 (M-H)⁻.

15 EXAMPLE J**Route 9: N-{6-[3,5-dibenzyloxybenzoyl]amino}-pyridin-3-yl}-2-hydroxy-2-methyl-propionamide**


- To a stirred suspension of N-{6-[3,5-dibenzyloxybenzoyl]amino}-pyridin-3-yl}-2-acetoxy-2-
- 20 methyl-propionamide (158 mg) in methanol (10 ml) was added a solution of $\text{LiOH} \cdot \text{H}_2\text{O}$ (30 mg) in water (1 ml) and THF (3 ml). The mixture was stirred at ambient temperature for 20 hours. Volatile material was removed by evaporation. To the residue was added water (10 ml). Made acidic with 2M hydrochloric acid. Precipitate filtered off, washed with ethyl acetate, and dried under high vacuum to give the title compound (120 mg) as a solid, $^1\text{H NMR } \delta$ (d_6 -

- 61 -

DMSO): 1.35 (s, 6H), 5.18 (s, 4H), 6.88 (s, 1H), 7.28 to 7.48 (m, 12H), 8.08 (d, 1H), 8.22 (d, 1H), 8.82 (s, 1H), 9.90 (s, 1H), 10.96 (s, 1H). MS ES⁺ 512.16 (M+H)⁺.

EXAMPLE K


5 Route 10: 3,5-dibenzoyloxy-N-(5-[(*tert*-butylamino)carbonyl]amino)pyridin-2-yl)benzamide



A solution of *tert*-butyl isocyanate (51 mg) in THF (5 ml) was treated with 2-[(3,5-dibenzoyloxybenzoyl)amino]-5-aminopyridine (212 mg), and stirred at ambient temperature for
 10 24 hours. More *tert*-butyl isocyanate (0.34 ml) added, and stirring continued at ambient temperature for a further 4 days. Volatile material was removed by evaporation and the residue was triturated under methanol to give the title compound (159 mg) as a solid, ¹H
 NMR δ (d₆-DMSO): 1.30 (s, 9H), 5.18 (s, 4H), 6.09 (s, 1H), 6.85 (s, 1H), 7.32 to 7.50 (m, 12H), 7.78 (dd, 1H), 8.04 (d, 1H), 8.38 (s, 1H), 8.44 (s, 1H), 10.65 (s, 1H). MS ES⁺ 525.61
 15 (M+H)⁺.

EXAMPLE L

Route 11: 3,5-di(2-cyanobenzoyloxy)-N-[5-[(2-methoxyethyl)amino]pyridin-2-yl)benzamide




20

To a stirred solution of *tert*-butyl 6-[(3,5-di(2-cyanobenzoyloxy)benzoyl)amino]pyridin-3-yl(2-methoxyethyl)carbamate (237 mg) in dichloromethane (10 ml) was added trifluoroacetic acid (3 ml). The solution was stirred at ambient temperature for three hours. Volatile material was removed by evaporation. The residue was diluted in DCM (100 ml), washed with 2M sodium
 25 Hydroxide (50 ml), brine (50 ml), dried over MgSO₄. Volatile material was removed by

evaporation to give the title compound (190 mg) as a foam, ^1H NMR δ (d_6 -DMSO): 3.22 (t, 2H), 3.28 (2, 3H), 3.50 (t, 2H), 5.31 (s, 4H), 6.92 (s, 1H), 7.12 (dd, 1H), 7.34 (s, 2H), 7.57 (m, 2H), 7.75 (m, 5H), 7.82 (d, 1H), 7.91 (d, 2H), 10.49 (br s, 1H). MS ES^+ 534.41 ($\text{M}+\text{H}$) $^+$.


5 The requisite starting materials were prepared as follows:

Preparation of *tert*-butyl 2-nitropyridin-5-yl(2-methoxyethyl)carbamate



To a suspension of Cs_2CO_3 (1430 mg) in toluene was added 2-nitro-5-bromopyridine (406 mg), $\text{Pd}(\text{Ac})_2$ (44 mg), 1,1-bis(diphenylphosphino)ferrocene (322 mg) and 2-methoxyethyl
10 amine (0.26 ml). The mixture was stirred at 85°C , under Nitrogen, for 16 hours. Allowed to cool to ambient temperature. Diluted with ethyl acetate (100 ml), and filtered through a celite plug. Volatile material was removed by evaporation, the residue was purified by flash chromatography on silica, eluted with 50-100% ethyl acetate in hexane to give a solid which was added to a solution of di-*tert*-butyl-dicarbonate (436mg) and N-dimethyl-aminopyridine
15 (cat) in THF (10 ml). The solution was stirred for 14 hours at 75°C . Allowed to cool to ambient temperature, then the volatile material was removed by evaporation. The residue was dissolved in ethyl acetate (100 ml), washed with water (50 ml), brine (50 ml), dried over MgSO_4 . Volatile material was removed by evaporation, the residue was purified by flash chromatography on silica, eluted with 20-40% ethyl acetate in hexane to give the title
20 compound (359 mg) as a gum, ^1H NMR δ (CDCl_3): 1.49 (s, 9H), 3.33 (s, 6H), 3.62 (t, 2H), 3.86 (t, 2H), 8.06 (dd, 1H), 8.21 (d, 1H), 8.65 (s, 1H). MS ES^+ 298.35 ($\text{M}+\text{H}$) $^+$.

Preparation of *tert*-butyl 2-aminopyridin-5-yl(2-methoxyethyl)carbamate




25 To a solution of *tert*-butyl 2-(6-nitropyridin-3-yl)-4-methoxybutanoate (350 mg) in ethanol (20 ml) and ethyl acetate (20 ml) was added 10 % Palladium on carbon (100 mg). The suspension was stirred at ambient temperature for 16 hours under Hydrogen. Filtered through celite, then volatile material removed by evaporation to give the title compound (299 mg) as a

- 63 -

solid, ^1H NMR δ (d_6 -DMSO): 1.32 (brs, 9H), 3.18 (s, 3H), 3.34 (t, 2H), 3.56 (t, 2H), 5.84 (s, 2H), 6.37 (d, 1H), 7.17 (dd, 1H), 7.70 (d, 1H). MS ES^+ 268.34 ($\text{M}+\text{H}$) $^+$.

EXAMPLE M

5 **Route 12: *N*-(5-aminopyridin-2-yl)-3-[(2-chlorobenzyl)oxy]-5-[(2-cyanobenzyl)oxy]benzamide**




The title compound was prepared from *N*-(5-nitropyridin-2-yl)-3-[(2-chlorobenzyl)oxy]-5-[(2-cyanobenzyl)oxy]benzamide using a method similar to that described in Route 7.

10


The requisite starting materials were prepared as follows:

Preparation of 3-[[[5-nitropyridin-2-yl)amino]carbonyl]-5-[(2-cyanobenzyl)oxy]phenyl
acetate




- 15 To a stirred solution of 3-acetoxy-5-(2-cyanobenzyl)benzoic acid (8760 mg) in THF (100 ml) was added Oxalyl chloride (3.6 ml) and DMF (0.5 ml). The mixture was stirred at ambient temperature for 3 hours. Volatile material was removed by evaporation. The residue was dissolved in a mixture of THF (60 ml) and pyridine (40 ml). 2-amino-5-nitropyridine (3919 mg) added. The stirred mixture was heated to 55°C for 16 hours. Volatile material was
- 20 removed by evaporation to leave a gum which was purified by flash chromatography on silica eluted with 1-5% ethyl acetate in hexane to give the title compound (6200 mg) as a solid, ^1H NMR δ (d_6 -DMSO): 2.29 (s, 3H), 5.37 (s, 2H), 7.17 (s, 1H), 7.45 (s, 1H), 7.58 (m, 1H), 7.70 (s, 1H), 7.76 (m, 2H), 7.92 (d, 1H), 8.40 (d, 1H), 8.65 (dm, 1H), 9.21 (m, 1H), 11.57 (s, 1H). MS ES^+ 433.48 ($\text{M}+\text{H}$) $^+$.

25

Preparation of *N*-(5-nitropyridin-2-yl)-3-[(2-cyanobenzyl)oxy]-5-hydroxybenzamide


A suspension of 3-[[[(5-nitropyridin-2-yl)amino]carbonyl]-5-[(2-cyanobenzyl)oxy]phenyl acetate (5710 mg) in THF (35 ml) was treated with 25% NaOMe in methanol (6 ml). Stirred at ambient temperature for 30 minutes. Acidified with 2M hydrochloric acid (25 ml), then extracted with ethyl acetate (100 ml). The extract was washed with water (50 ml), brine (50 ml), dried over MgSO₄. Volatile material was removed by evaporation to give a solid. This was washed with hot methanol to give the title compound (4358 mg) as a solid, LCMS rt = 2.38 min (90.5%). ES⁺ 391.45 (M+H)⁺.

10


Preparation of *N*-(5-nitropyridin-2-yl)-3-[(2-chlorobenzyl)oxy]-5-[(2-cyanobenzyl)oxy]benzamide

A solution of *N*-(5-nitropyridin-2-yl)-3-[(2-cyanobenzyl)oxy]-5-hydroxybenzamide (195 mg) in DMF (3 ml) was treated with Ag₂CO₃ (165 mg) and 2-Chlorobenzyl bromide (0.073 ml). Heated to 85°C and stirred for 17 hours under Nitrogen. Allowed to cool to ambient temperature. Water (25 ml) added. Extracted with ethyl acetate (50 ml), washed with brine (25 ml), dried over MgSO₄. Volatile material was removed by evaporation to give a solid, which was purified by flash chromatography on silica eluted with 0-5% ethyl acetate in dichloromethane to give the title compound (43 mg) as a solid, ¹H NMR δ (d₆-DMSO): 5.20 (s, 2H), 5.33 (s, 2H), 6.96 (s, 1H), 7.40 (m, 5H), 7.57 (m, 2H), 7.72 (m, 2H), 7.90 (d, 1H), 8.40 (d, 1H), 8.64 (dd, 1H), 9.22 (s, 1H), 11.50 (s, 1H). LCMS rt = 3.27 min (97.4%), ES⁺ 515.50 (M+H)⁺.

20

EXAMPLE N**Route 13: 6-([3,5-Di-(benzyloxy)benzoyl]amino)-N-[2-(dimethylamino)ethyl]nicotinamide**

- 5 Diisopropylethylamine (DIPEA, 0.23ml, 1.3mM) then 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide (EDC, 126mg, 0.66mM) were added to a solution of 2-dimethylaminoethylamine (0.57ml, 0.53mM) and 6-([3,5-Di-(benzyloxy)benzoyl]amino)nicotinic acid (0.20g, 0.44mM) in dichloromethane (10ml) under argon at ambient temperature. After 16 hours the reaction mixture was evaporated *in vacuo*
- 10 and then chromatographed on SiO₂ using a gradient elution of 10 to 25% methanol in dichloromethane. The fractions containing product were evaporated to give a cream solid (0.052g, 25%); ¹H NMR δ (d₆-DMSO): 2.67 (6H, s); 3.11 (2H, m); 3.62 (2H, m); 5.18 (4H, s); 6.88 (1H, s); 7.27-7.52 (12H, br m); 8.18-8.36 (2H, m); 8.90 (1H, s); 10.20 (1H, br s).


15 EXAMPLE O**Route 14: 2-[3,5-Di-(2-chlorobenzyloxy) benzoylamino]-5-hydroxymethyl pyridine**

- To a cold (-15 degC) solution of 2-[3,5-Di-(2-chlorobenzyloxy)benzoyl]amino-pyridine-5-carboxylic acid (305 mg, 0.59 mmol) in dimethoxy ethane (5ml) was added 4-methyl
- 20 morpholine (80μl, 1eq) and isobutyl chloroformate (76μl, 1.02 eq). The reaction mixture was stirred at -15deg C for 15 mins and then filtered; the residue was washed with dimethoxy ethane (5x1ml). The filtrate and washings were cooled to -15 deg C and treated with a suspension of sodium borohydride (22mg, 1eq) in water (1ml). After the effervescence had ceased, water (50ml) and ethyl acetate (30ml) were added; the reaction mixture was
- 25 evaporated to dryness and the residue absorbed onto silica. The required compound was isolated by flash chromatography (eluting with 5% methanol in dichloromethane) to give the

title compound as a colourless solid (97 mg), ^1H NMR δ (d_6 -DMSO): 4.5 (1H, d), 5.25 (s, 4H), 6.9 (s, 1H), 7.40 (m, 6H), 7.5 (m, 2H), 7.6 (m, 2H), 7.75 (dd, 1H), 8.10 (d, 1H), 8.3 (s, 1H), 10.8 (br s, 1H); LCMS r_t = 3.25 min (100%), ES^+ 509 ($\text{M}+\text{H}$) $^+$.

5 EXAMPLE P

Route 15: N-[6-[3,5-di-(2-chlorobenzoyloxybenzoyl)amino]-pyridin-2-yl]-2-acetamide




To a solution of 2-[(3,5-di-(2-chlorobenzoyloxybenzoyl)amino]-6-aminopyridine (220 mg, 0.45 mmol) in tetrahydrofuran (4 ml) was added pyridine (43 mg, 0.54 mmol) and acetyl chloride (42 mg, 0.54 mmol), and the reaction mixture stirred at ambient temperature for 16 hours. The reaction mixture was diluted with diethyl ether and washed successively with water, 1M citric acid, and water; the solution was dried over magnesium sulfate and the solvent removed *in vacuo* to give a yellow solid (154mg). Trituration with methanol gave the title compound (75mg), ^1H NMR δ (d_6 -DMSO): 3.3 (3H, s), 5.25 (s, 4H), 6.95 (s, 1H), 7.3 (d, 2H), 7.4 (m, 4H), 7.5 (m, 2H), 7.6 (m, 2H), 7.7 (m, 1H), 7.8 (m, 2H), 10.14 (br s, 1H), 10.36 (br s, 1H); ES^+ 536/538 ($\text{M}+\text{H}$) $^+$.

The starting material, 2-[(3,5-di-(2-chlorobenzoyloxybenzoyl)amino]-6-aminopyridine, is exemplified herein as Example number 106.

EXAMPLE Q

Route 16: 3,5-bis(benzyloxy)-N-[5-(1H-tetrazol-5-yl)pyridin-2-yl]benzamide




Tributyltin azide (156 μL , 0.57 mmol) was added to a suspension of 3,5-bis(benzyloxy)-N-(5-cyanopyridin-2-yl)benzamide (180 mg, 0.41 mmol) in toluene (3 mL). The mixture was

heated at reflux for 16 hours. The suspension was cooled and partitioned between ethyl acetate and hydrochloric acid (1M). The organic layer was concentrated in vacuo and the residue was purified by MPLC on silica MPLC (eluting with 1% methanol / DCM to 15% methanol / DCM). The tetrazole was obtained as a colourless solid (113 mg, 57%). ¹H NMR δ (d₆-DMSO): 5.19 (4H, s); 6.88 (1H, s); 7.26-7.48 (12H, m); 8.40 (1H, d); 8.46 (1H, dd); 9.04 (1H, s); 11.13 (1H, br s); ^{m/z} (LCMS; ESI+) 479 (MH⁺).

The requisite starting material was prepared as follows:

10 Preparation of 3,5-bis(benzyloxy)-N-(5-cyanopyridin-2-yl)benzamide




The title compound was prepared as described in Example A (route 1), starting from 2-amino-5-cyanopyridine and 3,5-bis(benzyloxy) benzoyl chloride, ¹H NMR δ (d₆-DMSO): 5.19 (4H, s); 6.89 (1H, m); 7.26-7.46 (12H, m); 8.27 (1H, dd); 8.33 (1H, d); 8.84 (1H, s); 11.23 (1H, br s); ^{m/z} (LCMS; ESI+) 436 (MH⁺).

The requisite 2-amino-5-cyanopyridine starting material may be purchased (Bionet Research, and other suppliers), or may be prepared according to the method given in WO95/06034.

20 EXAMPLE R

Route 17: 3,5-bis(benzyloxy)-N-[5-(5-oxo-4,5-dihydro-1,2,4-oxadiazol-3-yl)pyridin-2-yl]benzamide




- 68 -

Ethyl chloroformate (32 μ L, 0.33 mmol) was added to a solution of 3,5-bis(benzyloxy)-*N*-{5-[(hydroxyamino)(imino)methyl]pyridin-2-yl}benzamide (140 mg, 0.30 mmol) in pyridine (5 mL). This solution was heated at reflux overnight. The mixture was cooled and concentrated under reduced pressure. DCM and methanol were used to dissolve the remaining material and the solution was washed with water. The organic solution was concentrated under reduced pressure and the residue was purified on silica by MPLC (eluting firstly with 5% methanol / DCM then 10% methanol / DCM). The title compound was obtained as a colourless solid (103 mg, 70%). $^1\text{H NMR } \delta$ (d_6 -DMSO): 5.19 (4H, s); 6.87 (1H, s); 7.28-7.46 (12H, m); 8.21 (1H, dd); 8.38 (1H, d); 8.79 (1H, s); 11.14 (1H, br s); m/z (LCMS; ESI+) 495 (MH^+).

10

The requisite starting material was prepared as follows:


Preparation of 3,5-bis(benzyloxy)-*N*-{5-[(hydroxyamino)(imino)methyl]pyridin-2-yl}benzamide



15 A mixture of 3,5-bis(benzyloxy)-*N*-(5-cyanopyridin-2-yl)benzamide (212 mg, 0.49 mmol), triethylamine (170 μ L, 1.22 mmol) and hydroxylamine hydrochloride (85 mg, 1.22 mmol) in ethanol (5 mL) was heated at reflux overnight. The mixture was cooled and concentrated under reduced pressure. The residue was purified by MPLC on silica eluting with 5% methanol / DCM then 15% methanol / DCM. The title compound was obtained as a colourless solid (171 mg, 75%). $^1\text{H NMR } \delta$ (d_6 -DMSO): 5.19 (4H, s); 5.92 (2H, s); 6.87 (1H, s); 7.28-7.48 (12H, m); 8.06 (1H, dd); 8.17 (1H, d); 8.65 (1H, s); 9.68 (1H, s); 10.85 (1H, br s); m/z (LCMS; ESI+) 469 (MH^+).

The requisite 3,5-bis(benzyloxy)-*N*-(5-cyanopyridin-2-yl)benzamide was prepared as described in Example P (route 15).

25

EXAMPLE S**Route 18: [(2-[[3,5-bis(benzyloxy)benzoyl]amino]pyridin-5-yl)amino](oxo)acetic acid**

Methyl oxalyl chloride (37 μ L, 0.4 mmol) was added to a mixture of *N*-(5-aminopyridin-2-yl)-
5 3,5-bis(benzyloxy)benzamide (150 mg, 0.36 mmol) and triethylamine in DCM (5 mL). The
mixture was stirred for 1 hour under an atmosphere of nitrogen. The solution was diluted with
DCM and washed with water. The organics were concentrated under reduced pressure and the
residue was purified on silica by MPLC (eluting with 1% methanol / DCM to 15% methanol /
DCM) to give a colourless solid (110 mg). This material was dissolved in THF (2 mL). Water
10 (3 mL) and sodium hydroxide (0.5 mL, 2M, 1 mmol) were added. The mixture was stirred for
1 hour before being acidified with hydrochloric acid (2M) and diluted with water. The
resulting precipitate was isolated by filtration, washed with water and dried *in vacuo*. The title
compound was obtained as a colourless solid (88 mg, 50%). ^1H NMR δ (d_6 -DMSO): 5.18 (4H,
s); 6.88 (1H, s); 7.30-7.50 (12H, m); 8.17 (2H, s); 8.79 (1H, s); 10.79 (1H, s); 10.93 (1H, br
15 s); m/z (LCMS; ESI+) 498 (MH^+).




The requisite starting material was prepared according to Example H (route 7).

EXAMPLE T:








20 By analogous methods to those described above the following compounds, Example numbers
 T_1 to T_{20} , were also made.






Compound T_9 was prepared by Route 1b (multi-parallel synthesis), as follows. To the
appropriate acid (6.0 mmol) in dichloromethane (25 mls) was added 1 drop of
25 dimethylformamide and the mixture stirred at room temperature under argon. The oxalyl
chloride (0.867 mls) was added to the acid and stirred at room temperature for 2 hrs. The
solvent was removed in Genevac DD4, and resulting residue azeotroped with
dichloromethane (3 x 10 mls), then dried high vacuum for 2hrs. The resulting acid chloride
was then dissolved in THF (30 mls) and 5mls of the solution was added to one of the set of

six amines in THF / Pyridine (5mls). The resulting mixture was stirred overnight at room temperature, diluted with ethyl acetate (5mls). The resulting solution was transferred to the Alex automated extractor and washed with water (2x5mls), sodium hydrogen carbonate (5mls), 1M citric acid (5mls), brine (5mls) dried (magnesium sulphate) and evaporated in 5 Genevac DD4. The resulting gum was triturated with methanol (1-2 mls) and the resulting solid filtered, washed methanol and air-dried.


Example	Structure	Route	NMR
1		1	1H NMR d (d6-DMSO): 5.26 (4H, s); 6.96 (1H, m); 7.38-7.45 (6H, m); 7.53 (2H, m); 7.62 (2H, m); 8.43 (1H, d); 8.49 (1H, m); 9.42 (1H, m); 11.13 (1H, s).
2		1	1H NMR d (d6-DMSO): 5.25 (4H, s); 6.97 (1H, m); 7.38-7.45 (6H, m); 7.53 (2H, m); 7.63 (2H, m); 8.64 (1H, d); 9.26 (1H, d); 11.33 (1H, s).
3		1	1H NMR d (d6-DMSO): 5.24 (4H, s); 6.95 (1H, s); 7.35-7.40 (6H, m); 7.50 (2H, m); 7.60 (2H, m); 8.61 (1H, s); 9.22 (1H, s); 11.25 (1H, br s).

Example	Structure	Route	NMR
4		1	¹ H NMR d (d6-DMSO): 5.36 (4H, s); 7.00 (1H, m); 7.44 (2H, d); 7.55-7.64 (2H, m); 7.77 (4H, m); 7.93 (2H, d); 8.43 (1H, d); 8.49 (1H, m); 9.43 (1H, s); 11.17 (1H, s).
5		1	¹ H NMR d (d6-DMSO): 3.90 (3H, s); 5.24 (4H, s); 6.97 (1H, m); 7.39 (6H, m); 7.50 (2H, m); 7.60 (2H, m); 9.02 (1H, s); 9.52 (1H, s); 11.54 (1H, br s).
6		2	¹ H NMR d (d6-DMSO): 5.24 (4H, s); 6.96 (1H, m); 7.39 (6H, m); 7.51 (2H, m); 7.62 (2H, m); 8.98 (1H, s); 9.48 (1H, s); 11.44 (1H, br s).
7		2 *	¹ H NMR d (d6-DMSO): 5.34 (4H, s); 7.00 (1H, s); 7.57 (2H, m); 7.75 (4H, m); 7.91 (2H, d); 9.00 (1H, s); 9.52 (1H, s); 11.53 (1H, s); 13.43, (1H br s).
8		1	¹ H NMR d (d6-DMSO): 2.29 (3H, s); 2.33 (3H, s); 3.24 (m, 2H); 4.21 (2H, t); 5.12 (2H, s); 6.80 (1H, m); 7.21 (4H, m); 7.31 (1H, m); 7.40 (2H, m); 8.39 (1H, m); 8.45 (1H, m); 8.82 (1H, s); 9.38 (1H, s); 11.06 (1H, br s).


Example	Structure	Route		NMR
9		1b		¹ H NMR δ (d ₆ -DMSO): 1.25 (d, 12H), 4.7 (hept, 2H), 6.6 (d, 1H), 7.2 (d, 2H), 8.4 (d, 1H), 8.45 (t, 1H), 9.4 (s, 1H), 11.0 (br s, 1H).
10		1		¹ H NMR δ (d ₆ -DMSO): 2.37 (s, 3H), 3.24 (t, 2H), 4.23 (t, 2H), 4.65 (d, 2H), 5.28 (d, 1H), 5.42 (d, 1H), 6.05 (m, 1H), 6.75 (s, 1H), 7.23 (s, 2H), 8.43 (s, 1H), 8.84 (s, 1H), 9.40 (s, 1H), 11.07 (br s, 1H).
11		1		¹ H NMR δ (d ₆ -DMSO): 2.32 (s, 3H), 2.35 (s, 3H), 3.21 (t, 2H), 4.21 (t, 2H), 5.13 (s, 2H), 6.81 (s, 1H), 7.14-7.26 (m, 4H), 7.32 (1H, s), 7.41 (1H, d), 8.51 (s, 1H), 8.81 (s, 1H), 9.39 (s, 1H), 11.34 (brs, 1H).
12		1	364	¹ H NMR δ (d ₆ -DMSO): 2.12 (s, 6H), 3.81 (s, 3H), 5.05 (s, 2H), 6.95 (s, 1H), 7.05 (s, 2H), 7.1 (s, 1H), 7.72 (d, 1H), 7.78 (s, 1H), 8.36 (d, 1H), 8.43 (s, 1H), 9.4 (s, 1H), 10.92 (br s, 1H).
13		1b	412	410
14		1b	330	
15		1b	288	

Example	Structure	Route			NMR
16		1b	312		
17		1b	452		
18		1b	400		
19		2a, 1c	428	426	δ_H (500MHz, DMSO- d_6) 1.28 (6H, d), 3.07 (2H, t), 4.26 (2H, t), 4.70 (1H, m), 6.71 (1H, m), 7.12 (1H, m), 7.24 (2H, m), 7.30 (1H, m), 7.46 (1H, m), 8.98 (1H, d), 9.48 (1H, d), 11.33 (1H, s), 13.24 (1H, br s).
20		1a	382		

* For Example 7, the ester intermediate was prepared by route 1:




1H NMR δ (d_6 -DMSO): 3.90 (3H, s); 5.34 (4H, s); 7.01 (1H, s); 7.43 (2H, s); 7.58 (2H, m);
5 7.74 (4H, m); 7.91 (2H, d); 9.02 (1H, s); 9.52 (1H, s); 11.57 (1H, br s).

EXAMPLE U**2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-dimethylamino] benzoylamino]-[1,3,4]-thiadiazole****(Route 19)**


Formaldehyde (37% in water) (0.033ml, 0.44mM) was added to 2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-amino] benzoylamino]-[1,3,4]-thiadiazole (27mg 0.074mM) and 4A molecular sieves (0.2g) in methanol (4ml)/acetonitrile (3ml)/ g.AcOH (2 drops) under an inert atmosphere at room temperature. After 150 mins sodium cyanoborohydride (7mg, 0.12mM) was added and the reaction mixture stirred for 40 hrs. The reaction mixture was filtered, concentrated *in vacuo*, the residue acidified with 2M HCl to precipitate a colourless solid. Purified on silica gel (50 to 75% EtOAc/iso-hexane) gave the title compound as a colourless solid (25mg, 85%); ¹H NMR δ (d₆-DMSO): 2.35 (s, 3H), 2.93 (s, 6H), 3.22 (m, 2H), 4.19 (m, 2H), 6.41 (m, 1H), 6.98 (m, 1H), 7.06 (m, 1H), 8.80 (s, 1H), 9.17 (s, 1H).

15


The requisite 2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-amino] benzoylamino]-[1,3,4]-thiadiazole starting material was prepared as follows:




10% Palladium on carbon (80mg) was added under an argon atmosphere to a solution of 2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-nitro] benzoylamino]-[1,3,4]-thiadiazole (0.38g, 0.99 mM) in ethyl acetate (40ml). Hydrogen gas was introduced and the reaction mixture stirred vigorously for 18hrs before filtering through celite, concentration *in vacuo* and replacement of the catalyst (80mg). After stirring under hydrogen gas for a further 18hrs a final catalyst change was carried out. After which the crude aniline was purified on silica gel (1% to 4% MeOH/DCM) to give 2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-amino] benzoylamino]-[1,3,4]-thiadiazole as a colourless solid (0.1g, 28%); MS (M-H⁺)⁻ 360.



Oxalyl chloride (0.20ml, 2.35mM) was added to 3-(2-(4-methyl thiazol-5-yl) ethoxy)-5-nitro benzoic acid (0.72g, 2mM) in dichloromethane (30ml) containing DMF (2 drops) under an argon atmosphere at room temperature. After 3hrs the reaction mixture was concentrated *in vacuo* and azeotroped with toluene to give an off-white solid. The acid chloride and 2-amino-[1,3,4]-thiadiazole (0.19g, 1.9 mM) were dissolved in DCM (20ml) then DIPEA (0.96ml, 5.6 mM) and DMAP (0.04g, 0.3 mM) added. After stirring overnight under argon the reaction mixture was concentrated, purified on silica gel (50%to75%to100% EtOAc/iso-hexane) gave a pale yellow solid which was triturated with MeOH to give 2-[3-(2-(4-methyl thiazol-5-yl) ethoxy)-5-nitro] benzoylamino]-[1,3,4]-thiadiazole as a colourless solid (0.30g, 48%); ¹H NMR δ (d₆-DMSO): 2.37 (s, 3H), 3.26 (t, 2H), 4.35 (t, 2H), 7.89 (m, 1H), 8.09 (s, 1H), 8.47 (s, 1H), 8.81 (s, 1H), 9.24 (s, 1H).




DIAD (3.16ml, 16.1mM) was added to a stirred solution of methyl 3-nitro-5-hydroxy benzoate (2.11g, 10.7mM), 4-(2-hydroxy ethyl)-5-methylthiazole (1.55ml, 12.8mM), and triphenylphosphine (4.21g, 16.1 mM) in THF (50ml) under an argon atmosphere at room temperature. After 1hr reaction mixture concentrated *in vacuo*, residue triturated with diethyl ether to give a colourless solid (triphenylphosphine oxide). Diethyl ether conc. to give a dark brown gum, purification on silica gel (50% to 75% EtOAc/iso-hexane) gave the product contaminated with reduced DIAD and triphenylphosphine oxide (6.8g). The crude product was dissolved/suspended in MeOH (80ml), 2M NaOH (20ml, 40 mM) added, heated at 65°C for 4 hrs then cooled and concentrated. The residue was diluted with water (140ml)/ 2M NaOH (40ml), the precipitated triphenylphosphine oxide filtered, then acidified with c. HCl to pH1-2. The precipitate was filtered, washed with water, dried under high-vacuum to give 3-(2-(4-methyl thiazol-5-yl) ethoxy)-5-nitro benzoic acid as a colourless solid (3.12g, 79% over 2 steps); ¹H NMR δ (d₆-DMSO): 2.39 (s, 3H), 3.23 (t, 2H), 4.35 (t, 2H), 7.78 (s, 1H), 7.90 (m, 1H), 8.22 (s, 1H), 8.93 (s, 1H).

EXAMPLE V**2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-hydroxy] benzoylamino]-[1,3,4]-thiadiazole****(Route 20)**


A solution of 2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-allyloxy] benzoylamino]-[1,3,4]-thiadiazole (1.1g, 2.7 mmol) in tetrahydrofuran (40ml) was stirred under an argon atmosphere and treated with Meldrum's acid (0.79g, 5.4mmol) and tetrakis (triphenyl phosphine) palladium (0) (825mg, 0.7mmol, 0.25 eq) and the resulting yellow solution stirred at ambient temperature for 2 hours. Sequential triturations with dichloromethane and hot tetrahydrofuran gave 2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-hydroxy] benzoylamino]-[1,3,4]-thiadiazole as a colourless solid (0.59g, 59%), ¹H NMR δ (d₆-DMSO): 2.35 (s, 3H), 3.2 (t, 2H), 4.2 (t, 2H), 6.55 (m, 1H), 7.05 (s, 1H), 7.2 (s, 1H), 8.81 (s, 1H), 9.2 (s, 1H), 9.8 (br s, 1H); m/z 363 (M+H)⁺, 361 (M-H)⁻.

15



The requisite 2-[3-{2-(4-methyl thiazol-5-yl) ethoxy}-5-allyloxy] benzoylamino]-[1,3,4]-thiadiazole starting material was prepared according to the appropriate generic alkylation method, and the resulting benzoic acid coupled with 1,3,4-thiadiazole according to Route 1. The analytical data on all intermediates was consistent with the proposed structures.


20

EXAMPLE W**2-(3-isopropoxy-5-dimethylaminomethyl)benzoyl aminothiazole****(Route 21)**

- 5 A solution of 2-(3-isopropoxy-5-formyl)benzoyl aminothiazole (0.11g, 0.39mmol) in dichloromethane was treated with dimethylamine (0.074ml of an approx. 5.6M solution in ethanol, 0.41mmol, 1.1 eq) and stirred under argon for 10 mins. To the solution was added sodium tris-acetoxy borohydride (0.11g, 0.53mmol, 1.4 eq), and the resulting mixture stirred overnight at ambient temperature. Further reagents were then added (same quantities as
- 10 before) and the mixture again stirred overnight at ambient temperature. The solution was treated with saturated sodium bicarbonate solution (10ml) and stirred for 20 mins; it was then extracted twice with dichloromethane, the organic extracts dried over magnesium sulfate and evaporated *in vacuo* to give the product as a colourless oil. This was dissolved in ethyl acetate and the solution treated with an ethereal solution of HCl (excess of 1M); the precipitate thus
- 15 formed was filtered under argon and washed with diethyl ether to give 2-(3-isopropoxy-5-dimethylaminomethyl)benzoyl aminothiazole hydrochloride as a colourless solid (0.1g, 72%), ¹H NMR δ (d₆-DMSO): 1.31 (d, 6H), 2.71 (s, 6H), 4.26 (m, 2H), 4.76 (m, 1H), 7.29 (d, 1H), 7.42 (m, 1H), 7.55 (d, 1H), 7.70 (s, 1H), 10.66 (bs, 1H).
- 20 The requisite starting material was prepared as follows:

EXAMPLE X**2-(3-isopropoxy-5-formyl)benzoyl aminothiazole****(Route 22):**

25




A solution of 2-(3-isopropoxy-5-hydroxymethyl)benzoyl aminothiazole (0.115g, 0.39mmol) in tetrahydrofuran (8ml) was treated with manganese dioxide (0.27g, 3.1mmol, 8eq) and the resulting suspension stirred overnight at ambient temperature; additional oxidant (0.1g portions) was added until all the starting material was consumed (tlc). The suspension was
5 filtered, the residue washed well with ethyl acetate, and the combined filtrate and washings evaporated *in vacuo* to give the product as a pale yellow solid, ¹H NMR δ (d₆-DMSO): 1.31 (d, 6H), 4.82 (m, 1H), 7.26 (d, 1H), 7.56 (d, 1H), 7.59 (s, 1H), 7.94 (d, 1H), 8.15 (s, 1H), 10.00 (s, 1H), 12.77 (bs, 1H).

10 The requisite starting material was prepared as follows:

EXAMPLE Y


2-(3-isopropoxy-5-hydroxymethyl)benzoyl aminothiazole

(Route 23)




15

Standard ester cleavage of 2-(3-isopropoxy-5-acetoxymethyl)benzoyl aminothiazole (0.15g, 0.46 mM) using 2M NaOH/THF/MeOH for 1 hour gave 2-(3-isopropoxy-5-hydroxymethyl)benzoyl aminothiazole as a colourless solid (0.149g, 100%); ¹H NMR δ (d₆-DMSO): 1.28 (d, 6H), 4.51 (s, 2H), 4.71 (m, 1H), 7.05 (s, 1H), 7.25 (d, 1H), 7.50 (s, 1H), 7.53
20 (d, 1H), 7.58 (s, 1H), 12.50 (bs, 1H).



The requisite 2-(3-isopropoxy-5-acetoxymethyl)benzoyl aminothiazole was prepared by a standard coupling between 3-isopropoxy-5-acetoxymethyl benzoyl chloride and 2-
25 aminothiazole according to Route 1, to give the title compound as a pale yellow oil, δ (d₆-DMSO): 1.3 (d, 6H), 2.1 (s, 3H), 4.75 (hept, 1H), 5.1 (s, 2H), 7.15 (s, 1H), 7.25 (d, 1H), 7.65 (d, 1H), 7.6 (m, 2H), 12.6 (bs, 1H).


The requisite 3-isopropoxy-5-acetoxymethyl benzoic acid was prepared as follows:



A solution of 3-isopropoxy-5-hydroxymethyl benzoic acid (0.77g, 3.7mmol) in
 5 dichloromethane (20ml) was cooled (ice-bath) and stirred under argon; pyridine (1.18ml,
 14.6mmol, 4eq) was added followed dropwise by acetyl chloride (0.55ml, 7.7 mmol, 2.1 eq).
 The mixture was stirred for 5 mins, then allowed to warm to ambient temperature over 90
 mins. Water (20ml) was added, the mixture stirred for 2 hrs, then allowed to stand overnight.
 The organic layer was separated, the aqueous portion washed with dichloromethane, and the
 10 dichloromethane fractions combined and evaporated. The resulting pale yellow oil was
 dissolved in ethyl acetate and the solution washed with 0.05M aqueous HCl (20ml); the
 organic layer was separated, dried over magnesium sulfate and evaporated *in vacuo* to give the
 product as a pale yellow solid, ^1H NMR δ (d_6 -DMSO): 1.25 (d, 6H), 2.06 (s, 3H), 4.65 (hept,
 1H), 5.05 (s, 2H), 7.12 (s, 1H), 7.31 (d, 1H), 7.46 (s, 1H).


15

The requisite 3-isopropoxy-5-hydroxymethyl benzoic acid starting material was prepared as follows:



Standard 2M NaOH/THF/MeOH cleavage of methyl 3-isopropoxy-5-hydroxymethyl benzoate
 20 (1.12g, 5.0 mM) gave the title compound as a colourless solid (0.98g, 94%); ^1H NMR δ (d_6 -
 DMSO): 1.25 (d, 6H), 4.47 (s, 2H), 4.60 (m, 1H), 5.23 (bs, 1H), 7.06 (s, 1H), 7.24 (s, 1H),
 7.45 (s, 1H).


The requisite methyl 3-isopropoxy-5-hydroxymethyl benzoate starting material was prepared
 25 as follows:



- 80 -

Mono-methyl-5-isopropoxy-isophthalate (5.15g, 21.6 mM) was dissolved in THF (180ml), cooled to 2°C and borane:THF complex (72ml of 1.5M solution in THF, 0.11 mM) added dropwise over 15 mins, maintaining an internal temperature of < 5°C. After 15 mins the reaction mixture was warmed to ambient temperature, stirred for 3 hrs before cooling (ice bath) and quenching with pieces of ice. When no further reaction observed brine (150ml)/ diethyl ether (150ml) added. The organic layer was removed, aqueous extracted with additional diethyl ether (1x100ml), combined organics washed with brine (1x100ml), dried (MgSO₄), filtered and concentrated. Purified on silica gel (20-25% EtOAc/isohexane) to give the title compound as a colourless solid (3.57g, 74%); ¹H NMR δ (d₆-DMSO): 1.26 (d, 6H), 3.82 (s, 3H), 4.50 (d, 2H), 4.63 (m, 1H), 5.26 (t, 1H (-OH)), 7.10 (s, 1H), 7.25 (s, 1H), 7.47 (s, 1H).

The requisite mono-methyl-5-isopropoxy-isophthalate starting material was prepared as follows:




15

2M NaOH (1.03g, 25.9 mM) in methanol (9 ml) was added to a solution of dimethyl-5-isopropoxy-isophthalate (5.68g, 22.5 mM) in acetone (45ml) and stirred at ambient temperature overnight. The reaction mixture was concentrated, acidified (2M HCl) to pH1-2, filtered, washed with water and dried under high vacuum to give 14279/66/1 as a colourless solid (5.25g, 98%) (contains 15-20% diacid); MS (M-H⁺)⁻ 237.

20

The requisite dimethyl-5-isopropoxy-isophthalate starting material was prepared as follows:



Dimethyl-5-hydroxy-isophthalate (5.2g, 24.6 mM), potassium carbonate (4.07g, 29.5 mM), potassium iodide (0.82g, 4.9 mM) and 2-bromopropane (2.4ml, 25.8 mM) in DMF (50ml) was heated at 90°C for 3hrs, after which additional 2-bromopropane (2.4ml), potassium carbonate (2.2g) was added, heated for a further 4hrs then cooled to room temperature and concentrated. EtOAc (150ml) was added then washed with water, brine, dried (MgSO₄),


25

filtered and concentrated to give a pale yellow oil which solidified on standing (6.0g, 97%); MS (MH⁺) 253.

EXAMPLE Z

5 2-(3-isopropoxy-5-formyl)benzoyl aminothiazole-5-carboxylic acid

(Route 24)




A solution of 2-(3-isopropoxy-5-hydroxymethyl)benzoyl aminothiazole-5-carboxylic acid (0.42g, 1.25mmol) in tetrahydrofuran (50ml) was treated with Dess-Martin periodinane (DMP, 0.58g, 1.37mmol, 1.1 eq) and stirred at ambient temperature for 90 mins. The solvent was removed *in vacuo*, and the residue treated with dichloromethane and filtered. The residue was partitioned between ethyl acetate and sat'd sodium bicarbonate solution containing sodium thiosulfate solution (ca 7 eq of 2.1 M), and the resulting 2-phase mixture stirred vigorously before being acidified to ca pH6. The title compound was isolated by filtration as a
 15 colourless solid, (0.145g, 35%), ¹H NMR δ (d₆-DMSO): 1.32 (d, 6H), 4.79 (m, 1H), 7.62 (m, 1H), 7.92 (m, 1H), 8.13 (s, 1H), 8.18 (s, 1H), 10.03 (s, 1H).

The requisite 2-(3-isopropoxy-5-hydroxymethyl)benzoyl aminothiazole-5-carboxylic acid starting material was prepared according to the procedure given in Route 2a and is
 20 exemplified as Example II₈₁.

EXAMPLE AA

Z-[2-[3-isopropoxy-5-(3-methyl-but-1-enyl)]benzoyl aminothiazole-5-carboxylic acid]

(Route 25)



25 A solution of iso-butyl triphenyl phosphonium bromide (0.45g, 1.13mmol, 3.1 eq) in tetrahydrofuran (20ml) was treated with potassium t-butoxide (1.1ml of 1M in

- 82 -


tetrahydrofuran, 1.13mmol, 3.1 eq) and stirred at 0 deg C under argon. To this was added 2-(3-isopropoxy-5-formyl)benzoyl aminothiazole-5-carboxylic acid (0.122g, 0.36mmol), and the resulting solution stirred for 100 mins, allowing to warm to ambient temperature. Water was added and the solvent removed *in vacuo*; the residue was partitioned between water and ethyl acetate and the layers separated. The aqueous portion was neutralised (2M HCl) and extracted twice with ethyl acetate; the organic extracts were dried (MgSO₄), filtered and concentrated and the residue purified by chromatography on silica gel (10g Bondelut cartridge, eluting with dichloromethane containing methanol, 10% v/v) to give the title compound as a colourless solid (0.012g, 9%); ¹H NMR δ (d₆-DMSO): 1.01 (d, 6H), 1.29 (d, 6H), 2.81 (m, 1H), 4.72 (m, 1H), 6.53 (dd, 1H), 6.29 (d, 1H), 6.97 (s, 1H), 7.50 (s, 1H), 7.53 (s, 1H), 8.11 (s, 1H), 8.18 (s, 1H).

The requisite 2-(3-isopropoxy-5-formyl)benzoyl aminothiazole-5-carboxylic acid was prepared according to the procedure given under Example Z (Route 24); see Example II₈₉.


EXAMPLE BB

2-[3-isopropoxy-5-(4-methyl-1-piperidinocarbonylmethylenoxy)] benzoyl aminothiazole.


(Route 26)




This was prepared by a standard acid chloride coupling (Example A, Route 1), starting from 2-(3-isopropoxy-5-carboxymethylene oxy) benzoyl aminothiazole, to give the title compound, ¹H NMR δ (d₆-DMSO): 1.28 (d, 6H), 2.18 (s, 3H), 2.24 (m, 2H), 2.32 (m, 2H), 3.44 (ap t, 4H), 4.65 (m, 1H), 4.85 (s, 2H), 6.68 (ap t, 1H), 7.19 (m, 1H), 7.24 (ap d, 2H), 7.55 (ap d, 1H), 12.45 (bs, 1H); m/z 419 (M+H)⁺, 417 (M-H)⁻.



The requisite 2-(3-isopropoxy-5-carboxymethylene oxy) benzoyl aminothiazole was prepared from 2-(3-isopropoxy-5-methoxycarbonylmethylene oxy) benzoyl aminothiazole by standard ester hydrolysis (Route 2a); ^1H NMR δ (d_6 -DMSO): 1.28 (d, 6H), 4.69 (m, 1H), 4.73 (s, 2H), 5 6.66 (ap t, 1H), 7.22 (s, 1H), 7.27 (ap d, 2H), 7.53 (ap d, 1H); m/z 337.31 ($M+H$) $^+$ 335.27 ($M-H$) $^-$




The requisite 2-(3-isopropoxy-5-methoxycarbonylmethylene oxy) benzoyl aminothiazole 10 starting material was prepared from 3-isopropoxy-5-(methoxycarbonyl)methoxybenzoic acid and 2-aminothiazole (48% isolated yield) by a standard acid chloride coupling (Route 1); ^1H NMR δ (d_6 -DMSO): 1.27 (d, 6H), 3.70 (s, 3H), 4.71 (m, 1H), 4.86 (s, 2H), 6.99 (t, 1H), 7.23 (t, 1H), 7.26-7.27 (m, 2H), 12.53 (s, 1H); m/z 351.31 ($M+H$) $^+$, 349.28 ($M-H$) $^-$




The requisite starting material was prepared from 3-isopropoxy-5-(methoxycarbonyl methylene oxy) benzoic acid was prepared by monoesterification of 3-isopropoxy-5-(carboxymethylene oxy) benzoic acid (78% isolated yield) using the conditions of Ram and Charles, *Tetrahedron* **1997**, 53 (21), pp.7335-7340: ^1H NMR δ (d_6 -DMSO): 1.25 (d, 6H), 20 3.69 (s, 3H), 4.65 (m, 1H), 4.83 (s, 2H), 6.71 (ap t, 1H), 6.98 (s, 1H), 7.01 (s, 1H), 12.97 (bs, 1H); m/z 554.27 ($2M+NH_4$) $^+$, 267.26 ($M-H$) $^-$

3-isopropoxy-5-(carboxymethoxy)benzoic acid



The title compound was prepared from methyl (3-isopropoxy-5-(t-butyloxycarbonyl)methoxy)benzoate (56% isolated yield) using standard hydrolysis method

- 5 2a. $^1\text{H NMR } \delta$ (d_6 -DMSO): 1.25 (d, 6H), 4.62 (m, 1H), 4.69 (s, 2H), 6.67 (ap t, 1H), 6.96 (s, 1H), 7.02 (s, 1H), 12.95 (bs, 1H); m/z 253.27 (M-H) $^-$




The requisite methyl (3-isopropoxy-5-(t-butyloxycarbonyl)methoxy)benzoate was prepared

- 10 according to generic Alkylation Method B. The analytical data on all intermediates was consistent with the proposed structures.

EXAMPLE CC**3-amino-6-(3-isobutoxy-5-isopropoxy benzoyl) aminopyridine**


- 15 **(Route 7b)**



- To a solution of 2-(3-isobutoxy-5-isopropoxybenzoyl)amino-5-nitropyridine (1.74g, 4.66mmol) in ethanol (20ml) was added 10% Pd/C under an inert atmosphere. The reaction mixture was placed under a hydrogen atmosphere and stirred vigorously for 16h. The reaction mixture was flooded with argon, and then diluted with water (20ml) and acidified with 2M HCl (5ml). The suspension was filtered through celite, and the filtrate evaporated *in vacuo*.

The residue was partitioned between ethyl acetate (25ml) and saturated sodium bicarbonate (25ml), and the organic extract dried over MgSO_4 . Evaporation *in vacuo* afforded the title compound as a brown solid (1.30g, 81%).


$^1\text{H NMR } \delta$ (d_6 -DMSO): 0.97 (d, 6H), 1.26 (d, 6H), 2.00 (m, 1H), 3.78 (d, 2H), 4.69 (m, 1H), 5.12 (s, 2H), 6.58 (t, 1H), 6.99 (dd, 1H), 7.1 (ap d, 2H), 7.73-7.78 (m, 2H), 10.24 (bs, 1H); m/z 344.41 ($\text{M}+\text{H}$) $^+$



- 10 The requisite 2-(3-isobutoxy-5-isopropoxy) benzoyl amino-5-nitropyridine was prepared according to Route 1 (see Example 10 in Pyridine table); $^1\text{H NMR } \delta$ (d_6 -DMSO): 0.98 (d, 6H), 1.27 (d, 6H), 2.01 (m, 1H), 3.60 (d, 2H), 4.71 (m, 1H), 6.67 (ap t, 1H), 7.17 (ap d, 2H), 8.39 (d, 1H), 8.63 (dd, 1H), 9.20 (d, 1H), 11.43 (bs, 1H); m/z 374 ($\text{M}+\text{H}$) $^+$, 372 ($\text{M}-\text{H}$) $^-$.

15 EXAMPLE DD

2-[(3-isobutoxy-5-isopropoxy) benzoyl] amino -5-(N-methylsulfonyl)-carboxamido pyridine (Route 27)



- 20 2-[(3-isobutoxy-5-isopropoxy) benzoyl] aminopyridine-5-carboxylic acid (95mg, 0.255mmol) was stirred with EDC (59mg, 0.306mmol), DMAP (37mg, 0.306mmol) and methanesulfonamide (36mg, 0.378mmol) in DCM (3ml) under an inert atmosphere for 16h. The reaction mixture was diluted with further DCM (10ml) and extracted with water (2x5ml). 1M citric acid (5ml) and brine (5ml). Filtration through a PTFE membrane and evaporation *in vacuo* afforded the title compound as a colourless crystalline solid (90mg, 79%). $^1\text{H NMR } \delta$ (d_6 -DMSO): 0.97 (d, 6H), 1.26 (d, 6H), 2.03 (m, 1H), 3.01 (s, 3H), 3.79 (d, 2H), 4.70 (m, 1H),
- 25

- 86 -

6.63 (ap t, 1H), 7.14 (ap d, 2H), 7.70 (dd, 1H), 8.12 (d, 1H), 8.34 (ap d, 1H), (9.83, s, 1H), 10.81 (bs, 1H);
 m/z 422.37 (M+H)⁺, 420.30 (M-H)⁻


- 5 The requisite 2-[(3-isobutyloxy-5-isopropoxy) benzoyl] aminopyridine-5-carboxylic acid starting material was prepared from methyl 2-[(3-isobutyloxy-5-isopropoxy) benzoyl] aminopyridine-5-carboxylate by standard hydrolysis (Route 2a);

The requisite methyl 2-(3-isobutyloxy-5-isopropoxy) benzoyl aminopyridine-5-carboxylate
 10 was prepared by standard acid chloride coupling (Route 1);


EXAMPLE EE

2-[3-isopropoxy-5-[1-methyl-1-(5-carboxy-thiazol-2-yl aminocarbonyl)] ethoxy benzoyl] aminothiazole-5-carboxylic acid

15 (Route 28)




- Ethyl 2-[3-isopropoxy-5-[1-methyl-1-(5-ethoxycarbonyl-thiazol-2-yl aminocarbonyl)] ethoxy benzoyl] aminothiazole-5-carboxylate was hydrolysed by a standard method according to Example B Route 2a to give 2-[3-isopropoxy-5-[1-methyl-1-(5-carboxy-thiazol-2-yl aminocarbonyl)] ethoxy benzoyl] aminothiazole-5-carboxylic acid, ¹H NMR δ (d₆-DMSO):
 20 1.22 (d, 6H), 1.61 (s, 6H), 4.58-4.64 (m, 1H), 6.62 (s, 1H), 7.19 (s, 1H), 7.40 (s, 1H), 8.05 (s, 1H), 8.12 (s, 1H), m/z 533
 (M-H)⁻.



The requisite ethyl 2-{3-isopropoxy-5-[1-methyl-1-(5-ethoxycarbonyl-thiazol-2-yl)aminocarbonyl] ethoxy benzoyl} aminothiazole-5-carboxylate starting material was prepared by a standard acid chloride method according to Example A Route 1, starting from 3-isopropoxy-5-[(1-methyl-1-carboxy) ethoxy] benzoic acid.

5







The requisite 3-isopropoxy-5-[(1-methyl-1-carboxy) ethoxy] benzoic acid starting material was prepared according to the procedure described by Corey et al, JACS 91 p4782 (1969), starting from methyl 3-isopropoxy-5-hydroxy benzoate. The methyl ester was hydrolysed under the reaction conditions, and the product was isolated by extraction into aqueous sodium bicarbonate solution followed by acidification and extraction into ethyl acetate. The organic extracts were dried (MgSO₄), filtered and concentrated *in vacuo* to give the crude product as a pale yellow solid. Recrystallisation from hexane gave the title compound as a colourless solid; ¹H NMR δ (d₆-DMSO): 1.15 (d, 6H), 1.5 (s, 6H), 4.55 (hept, 1H), 6.55 (dd, 1H), 6.95 (m, 1H), 7.05 (m, 1H), 13.0 (br s, 1H); m/z 283 (M+H)⁺, 281 (M+H)⁺.

EXAMPLE FF:


By analogous methods to those described above the following pyridazine compounds, Example numbers FF₁ to FF₅, were also made.

20

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
1		1			¹ H NMR d (d ₆ -DMSO): 3.95 (3H, s); 5.25 (4H, s); 6.95 (1H, s); 7.4 (6H, m); 7.5 (2H, m); 7.65 (2H, m); 8.25 (1H, d); 8.6 (1H, d); 11.85 (1H, br s).

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
2		2	524/526	522	¹ H NMR d (d ₆ -DMSO): 2.0 (1H, s); 5.25 (4H, s); 6.95 (1H, s); 7.4 (6H, m); 7.5 (2H, m); 7.6 (2H, m); 8.25 (1H, d); 8.55 (1H, d); 11.8 (1H, br s). MS and NMR contained signals due to acid starting material (~20 mol%); NMR contained signals due to ethyl acetate, (~33 mol%)
3		1			¹ H NMR d (d ₆ -DMSO): 5.24 (4H, s); 6.93 (1H, m); 7.37 (6H, m); 7.50 (2H, m); 7.61 (2H, m); 7.71 (1H, dd); 8.36 (1H, d); 9.00 (1H, d).
4		2 *	524/526	522/524	¹ H NMR δ (d ₆ -DMSO): 5.2 (4H, s); 6.95 (1H, m); 7.15 (1H, s); 7.3 (1H, d); 7.4 (4H, m); 7.5 (2H, m); 7.6 (2H, m); 9.1 (2H, s); 11.35 (1H, br s); the spectrum also contains signals due to acid starting material (~40 mol%)
5		2a, 1c (c)	428		δ _H (300MHz, DMSO-d ₆) 1.29 (6H, d), 3.08 (2H, t), 4.30 (2H, t), 4.74 (1H, m), 6.73 (1H, s), 7.13 (1H, m), 7.24 (1H, s), 7.27 (1H, s), 7.34 (1H, m), 7.52 (1H, m), 8.25 (1H, d), 8.56 (1H, d), 11.75 (1H, s), 13.66 (1H, br s).

* For Example 15, the ester intermediate was prepared by route 1 and is exemplified as Example 12:


**EXAMPLE GG:**

5 By analogous methods to those described above the following compounds, Example numbers GG₁ to GG₇, were also made.

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
1		2 *			¹ H NMR d (d ₆ -DMSO): 5.22 (4H, s); 6.54 (1H, d); 6.93 (1H, d); 7.27 (1H, d); 7.32-7.44 (6H, m); 7.53 (2H, m); 7.63 (2H, m); 11.85 (1H, s); 12.86 (1H, br s).
2					¹ H NMR d (d ₆ -DMSO): 3.75 (3H, s); 5.21 (4H, s); 6.55 (1H, d); 6.86 (1H, m); 7.31 (1H, m); 7.38 (4H, m); 7.38 (2H, m); 7.56 (1H, m); 7.59 (2H, m); 10.80 (1H, br s).
3		1a	331		
4		1a	332.53	330.51	δ_H (300MHz, CDCl ₃) 1.02 (6H, d), 1.36 (6H, d), 2.08 (1H, m), 2.30 (3H, s), 3.75 (d, 2H), 4.60 (1H, hept), 6.66 (2H, m), 7.08 (2H, m), 9.85 (1H, br s).
5		1a	376.47	374.45	δ_H (300MHz, DMSO-d ₆) 0.98 (6H, d), 1.27 (6H, d), 2.02 (1H, m), 3.80 (2H, d), 3.84 (3H, s), 4.68 (1H, hept), 6.62 (1H, s), 7.12 (3H, m), 10.95 (1H, br s), 13.65 (1H, br s).

Example	Structure	Route	(M+H)+	(M-H)-	NMR
6		1b (HA TU)	386.47		δ_H (300MHz, $CDCl_3$) 1.35 (6H, d), 3.13 (2H, t), 3.72 (3H, s), 4.16 (2H, t), 4.53 (1H, hept), 6.60 (1H, s), 6.83 (1H, s), 7.00 (4H, m) 7.28 (2H, m), 8.98 (1H, br s).
7		1a	384		

* For GG₁, the ester intermediate was prepared by route 1:



5 1H NMR δ (d_6 -DMSO): 3.80 (3H, s); 5.23 (1H, m); 6.61 (1H, d); 6.95 (1H, s); 7.33-7.43 (7H, m); 7.50-7.55 (2H, m); 7.60-7.63 (2H, m); 11.90 (1H, br s).

EXAMPLE HH:

By analogous methods to those described above the following compounds, Example numbers







10 HH₁ to HH₃₃, were also made.









Example	Structure	Route	(M+H)+	(M-H)-	NMR
1		1	484		1H NMR δ (d_6 -DMSO): 5.26 (4H, s); 7.02 (1H, s); 7.40 (4H, m); 7.46 (2H, m); 7.54 (2H, m); 7.63 (2H, m); 9.24 (1H, s); 13.08 (1H, br s).


Example	Structure	Route	(M+ H)+	(M- H)-	NMR
2		1			¹ H NMR d (d6-DMSO): 2.63 (3H, s); 5.24 (4H, s); 6.96 (1H, s); 7.35-7.45 (6H, m); 7.51 (2H, m); 7.61 (2H, m); 12.84 (1H, br s).
3		1			¹ H NMR d (d6-DMSO): 1.38 (3H, t), 3.25 (2H, q); 5.25 (4H, s); 6.97 (1H, s); 7.41 (6H, m); 7.54 (2H, m); 7.64 (2H, m); 13.13 (1H, br s).
4		1			¹ H NMR d (d6-DMSO): 1.32 (3H, t), 4.32 (2H, q); 5.20 (4H, s); 6.78 (1H, s); 7.39 (4H, m); 7.46 (2H, m); 7.53 (2H, m); 7.64 (2H, m).
5		1			¹ H NMR d (d6-DMSO): 4.20 (3H, s); 5.28 (4H, s); 6.98 (1H, s); 7.42 (6H, m); 7.53 (2H, m); 7.62 (2H, m); 12.78 (1H, br s).
6		2	530, 532		¹ H NMR d (d6-DMSO): 5.24 (4H, s); 6.96 (1H, s); 7.37 (4H, m); 7.33 (2H, m); 7.53 (2H, m); 7.62 (2H, m).
7		1			¹ H NMR d (d6-DMSO): 5.25 (4H, s); 6.74 (1H, m); 6.99 (1H, s); 7.23 (1H, m); 7.41 (4H, m); 7.49 (2H, m); 7.53 (2H, m); 7.65 (2H, m); 7.97 (1H, s); 13.20 (1H, br s).

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
8		1			¹ H NMR d (d6-DMSO): 5.34 (4H, s); 7.03 (1H, s); 7.49 (2H, m); 7.57 (2H, m); 7.75 (4H, m); 7.91 (2H, d); 9.22 (1H, s); 13.06 (1H, br s).
9		19		564, 566	¹ H NMR d (d6-DMSO): 5.20 (4H, s); 6.68 (1H, m); 7.37 (4H, m); 7.45 (2H, m); 7.50 (2H, m); 7.62 (2H, m).
10		1	566	564, 566	¹ H NMR d (d6-DMSO): 5.22 (4H, s); 6.99 (1H, m); 7.39 (4H, m); 7.45 (2H, m); 7.51 (2H, m); 7.60 (2H, m); 13.34 (1H, br s).
11		1			¹ H NMR d (d6-DMSO): 2.33 (3H, s), 2.37 (3H, s); 3.25 (2H, m); 4.21 (2H, t); 5.14 (2H, s); 6.84 (1H, m); 7.22 (3H, m); 7.31 (1H, s); 7.40 (2H, m); 8.83 (1H, s); 9.21 (1H, s); 12.99 (1H, br s).
12		1			¹ H NMR d (d6-DMSO): 1.33 (3H, t); 2.32 (3H, s), 2.35 (3H, s); 3.22 (2H, m); 4.21 (2H, t); 4.40 (2H, q); 5.13 (2H, s); 6.87 (1H, m); 7.22 (3H, m); 7.33 (1H, m); 7.41 (2H, m); 8.82 (1H, s); 13.46 (1H, br s).

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
13				516, 518	¹ H NMR δ (d ₆ -DMSO): 5.21 (4H, s); 6.98 (1H, m); 7.34-7.40 (6H, m); 7.50 (2H, m); 7.59 (2H, m).
14			398		¹ H NMR δ (d ₆ -DMSO): 1.0 (d, 6H), 2.0 (hept, 1H), 2.35 (s, 3H), 3.8 (d, 2H), 5.2 (s, 2H), 6.85 (d, 1H), 7.15-7.25 (m, 3H), 7.30 (d, 1H), 7.4 (2H, m), 9.2 (s, 1H), 11.6 (br s, 1H).
15			402		¹ H NMR δ (d ₆ -DMSO): 1.0 (d, 6H), 2.0 (hept, 1H), 3.8 (d, 2H), 5.2 (s, 2H), 6.85 (s, 1H), 7.2-7.3 (m, 2H), 7.35 (s, 1H), 7.4 (m, 2H), 7.6 (t, 1H), 9.2 (s, 1H), 13.0 (br s, 1H).
16			350		
17			322		¹ H NMR δ (d ₆ -DMSO): 1.3 (d, 12H), 4.7 (hept, 2H), 6.65 (s, 1H), 7.25 (s, 2H), 9.2 (s, 1H), 12.95 (br s, 1H).
18					¹ H NMR δ (d ₆ -DMSO): 2.34 (s, 3H), 3.23 (t, 2H), 4.21 (t, 2H), 4.62 (d, 2H), 5.26 (d, 1H), 5.40 (d, 1H), 6.05 (m, 1H), 6.75 (s, 1H), 7.31 (s, 2H), 8.83 (s, 1H), 9.20 (s, 1H), 12.48 (br s, 1H).




Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
19					¹ H NMR δ (d ₆ -DMSO): 2.31 (s, 3H), 2.34 (s, 3H), 3.22 (t, 2H), 4.21 (t, 2H), 5.13 (s, 2H), 6.84 (s, 1H), 7.15-7.25 (m, 3H), 7.26 (1H, m), 7.39 (2H, m), 8.81 (s, 1H).
20					¹ H NMR δ (d ₆ -DMSO): 2.37 (s, 3H), 2.42 (s, 3H), 3.29 (t, 2H), 4.29 (t, 2H), 5.21 (s, 2H), 5.58 (s, 2H); 6.92 (s, 1H), 7.22-7.31 (m, 3H), 7.40 (1H, bs), 7.47 (2H, m), 8.90 (s, 1H). MS ES ⁺ 547.2, 549.1 (M+H) ⁺ .
21		19			¹ H NMR δ (d ₆ -DMSO): 2.35 (s, 3H), 2.93 (s, 6H), 3.22 (m, 2H), 4.19 (m, 2H), 6.41 (m, 1H), 6.98 (m, 1H), 7.06 (m, 1H), 8.80 (s, 1H), 9.17 (s, 1H).
22		19			¹ H NMR δ (d ₆ -DMSO): 2.58 (m, 6H), 3.43 (t, 2H), 4.37 (t, 2H), 4.50 (d, 2H), 6.41 (m, 1H), 6.61 (m, 1H), 7.16 (m, 2H), 7.34- 7.45 (m, 3H), 7.50 (m, 1H), 9.05 (s, 1H), 9.42 (s, 1H).
23		1	358		¹ H NMR δ (d ₆ -DMSO): 3.81 (s, 3H), 5.15 (s, 2H), 7.18 (t, 1H), 7.2-7.3 (m, 3H), 7.38 (d, 1H), 7.39- 7.43 (m, 1H), 7.55 (t, 1H), 12.27 (br s, 1H)
24		20	363	361	¹ H NMR δ (d ₆ -DMSO): 2.35 (s, 3H), 3.2 (t, 2H), 4.2 (t, 2H), 6.55 (m, 1H), 7.05 (s, 1H), 7.2 (s, 1H), 8.81 (s, 1H), 9.2 (s, 1H), 9.8 (br s, 1H).

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
25		1b	336		
26		1b	405		
27		2a, 1c (b)	388	386	δ_H (500MHz, DMSO- d_6) 1.27 (6H, d), 4.73 (1H, m), 5.21 (2H, s), 6.82 (1H, s), 7.20-7.31 (3H, br m), 7.36- 7.47 (2H, brm), 7.58 (1H, t), 9.23 (1H, s), 12.97 (1H, br s).
28		2a, 1c (b)	389		δ_H (500MHz, DMSO- d_6) 1.28 (6H, d), 3.06 (2H, t), 4.27 (2H, t), 4.72 (1H, m), 6.72 (1H, s), 7.12 (1H, d), 7.26 (1H, s), 7.31 (2H, m), 7.48 (1H, m), 9.20 (1H, s).
29		2a, 1a (d)	434		δ_H (300MHz, DMSO- d_6) 1.26 (6H, d), 3.07 (2H, t), 4.15 (2H, t), 4.70 (1H, m), 6.68 (1H, s), 7.11 (1H, d), 7.22 - 7.34 (3H, br m), 7.47 (1H, m).
30		1b (HA TU)	402 .42	400 .39	δ_H (300MHz, DMSO- d_6) 1.27 (6H, d), 2.63 (3H, s), 4.70 (1H, hept), 5.20 (2H, s), 6.82 (1H, s), 7.24 (3H, m), 7.39 (2H, m), 7.56 (1H, t), 12.80 (1H, br s).
31		1b (HA TU)	404 .40	402 .37	δ_H (300MHz, DMSO- d_6) 1.27 (6H, d), 2.63 (3H, s), 3.06 (2H, t), 4.25 (2H, t), 4.70 (1H, hept), 6.72 (1H, s), 7.12 (1H, d), 7.28 (3H, m), 7.47 (1H, m), 12.77 (1H, br s).
32		1b (HA TU)	468 .39	466 .37	δ_H (300MHz, DMSO- d_6) 2.63 (3H, s), 5.23 (4H, s), 6.97 (1H, s), 7.24 (4H, m), 7.43 (4H, m), 7.57 (2H, t), 12.84 (1H, br s).







Example	Structure	Route	(M+ H)+	(M- H)-	NMR
33		1a	336 .44	334 .40	

EXAMPLE II:







- By analogous methods to those described above the following compounds, Example numbers 5 II₁ to II₁₆₆, were also made. Some compounds were prepared by Route 1b (multi-parallel synthesis), as described in Example T. For compounds made by Route 2a (hydrolysis of esters), the requisite starting materials may be prepared by Route 1 or 1b.







Example	Structure	Route	(M+ H)+	(M- H)-	NMR
1		1	485, 487		¹ H NMR d (d6-DMSO): 5.24 (4H, s); 6.93 (1H, s); 7.26 (1H, d); 7.36-7.43 (6H, m); 7.50 (2H, m); 7.55 (1H, d); 7.61 (2H, m); 12.60 (1H, br s).
2		2a ****			¹ H NMR d (d6-DMSO): 5.25 (4H, s); 7.0 (1H, s); 7.4 (6H, m); 7.5 (2H, m); 7.6 (2H, m); 8.2 (1H, d).
3		1			¹ H NMR d (d6-DMSO): 3.62 (3H, s); 3.76 (2H, s); 5.24 (4H, s); 6.94 (1H, m); 7.06 (1H, s); 7.38-7.47 (6H, m); 7.54 (2H, m); 7.63 (2H, m); 12.69 (1H, br s).








Example	Structure	Route	(M+ H)+	(M- H)-	NMR
4		1		531	¹ H NMR d (d6-DMSO): 4.77 (2H, s); 5.25 (4H, s); 6.94 (1H, m); 7.31 (1H, s); 7.36-7.48 (6H, m); 7.53 (2H, m); 7.63 (2H, m); 12.83 (1H, br s) (+ 0.4 eq. iPr2NEt).
5		3	528, 530		¹ H NMR d (d6-DMSO): 2.63 (3H, m); 4.16 (2H, m); 5.24 (4H, s); 6.99 (1H, s); 7.38-7.44 (7H, m); 7.52 (2H, m); 7.62 (2H, m); 9.06 (1H, br s); 12.75 (1H, br s).
6		3			¹ H NMR d (d6-DMSO): 2.57 (3H, m); 3.48 (2H, m); 5.25 (4H, s); 6.95 (2H, m); 7.36-7.44 (6H, m); 7.53 (2H, m); 7.62 (2H, m); 7.83 (1H, m); 12.60 (1H, br s).
7		2a ****		497, 499 (- CO2)	¹ H NMR d (d6-DMSO): 3.64 (2H, s); 5.26 (4H, s); 6.95 (1H, s); 7.04 (1H, s); 7.37-7.46 (6H, m); 7.54 (2H, m); 7.63 (2H, m); 12.40 (1H, br s); 12.68 (1H, br s) (.HCl).
8		2a ****		459, 415 (- CO2)	¹ H NMR d (d6-DMSO): 5.15 (4H, s); 6.9 (1H, s); 7.2-7.5 (12H, m); 8.1 (1H, s).
9		1			¹ H NMR d (d6-DMSO): (iPr2NEt salt) 1.24 (15H, m); 3.12 (2H, m); 3.80 (2H, m); 5.24 (4H, s); 6.93 (1H, m); 7.36-7.45 (7H, m); 7.51 (2H, m); 7.61 (2H, m); 12.56 (1H, br s).

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
10		3			¹ H NMR d (d6-DMSO): 2.45 (4H, m); 3.55 (2H, s); 3.61 (4H, m); 5.29 (4H, s); 7.00 (1H, m); 7.11 (1H, s); 7.43-7.51 (6H, m); 7.58 (2H, m); 7.67 (2H, m); 12.66 (1H, br s).
11		4	550, 552		¹ H NMR d (d6-DMSO): 5.19 (2H, br s); 5.23 (4H, s); 6.72 (1H, dd); 6.93 (1H, m); 7.03 (1H, m); 7.35-7.44 (7H, m); 7.51 (2H, m); 7.61 (2H, m); 12.46 (1H, br s).
12		3	558, 560		¹ H NMR d (d6-DMSO): 2.60 (2H, t); 3.45 (2H, t); 3.72 (2H, s); 5.22 (4H, s); 6.91 (1H, m); 6.96 (1H, s); 7.35-7.30 (7H, m); 7.50 (2H, m); 7.60 (2H, m).
13		3	586, 588		¹ H NMR d (d6-DMSO): 3.11 (2H, q); 3.37 (2H, q); 3.50 (2H, s); 3.61 (1H, t); 5.22 (4H, s); 6.92 (2H, m); 7.34-7.42 (6H, m); 7.49 (2H, m); 7.60 (2H, m); 7.88 (1H, br s).
14		3	554, 556		¹ H NMR d (d6-DMSO): 0.29 (2H, m); 0.40 (2H, m); 2.16 (1H, m); 3.79 (2H, s); 5.27 (4H, s); 6.98 (2H, m); 7.40-7.48 (7H, m); 7.56 (2H, m); 7.66 (2H, m).
15		2b ****	366	364	¹ H NMR d (d6-DMSO): 7.05 (1H, d); 7.35 (1H, t); 7.45 (1H, dd); 7.6 - 7.75 (2H, m); 7.85 (1H, m); 7.9 - 8.0 (2H, m); 8.15 (1H, s); 13.1 (2H, br s).

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
16		6			¹ H NMR d (d6-DMSO): 2.68 (3H, s); 3.81 (1H, s); 5.15 (2H, s); 6.38 (1H, s); 6.87 (1H, s); 7.00 (1H, s); 7.37 (2H, m); 7.49 (1H, m); 7.58 (1H, m); 8.10 (1H, s); 8.21 (1H, s).
17		6			¹ H NMR d (d6-DMSO): 1.32 (6H, d); 4.88 (1H, m); 7.87 (1H, s); 8.05 (1H, s); 8.14 (1H, s); 8.45 (1H, s).
18		6		400, 402 (-CO2)	¹ H NMR d (d6-DMSO): 1.22 (6H, d); 4.36 (2H, m); 4.58 (1H, m); 6.24 (1H, s); 6.47 (1H, m); 6.84 (2H, m); 7.26 (3H, m); 7.37 (2H, m); 7.45 (1H, m); 7.76 (1H, br s).
19		6			¹ H NMR d (d6-DMSO): 1.21 (6H, d); 4.28 (2H, m); 4.55 (1H, m); 6.26 (1H, s); 6.43 (1H, m); 6.83 (1H, s); 6.89 (1H, s); 7.20 (1H, m); 7.26-7.37 (4H, m); 7.74 (1H, br s).
20		6		367 (-CO2)	¹ H NMR d (d6-DMSO): 1.23 (6H, d); 4.38 (2H, s); 4.60 (1H, m); 6.33 (1H, m); 6.89 (2H, m); 7.47 (1H, dd); 7.89 (1H, d); 8.10 (1H, s); 8.51 (1H, dd); 8.63 (1H, d).
21		6		396 (-CO2)	¹ H NMR d (d6-DMSO): 1.21 (6H, d); 3.81 (3H, s); 4.24 (2H, m); 4.55 (1H, m); 6.26 (2H, m); 6.84 (3H, m); 6.97 (1H, m); 7.20 (2H, m).









Example	Structure	Route	(M+ H)+	(M- H)-	NMR
22		6		464, 420 (- CO2)	
23		6			1H NMR d (d6-DMSO): 0.28 (2H, m); 0.52 (2H, m); 1.09 (1H, m); 1.32 (6H, d); 3.02 (2H, d); 4.69 (1H, m); 6.50 (1H, s); 6.99 (2H, s); 8.20 (1H, s).
24		6			1H NMR d (d6-DMSO): 1.24 (6H, d); 3.29 (2H, m); 3.56 (2H, t); 4.50 (2H, s); 4.58 (1H, m); 6.37 (1H, m); 6.85 (1H, s); 6.90 (1H, s); 7.26 (2H, m); 7.13 (3H, m); 8.10 (1H, s).
25		6		348	1H NMR d (d6-DMSO): 1.27 (6H, d); 2.96 (6H, s); 4.69 (1H, m); 6.39 (1H, m); 6.97 (1H, s); 7.04 (1H, s); 8.13 (1H, s); 12.89 (1H, br s).
26		2a ****	389, 391		1H NMR d (d6-DMSO): 5.21 (2H, s); 7.29-7.49 (6H, m); 7.74 (2H, s); 8.13 (1H, s); 13.1 (1H, br s).
27		1			1H NMR d (d6-DMSO): 2.31 (3H, s); 2.35 (3H, s); 3.22 (2H, t); 4.21 (2H, t); 5.12 (2H, s); 6.79 (1H, m); 7.18-7.28 (4H, m); 7.30 (1H, m); 7.54 (1H, d); 8.82 (1H, s); 12.48 (1H, br s).

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
28		1			¹ H NMR d (d ₆ -DMSO): 2.32 (3H, s); 2.37 (3H, s); 3.24 (2H, t); 4.22 (2H, t); 5.13 (2H, s); 6.80 (1H, m); 7.19 (3H, m); 7.29 (1H, s); 7.37-7.45 (3H, m); 9.06 (1H, s); 12.48 (1H, br s).
29		1			¹ H NMR d (d ₆ -DMSO): 1.28 (3H, t); 2.32 (3H, s); 2.37 (3H, s); 3.24 (2H, t); 4.14-4.29 (4H, m); 5.13 (2H, s); 6.84 (1H, m); 7.21 (4H, m); 7.29 (1H, s); 7.38 (2H, m); 8.20 (1H, s); 8.81 (1H, s).
30		2a (1)			¹ H NMR d (d ₆ -DMSO): 1.26 (d, 6H), 4.69 (m, 1H), 5.14 (s, 2H), 6.75 (s, 1H), 7.26-7.48 (m, 7H), 8.01 (s, 1H).
31		2a (1b)		391	¹ H NMR d (d ₆ -DMSO): 1.0 (d, 12H), 2.0 (m, 2H), 3.8 (d, 4H), 6.75 (s, 1H), 7.25 (d, 2H), 8.15 (s, 1H).
32		1			¹ H NMR δ (d ₆ -DMSO): 1.26 (d, 6H), 4.69 (m, 1H), 5.16 (s, 2H), 6.74 (s, 1H), 7.26 (d, 1H), 7.31-7.47 (m, 7H), 8.54 (d, 1H), 12.47 (bs, 1H).
33		2a (1)			¹ H NMR δ (d ₆ -DMSO): 1.26 (d, 6H), 2.38 (s, 3H), 4.69 (m, 1H), 5.18 (s, 2H), 6.31 (s, 1H), 6.76 (s, 1H), 7.30 (s, 1H), 7.35 (s, 1H), 8.00 (s, 1H).

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
34		2a (1)			¹ H NMR δ (d ₆ -DMSO): 1.26 (d, 6H), 2.39 (s, 3H), 4.70 (m, 1H), 5.20 (s, 2H), 6.31 (s, 1H), 6.79 (s, 1H), 7.27 (s, 1H), 7.32 (s, 1H), 8.12 (s, 1H).
35		1b	397		
36		1b	401		
37		1			¹ H NMR δ (d ₆ -DMSO): 1.27 (d, 6H), 2.39 (s, 3H), 4.69 (m, 1H), 5.18 (s, 2H), 6.31 (s, 1H), 6.76 (s, 1H), 7.26 (m, 2H), 7.32 (s, 1H), 8.53 (d, 1H).
36		2a (1)	379	377	¹ H NMR δ (d ₆ -DMSO): 12.98(bs, 1H), 8.12 (s, 1H), 7.24(s, 1H), 6.66(s, 1H), 4.70(m, 1H), 3.79 (d, 2H), 2.01 (m, 1H), 1.28 (d, 6H), 0.98 (d, 6H).
37		2a (1b)	365		¹ H NMR d (d ₆ -DMSO): 1.25 (d, 12H), 4.7 (hept, 2H), 6.65 (s, 1H), 7.2 (s, 2H), 8.15 (s, 1H).
38		2a (1)			¹ H NMR δ (d ₆ -DMSO): 2.64 (s, 3H), 5.16 (s, 4H), 6.90 (s, 1H), 7.29-7.47 (m, 7H), 7.53 (s, 1H), 8.03 (m, 1H), 12.90 (bs, 1H).

Example	Structure	Route	(M+ H) ⁺	(M- H) ⁻	NMR
39		2a (1)			¹ H NMR δ (d ₆ -DMSO): 2.64 (s, 3H), 5.17 (s, 4H), 6.93 (s, 1H), 7.29-7.45 (m, 7H), 7.53 (s, 1H), 8.13 (m, 1H).
40		1			¹ H NMR δ (d ₆ -DMSO): 2.64 (s, 3H), 5.14 (s, 4H), 6.90 (s, 1H), 7.26 (d, 1H), 7.31-7.47 (m, 7H), 7.49 (m, 1H), 7.55 (d, 1H), 12.56 (bs, 1H).
41		1b	349		
42		2a (1)			¹ H NMR δ (d ₆ -DMSO): 5.22 (s, 4H), 6.96 (s, 1H), 7.20-7.29 (m, 4H), 7.37- 7.44 (m, 4H), 7.55 (m, 2H), 8.12 (s, 1H).
43		2a (1)			¹ H NMR δ (d ₆ -DMSO): 5.21 (s, 4H), 6.93 (s, 1H), 7.19-7.29 (m, 4H), 7.38- 7.46 (m, 4H), 7.56 (m, 2H), 8.03 (s, 1H).
44		2a (1)			¹ H NMR δ (d ₆ -DMSO): 2.38 (s, 3H), 3.25 (t, 2H), 4.24 (t, 2H), 4.65 (d, 2H), 5.27 (d, 1H), 5.42 (d, 1H), 6.05 (m, 1H), 6.78 (s, 1H), 7.32 (s, 2H), 8.15 (s, 1H), 8.90 (s, 1H), 12.94 (br s, 1H).
45		2a (1)			¹ H NMR δ (d ₆ -DMSO): 2.32 (s, 3H), 2.34 (s, 3H), 3.22 (t, 2H), 4.21 (t, 2H), 5.13 (s, 2H), 6.82 (s, 1H), 7.16-7.25 (m, 3H), 7.30 (1H, s), 7.39 (1H, m), 7.98 (s, 1H), 8.81 (s, 1H).

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
46		2a *	419	417	¹ H NMR δ (d ₆ -DMSO): 3.8 (s, 3H), 5.3 (s, 2H), 7.15 (dd, 1H), 7.2-7.4 (m, 4H), 7.5 (d, 1H), 7.6 (d, 1H), 8.0 (s, 1H).
47		2a *	427	425	¹ H NMR δ (d ₆ -DMSO): 1.1 (d, 6H), 2.85 (hept, 1H), 3.75 (s, 3H), 5.2 (s, 2H), 7.0-7.3 (m, 6H), 7.4 (d, 1H), 8.0 (s, 1H).
48		2a **	405	403	¹ H NMR δ (d ₆ -DMSO): 2.34 (s, 3H), 3.20 (t, 2H), 4.13 (t, 2H), 6.43 (s, 1H), 6.92 (s, 1H), 6.97 (s, 1H), 8.09 (s, 1H), 8.83 (s, 1H), 12.75 (bs, 1H).
49		2a *			¹ H NMR δ (d ₆ -DMSO): 2.33 (s, 3H), 2.36 (2.36, 3H), 3.23 (t, 2H), 4.22 (t, 2H), 5.15 (s, 2H), 7.21 (s, 1H), 7.02-7.44 (m, 6H), 8.13 (s, 1H), 8.85 (s, 1H), 12.92 (bs, 1H).
50		6 **			¹ H NMR δ (d ₆ -DMSO): 2.32 (s, 3H), 2.34 (s, 3H), 3.19 (t, 2H), 4.12 (t, 2H), 4.25 (s, 2H), 6.37 (s, 1H), 6.92 (d, 2H), 7.08-7.21 (m, 3H), 7.25 (dd, 1H), 8.10 (s, 1H), 8.85 (s, 1H), 12.76 (bs, 1H).
51		6 **			
52		1			¹ H NMR δ (d ₆ -DMSO): 1.28 (t, 3H), 2.35 (s, 3H), 3.22 (t, 2H), 4.11 (t, 2H), 4.27 (q, 2H), 4.63 (d, 2H), 5.26 (dd, 1H), 5.39 (d, 1H), 6.04 (m, 1H), 6.76 (t, 1H), 7.28 (d, 2H), 8.21 (s, 1H), 8.81 (s, 1H), 13.02 (bs, 1H).
53		1b	261	259	¹ H NMR δ (CDCl ₃): 4.58 (d, 2H), 5.31 (dd, 1H), 5.45 (dd, 1H), 6.04 (m, 1H), 6.95 (d, 1H), 7.11 (d, 1H), 7.18 (m, 1H), 7.41 (t, 1H), 7.55 (m, 2H), 12.09 (br s, 1H).

Example	Structure	Route	(M+H)+	(M-H)-	NMR
54		2a	445		¹ H NMR δ (d ₆ -DMSO): 0.98 (d, 6H), 1.98 – 2.05 (m, 1H), 3.81 (d, 2H), 5.20 (s, 2H), 6.81 (s, 1H), 7.0-7.1 (m, 2H), 7.35 (s, 1H), 7.38-7.45 (m, 2H), 7.58 (t, 1H), 8.03 (s, 1H), 12.90 (br s, 1H).
55		2a	441		¹ H NMR δ (d ₆ -DMSO): 0.98 (d, 6H), 1.98-2.05 (m, 1H), 2.36 (s, 3H), 3.81 (d, 2H), 5.17 (s, 2H), 6.81 7.17-7.23 (m, 3H), 7.32 (s, 1H), 7.40 (ap d, 2H), 8.01 (s, 1H).
56		2a			¹ H NMR δ (d ₆ -DMSO): 1.27 (d, 6H), 4.71 (sept, 1H), 5.16 (d, 2H), 6.78 (d, 1H), 7.25-7.51 (m, 7H), 8.12 (s, 1H), 12.98 (bs, 1H).
57		2a	434	432	¹ H NMR δ (d ₆ -DMSO): 0.98 (d, 6H), 1.98-2.05 (m, 1H), 3.81 (d, 2H), 5.26 (s, 2H), 6.83 (ap t, 1H), 7.30 (s, 1H), 7.39 (s, 1H), 7.79 (s, 1H), 8.12 (s, 1H), 9.1 (s, 1H).
58		1b	335		
59		1b	293		
60		1			¹ H NMR δ (d ₆ -DMSO): 1.29 (d, 6H), 4.74 (sept, 1H), 5.22 (s, 2H), 6.79 (t, 1H), 7.19-7.32 (m, 4H), 7.37 (t, 1H), 7.43 (m, 1H), 7.56 (m, 2H), 12.61 (bs, 1H).
61		2a			¹ H NMR δ (d ₆ -DMSO) 1.26 (d, 6H), 4.64-4.76 (m, 1H), 5.20 (s, 2H), 6.78 (s, 1H), 7.18-7.34 (m, 3H), 7.36-7.46 (m, 2H), 7.50-7.60 (m, 1H), 7.98 (s, 1H).

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
62		2a			¹ H NMR δ (d ₆ -DMSO): 1.27 (d, 6H), 4.71 (m, 1H), 5.20 (s, 2H), 6.78-6.84 (m, 1H), 7.18-7.31 (m, 3H), 7.34-7.49 (m, 2H), 7.52-7.61 (m, 1H), 8.12 (s, 1H), 12.98 (bs, 1H)
63		2a	377		¹ H NMR δ (d ₆ -DMSO): 0.0-0.2 (m, 2H), 0.22-0.3 (m, 2H), 0.98 (d, 6H), 3.59 (d, 2H), 4.35-4.42 (m, 1H), 6.4 (s, 1H), 6.93 (s, 2H), 7.82 (s, 1H).
64		2a	403		¹ H NMR δ (d ₆ -DMSO): 1.29 (d, 6H), 4.78 (m, 1H), 4.86 (q, 2H), 6.89 (ap t, 1H), 7.36 (ap t, 2H), 8.17 (s, 1H), 13.05 (bs)
65		1 ***			¹ H NMR δ (d ₆ -DMSO): 1.29 (d, 6H), 4.72 (m, 1H), 5.19 (s, 2H), 6.88-6.97 (m, 1H), 7.09 (m, 1H), 7.16-7.26 (m, 4H), 7.54 (d, 1H), 7.61 (s, 1H), 7.70 (s, 1H), 12.05 (bs, 1H).
66		2a ***			¹ H NMR δ (d ₆ -DMSO): 1.29 (d, 6H), 4.74 (m, 1H), 5.18 (s, 2H), 6.87-6.97 (m, 1H), 7.11 (m, 1H), 7.16-7.26 (m, 3H), 7.63 (s, 1H), 7.71 (s, 1H), 8.11 (s, 1H).
67		2a ***			¹ H NMR δ (d ₆ -DMSO): 1.29 (d, 6H), 4.74 (m, 1H), 5.18 (s, 2H), 6.89-6.97 (m, 1H), 7.09 (m, 1H), 7.17-7.26 (m, 3H), 7.66 (s, 1H), 7.74 (s, 1H), 7.99 (s, 1H).
68		1b	457		
69		1b	404		
70		23			¹ H NMR δ (d ₆ -DMSO): 1.28 (d, 6H), 4.51 (s, 2H), 4.71 (m, 1H), 7.05 (s, 1H), 7.25 (d, 1H), 7.50 (s, 1H), 7.53 (d, 1H), 7.58 (s, 1H), 12.50 (bs, 1H).









Example	Structure	Route	(M+ H)+	(M- H)-	NMR
71		2a	405	403	¹ H NMR δ (d ₆ -DMSO): 1.14 (d, 6H), 1.3-1.4 (m, 2H), 1.42-1.62 (m, 4H), 1.65-1.82 (m, 2H), 3.9 (d, 2H), 4.62-4.78 (m, 1H), 6.68 (s, 1H), 7.22 (s, 2H), 8.12 (s, 1H).
72		2a	381	379	¹ H NMR δ (d ₆ -DMSO): 1.25 (d, 6H), 3.3 (s, 3H), 3.7 (t, 2H), 4.15 (t, 2H), 4.6-4.8 (hept, 1H), 6.75 (t, 1H), 7.25 (d, 2H), 8.15 (s, 1H), 13.0 (bs, 2H).
73		2a	379	377	¹ H NMR δ (d ₆ -DMSO): 3.85 (s, 3H), 5.25 (s, 2H), 6.9 (m, 1H), 7.2-7.35 (m, 3H), 7.4-7.5 (m, 2H), 7.6-7.7 (t of d, 1H), 8.15 (s, 1H), 13.0 (bs, 2H).
74		2a	401		¹ H NMR δ (d ₆ -DMSO): 0.9 (t, 3H), 1.2-1.3 (d, 3H + d, 6H), 1.5-1.75 (m, 2H), 4.45 (hex, 1H), 4.75 (hept, 1H), 6.7 (t, 1H), 7.2 (d, 2H), 8.15 (s, 1H), 13.0 (bs, 2H).
75		22			¹ H NMR δ (d ₆ -DMSO): 1.31 (d, 6H), 4.82 (m, 1H), 7.26 (d, 1H), 7.56 (d, 1H), 7.59 (s, 1H), 7.94 (d, 1H), 8.15 (s, 1H), 10.00 (s, 1H), 12.77 (bs, 1H).
76		2a			¹ H NMR δ (d ₆ -DMSO): 0.97 (d, 3H), 1.26 (s, 6H), 1.72 (t, 2H), 3.85-4.20 (m, 2H), 4.56-4.83 (m, 1H), 6.69 (s, 1H), 7.00 (s, 1H), 7.26 (s, 1H), 8.11 (s, 1H).
77		2a	359		¹ H NMR δ (d ₆ -DMSO): 1.30 (d, 6H), 3.30 (s, 1H), 4.74 (m, 1H), 4.88 (s, 2H), 6.80 (s, 1H), 7.31 (app d, 2H), 8.15 (s, 1H), 10.01 (bs, 1H).
78		2a	407	405	¹ H NMR δ (d ₆ -DMSO): 0.91 (t, 6H), 1.29 (d, 6H), 1.37-1.53 (m, 4H), 1.56-1.70 (m, 1H), 3.30 (d, 2H), 4.73 (m, 1H), 6.72 (s, 1H), 7.26 (app d, 2H), 8.14 (s, 1H), 13.00 (bs, 1H).









Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
79		1	378		¹ H NMR δ (d ₆ -DMSO): 0.98 (d, 6H), 1.28 (d, 6H), 2.02 (m, 1H), 3.80 (d, 2H), 4.65 (m, 1H), 6.75 (ap t, 1H), 7.25 (ap d, 2H), 8.68 (s, 1H)
80		28	533		¹ H NMR δ (d ₆ -DMSO): 1.22 (d, 6H), 1.61 (s, 6H), 4.58-4.64 (m, 1H), 6.62 (s, 1H), 7.19 (s, 1H), 7.40 (s, 1H), 8.05 (s, 1H), 8.12 (s, 1H).
81		2a			¹ H NMR δ (d ₆ -DMSO): 1.29 (d, 6H), 4.50 (m, 2H), 4.71 (m, 1H), 5.26 (bs, 1H), 7.08 (s, 1H), 7.53 (s, 1H), 7.60 (s, 1H), 8.01 (s, 1H), 13.00 (bs, 1H).
82		21			¹ H NMR δ (d ₆ -DMSO): 1.32 (d, 6H), 2.80 (s, 3H), 3.37-3.63 (m, 4H), 3.95-4.10 (m, 4H), 4.39 (m, 2H), 4.76 (m, 1H), 7.29 (d, 1H), 7.53 (m, 3H), 7.68 (s, 1H), 7.79 (s, 1H), 12.77 (bs, 1H).
83		21			¹ H NMR δ (d ₆ -DMSO): 1.31 (d, 6H), 2.71 (s, 6H), 4.26 (m, 2H), 4.76 (m, 1H), 7.29 (d, 1H), 7.42 (m, 1H), 7.55 (d, 1H), 7.70 (s, 1H), 10.66 (bs, 1H).
84		21			¹ H NMR δ (d ₆ -DMSO): 1.31 (d, 6H), 3.03-3.16 (m, 4H), 3.71-3.95 (m, 4H), 4.34 (m, 2H), 4.77 (m, 1H), 7.47 (m, 1H), 7.72 (m, 2H), 8.13 (s, 1H).
85		21			¹ H NMR δ (d ₆ -DMSO): 0.41 (m, 2H), 0.60 (m, 2H), 1.14 (m, 1H), 1.35 (d, 6H), 2.85 (m, 2H), 4.19 (m, 2H), 4.81 (m, 1H), 7.32 (d, 1H), 7.46 (s, 1H), 7.60 (d, 1H), 7.72 (s, 1H), 7.80 (s, 1H), 9.35 (bs, 2H).
86		21			¹ H NMR δ (d ₆ -DMSO): 1.27(m, 12H), 3.26 (m, 2H), 4.14 (m, 2H), 4.76 (m, 1H), 7.26 (d, 1H), 7.45 (s, 1H), 7.55 (d, 1H), 7.68 (s, 1H), 7.76 (s, 1H), 9.18 (bs, 2H).










Example	Structure	Route	(M+ H) ⁺	(M- H) ⁻	NMR
87		21			¹ H NMR δ (d ₆ -DMSO): 0.72 (m, 2H), 0.89 (m, 2H), 1.32 (d, 6H), 2.66 (m, 1H), 4.21 (m, 2H), 4.75 (m, 1H), 7.26 (d, 1H), 7.42 (s, 1H), 7.55 (d, 1H), 7.68 (s, 1H), 7.76 (s, 1H), 9.53 (bs, 2H).
88		1 (See Ex 26)	351	349	¹ H NMR δ (d ₆ -DMSO): 1.27 (d, 6H), 3.70 (s, 3H), 4.71 (m, 1H), 4.86 (s, 2H), 6.99 (t, 1H), 7.23 (t, 1H), 7.26-7.27 (m, 2H), 12.53 (s, 1H).
89		24			¹ H NMR δ (d ₆ -DMSO): 1.32 (d, 6H), 4.79 (m, 1H), 7.62 (m, 1H), 7.92 (m, 1H), 8.13 (s, 1H), 8.18 (s, 1H), 10.03 (s, 1H).
90		26	419	417	¹ H NMR δ (d ₆ -DMSO): 1.28 (d, 6H), 2.18 (s, 3H), 2.24 (m, 2H), 2.32 (m, 2H), 3.44 (ap t, 4H), 4.65 (m, 1H), 4.85 (s, 2H), 6.68 (ap t, 1H), 7.19 (m, 1H), 7.24 (ap d, 2H), 7.55 (ap d, 1H), 12.45 (bs, 1H).
91		25			¹ H NMR δ (d ₆ -DMSO): 1.01 (d, 6H), 1.29 (d, 6H), 2.81 (m, 1H), 4.72 (m, 1H), 6.53 (dd, 1H), 6.29 (d, 1H), 6.97 (s, 1H), 7.50 (s, 1H), 7.53 (s, 1H), 8.11 (s, 1H), 8.18 (s, 1H).
92		1			¹ H NMR δ (d ₆ -DMSO): 1.29 (d, 9H), 4.28 (q, 2H), 4.53 (d, 2H), 4.71 (m, 1H), 5.26 (t, 1H (-OH)), 7.10 (s, 1H), 7.53 (s, 1H), 7.60 (s, 1H), 8.20 (s, 1H), 13.01 (bs, 1H).
93		1			¹ H NMR δ (d ₆ -DMSO): 1.34 (d, 3H), 1.39 (m, 6H), 4.30 (q, 2H), 4.84 (m, 1H), 7.58 (s, 1H), 7.97 (s, 1H), 8.17 (s, 1H), 8.26 (s, 1H), 10.09 (s, 1H).
94		1a	307		









Example	Structure	Route	(M+ H)+	(M- H)-	NMR
95		1a	307		
96		2a, 1c	389	387	δ_H (300MHz, DMSO- d_6) - 0.04-0.06 (4H,m); 0.22-0.35 (4H,m); 0.85-1.05 (2H,m); 3.54-4.64 (4H,d); 6.44 (1H, m); 6.93 (6.93-6.97 (2H, m); 7.84 (1H, s)
97		1b (HA TU)	389 .38	387 .34	δ_H (300MHz, DMSO- d_6) 1.30 (6H, d), 3.08 (2H, t), 4.25 (2H, t), 4.73 (1H, hept), 6.70 (1H, s), 7.14 (1H, d), 7.3 (4H, m), 7.48 (1H, m), 7.57 (1H, d), 12.55 (1H, br s).
98		1a	349		
99		1b (HA TU)	374 .43	372 .39	δ_H (300MHz, DMSO- d_6) 0.98 (6H, d), 1.27 (6H, d), 2.00 (1H, m), 3.80 (2H, d), 4.24 (2H, s), 4.70 (1H, hept), 6.66 (1H, t), 7.23 (2H, d), 7.46 (1H, s), 12.59 (1H, br s).
100		1a	401		
101		1a	415		
102		3 (e) (CM 1a)	395 .19	393 .19	δ_H (300MHz, CDCl ₃) 1.02 (6H, d), 1.35 (6H, d), 2.08 (4H, m), 3.74 (4H, m), 4.60 (1H, hept), 6.64 (1H, m), 6.78 (1H, s), 7.00 (1H, m).









Example	Structure	Route	(M+ H) ⁺	(M- H) ⁻	NMR
103		3 (e) (CM 1a)	393 .22	391 .21	δ_H (300MHz, $CDCl_3$) 1.02 (6H, d), 1.26 (3H, t), 1.35 (6H, d), 2.08 (1H, m), 3.60 (2H, q), 3.74 (d, 2H), 4.47 (2H, s), 4.58 (1H, hept), 6.64 (1H, m), 6.88 (1H, s), 7.02 (1H, m).
104			411 .42	409 .38	δ_H (300MHz, $DMSO-d_6$) 0.98 (6H, d), 1.27 (6H, d), 2.02 (1H, m), 2.55 (3H, s), 3.80 (2H, d), 4.14 (2H, s), 4.70 (1H, hept), 6.66 (1H, s), 7.23 (3H, m), 12.62 (1H, br s).
105		1a	427 .39	425 .38	δ_H (300MHz, $CDCl_3$) 1.02 (6H, d), 1.36 (6H, d), 2.08 (1H, m), 3.75 (2H, d), 4.60 (1H, hept), 6.68 (1H, m), 7.00 (2H, m), 7.69 (1H, s).
106		1b (HA TU)	349 .45	347 .43	δ_H (300MHz, $CDCl_3$) 0.95 (6H, d), 1.25 (6H, d), 1.95-2.05 (1H, m), 2.2 (3H, s), 3.65 (2H, d), 6.7 (1H, m), 6.98 (1H, m), 7.02 (1H, m).
107		1b (HA TU)	403 .39	401 .37	δ_H (300MHz, $DMSO-d_6$) 1.25 (6H, d), 2.38 (3H, s), 3.05 (2H, t), 4.6-4.8 (1H, m), 7.05 (1H, d), 7.10-7.12 (3H, m), 7.15, (1H, m), 7.42-7.45 (1H, m).
108		1b (HA TU)	401 .42	399 .39	δ_H (300MHz, $CDCl_3$) 1.25 (6H, d), 2.3 (3H, s), 4.4-4.6 (1H, m), 5.05 (2H, s), 6.65 (1H, m), 6.85 (1H, s), 7.0-7.15 (4H, m), 7.2-7.3 (1H, m), 7.38-7.42 (1H, m).
109		1b (HA TU)	467 .38	465 .37	δ_H (300MHz, $DMSO-d_6$) 2.35 (3H, s), 5.2 (4H, s), 6.95 (1H, s), 7.2-7.3 (5H, m), 7.4-7.45 (4H, m), 7.5-7.6 (2H, m).
110		1b (HA TU)	467 .37	465 .38	δ_H (300MHz, $CDCl_3$) 1.9 (3H, s), 4.95 (4H, s), 6.4 (1H, s), 6.9-7.1 (6H, m), 7.15-7.25 (2H, m), 7.3-7.4 (2H, m).









Example	Structure	Route	(M+ H) ⁺	(M- H) ⁻	NMR
111		2a, 1a	433	431	δ_H (500MHz, DMSO- d_6) 1.27 (6H, d), 3.06 (2H, t), 4.25 (2H, t), 4.72 (1H, m), 6.71 (1H, s), 7.12 (1H, d), 7.23-7.32 (3H, br m), 7.46 (1H, m), 8.10 (1H, s).
112		2a, 1a	433	431	δ_H (500MHz, DMSO- d_6) 1.28 (6H, d), 3.06 (2H, t), 4.24 (2H, t), 4.72 (1H, m), 6.69 (1H, s), 7.12 (1H, d), 7.27 (1H, s), 7.31 (2H, s), 7.47 (1H, m), 8.02 (1H, s).
113		21	439 .44	437 .39	δ_H (300MHz, DMSO- d_6) 1.25 (6H, d), 3.0-3.2 (2H, m), 3.3-3.55 (4H, m), 4.3-4.5 (4H, m), 4.75-4.85 (1H, m), 7.25 (1H, d), 7.55-7.6 (2H, m), 7.65 (1H, s), 7.75 (1H, s), 7.95 (1H, s), 8.1 (1H, s), 8.4 (1H, s).
114		3	430 .40	428 .38	δ_H (300MHz, CDCl ₃) 1.25 (6H, d), 2.42 (3H, s), 3.82 (2H, s), 4.45-4.6 (1H, m), 5.05 (2H, s), 6.6 (1H, s), 6.95-7.15 (3H, m), 7.2-7.25 (2H, m), 7.35-7.45 (1H, m).
115		3	474 .42	472 .40	
116		21	419 .47	417 .44	δ_H (300MHz, DMSO- d_6) 1.25 (6H, d), 3.25 (3H, s), 3.3-3.75 (12H, m), 4.3-4.45 (2H, m), 4.75-4.8 (1H, m), 7.25 (1H, d), 7.5-7.6 (2H, m), 7.7 (1H, s), 7.8 (1H, s).
117		21	453 .39	451 .37	
118		3	458 .39	456 .42	

Example	Structure	Route	(M+ H) ⁺	(M- H) ⁻	NMR
119		21	495 .43		δ_H (300MHz, DMSO- d_6) 1.25 (6H, d), 3.3-3.65 (8H, m), 4.2-4.5 (2H, m), 4.7-4.8 (1H, m), 6.05 (2H, s), 6.95 (1H, d), 7.05 (1H, d), 7.25 (2H, m), 7.55 (2H, m), 7.7 (1H, s), 7.8 (1H, s).
120		3	490 .43	488 .42	
121		3	470 .48	468 .47	
122		3	488 .49	486 .47	
123		3	486 .51	484 .51	
124		21	467 .50	465 .49	
125		21	455 .48 453 .46		
126		21	467 .50	465 .48	








Example	Structure	Route	(M+ H)+	(M- H)-	NMR
127		21	453 .49	451 .47	
128		21	459 .49	457 .47	
129		21	390 .51	388 .47	
130		21	446 .51	444 .49	
131		21	431 .55	429 .51	
132		1b (HA TU)	401 .37	399 .33	δ_H (300MHz, DMSO- d_6) 2.08 (3H, s), 5.12 (2H, s), 5.24 (2H, s), 7.23 (4H, m), 7.42 (1H, m), 7.56 (2H, m), 7.68 (1H, s), 7.76 (1H, s), 12.64 (1H, br s).
133		2a (f)	359 .43	357 .39	δ_H (300MHz, DMSO- d_6) 4.55 (2H, d), 5.23 (2H, s), 7.23 (4H, m), 7.42 (1H, m), 7.56 (2H, m), 7.68 (2H, m), 12.56 (1H, br s).
134		3	474 .48	472 .47	
135		3	460 .46	458 .43	

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
136		3	458 .48	456 .47	
137		3	472 .51	470 .49	
138		3	488 .51	486 .52	
139		3	486 .49	484 .47	
140		3	486 .50	484 .49	
141		3	444 .45	442 .41	
142		21	441 .43	439 .42	δ_H (300MHz, DMSO- d_6) 2.82 (3H, s), 3.49 (8H, m), 4.54 (1H, d), 5.24 (3H, m), 7.30 (3H, m), 7.45 (2H, m), 7.59 (2H, m), 7.81 (2H, m), 12.65 (1H, br s).
143		21	505 .45	503 .38	δ_H (300MHz, DMSO- d_6) 3.15 (2H, m), 3.45 (2H, m), 4.25 (4H, m), 4.52 (1H, d), 5.25 (3H, m), 7.27 (3H, m), 7.45 (1H, m), 7.62 (3H, m), 7.90 (3H, m), 8.16 (1H, s), 8.42 (1H, s), 12.70 (1H, br s).

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
144		21	521 .43		δ_H (300MHz, DMSO- d_6) 3.33 (8H, m), 4.52 (1H, d), 5.27 (3H, m), 7.03 (5H, m), 7.28 (3H, m), 7.45 (1H, m), 7.65 (3H, m), 7.89 (1H, m), 9.20 (1H, br s), 12.69 (1H, br s).
145		21	361 .50	359 .46	δ_H (300MHz, CDCl ₃) 1.36 (6H, d), 2.56 (4H, m), 3.04 (4H, m), 3.53 (2H, s), 4.61 (1H, hept), 6.95 (1H, d), 7.07 (1H, m), 7.24 (1H, m), 7.44 (2H, m).
146		21	382 .12	380 .13	1H NMR δ (CDCl ₃): 1.37 (d, 6H), 2.3 (m, 2H), 2.7 (m, 2H), 2.7 (m, 2H), 2.85 (m, 2H), 4.6 (m, 1H), 6.95 (m, 1H), 7.1 (m, 1H), 7.2 (m, 1H), 7.4 (m, 2H)
147		21	396 .45	394 .4	1H NMR δ (CDCl ₃): 1.37 (d, 6H), 1.95 (m, 4H), 2.5 (m, 4H), 3.55 (s, 2H), 4.6 (m, 1H), 7.0 (d, 1H), 7.1 (m, 1H), 7.6 (m, 1H)
148		1b (HAT U)	382 .12	380 .13	1H NMR δ (CDCl ₃): 1.37 (d, 6H), 2.3 (m, 2H), 2.7 (m, 2H), 2.7 (m, 2H), 2.85 (m, 2H), 4.6 (m, 1H), 6.95 (m, 1H), 7.1 (m, 1H), 7.2 (m, 1H), 7.4 (m, 2H)
149		1b (HAT U)	403 .39	401 .36	δ_H (300MHz, DMSO- d_6) 2.09 (3H, s), 3.26 (2H, t), 4.30 (2H, t), 5.08 (2H, s), 6.98 (2H, m), 7.17 (1H, s), 7.26 (1H, d), 7.35 (1H, m), 7.54 (1H, d), 7.64 (2H, br s), 12.62 (1H, br s).
150		2a (g)	361 .41	359 .38	
151		3	432 .40	430 .37	

Example	Structure	Route	(M+ H)+	(M- H)-	NMR
152		3	476 .48	474 .47	
153		3	472 .48	470 .45	
154		3	462 .45	460 .43	
155		21	462 .41	460 .38	
156		21	521 .42	519 .40	
157		21	507 .48		
158		21	453 .52	451 .49	¹ H NMR δ (CDCl ₃): 1.35 (d, 6H), 2.5 ((m, 2H), 3.65 (m, 4H), 4.65 (m, 1H), 6.3 (d, 1H), 6.95 (d, 1H), 7.1 (m, 1H), 7.35 (d, 1H), 7.5 (m, 1H), 7.58 (s, 1H), 8.1 (d, 1H)
159		21	461 .49	459 .48	

- 118 -

Example	Structure	Route	(M+ H) ⁺	(M- H) ⁻	NMR
160		1b (HAT U) (h)	453 .44	451 .40	
161		1b (HAT U) (h)	406 .40		
162		21 (i)	467 .50	465 .49	
163		21 (i)	506 .47	504 .46	
164		21 (i)	505 .46	503 .43	
165		21 (i)	541 .39	539 .35	
166		21	429 .54	427 .51	


Notes:

* Final products prepared by hydrolysis method 2a; requisite starting materials prepared according to generic alkylation methodology followed by coupling (Route 1).


** Final products prepared by reductive amination method 6 method ; requisite starting materials prepared according to generic alkylation methodology followed by coupling (Route 1) and hydrolysis (Route 2a).

*** Final products prepared by hydrolysis (Route 2a) or acid chloride coupling (Route 1); requisite starting materials prepared according to generic alkylation methodology followed by coupling (Route 1).

5 **** For Examples II₂, II₇, II₈, II₁₅ and II₂₆, the ester intermediates were prepared by route 1:




¹H NMR δ (d₆-DMSO): 1.3 (3H, t); 4.3 (2H, q); 5.25 (4H, s); 7.0 (1H, t); 7.4 (6H, m); 7.5 (2H, m); 7.6 (2H, m); 8.2 (1H, s).




exemplified as Example II₃.

10




¹H NMR δ (d₆-DMSO): 1.3 (3H, t); 4.3 (2H, q); 5.2 (4H, s); 6.95 (1H, t); 7.2-7.5 (12H, m); 8.2 (1H, s); 13.05 (1H, br s); the spectrum also contains signals due to trace amounts of 2-aminothiazole.

15



not characterised.














20 MH⁺ = 389, 391
M-H = 387, 389






EXAMPLE JJ:

By analogous methods to those described above the following compounds, Example numbers








JJ₁ to JJ₅₇, were also made.










Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
1		7	426.19	424.25	¹ H NMR d (d ₆ -DMSO): 5.17 (m, 6H) 6.80 (s, 1H), 7.00 (d, 1H), 7.26 to 7.46 (m, 12H), 7.71 (s, 1H), 7.78 (d, 1H), 10.28 (br s, 1H)
2		8		552.22	¹ H NMR d (d ₆ -DMSO): 1.55 (s, 6H), 2.08 (s, 3H), 5.18 (s, 4H), 6.85 (s, 1H), 7.29 to 7.50 (m, 12H), 7.98 (dd, 1H), 8.13 (d, 1H), 8.61 (s, 1H), 9.70 (s, 1H), 10.72 (s, 1H)
3		9	512.16	510.22	¹ H NMR d (d ₆ -DMSO): 1.35 (s, 6H), 5.18 (s, 4H), 6.88 (s, 1H), 7.28 to 7.48 (m, 12H), 8.08 (d, 1H), 8.22 (d, 1H), 8.82 (s, 1H), 9.90 (s, 1H), 10.96 (s, 1H)
4		8		502.49	¹ H NMR d (d ₆ -DMSO): 3.02 (s, 3H), 5.17 (s, 4H), 6.86 (s, 1H), 7.29 to 7.58 (m, 12H), 7.70 (d, 1H), 8.13 (d, 1H), 8.24 (s, 1H), 9.83 (s, 1H), 10.83 (s, 1H)
5		8	526.41	524.45	¹ H NMR d (d ₆ -DMSO): 2.13 (s, 3H), 4.65 (s, 2H), 5.18 (s, 4H), 6.84 (s, 1H), 7.27 to 7.48 (m, 12H), 7.96 (d, 1H), 8.13 (d, 1H), 8.61 (s, 1H), 10.24 (s, 1H), 10.73 (s, 1H)

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
6		8	498.55	496.55	¹ H NMR d (d6-DMSO): 3.39 (s, 3H), 4.01 (s, 1H), 5.18, (s, 4H), 6.85 (s, 1H), 7.28 to 7.50 (m, 12H), 8.07 (m, 2H), 8.67 (s, 1H), 9.95 (s, 1H), 10.71 (s, 1H)
7		8	540.58	538.63	¹ H NMR d (d6-DMSO): 1.20 (t, 3H), 3.47 (s, 2H), 4.11 (q, 2H), 5.17 (s, 4H), 6.83 (s, 1H), 7.28 to 7.48 (m, 12H), 7.95 (d, 1H), 8.13 (d, 1H), 8.60 (s, 1H), 10.35 (s, 1H), 10.73 (s, 1H)
8		8	526.53	524.61	¹ H NMR d (d6-DMSO): 1.30 (t, 3H), 4.30 (q, 2H), 5.17 (s, 4H), 6.86 (s, 1H), 7.28 to 7.50 (m, 12H), 8.14 (s, 2H), 8.74 (s, 1H), 10.78 (s, 1H), 10.97 (s, 1H)
9		10	525.61	523.66	¹ H NMR d (d6-DMSO): 1.30 (s, 9H), 5.18 (s, 4H), 6.09 (s, 1H), 6.85 (s, 1H), 7.32-7.50 (m, 12H), 7.78 (dd, 1H), 8.04 (d, 1H), 8.38 (s, 1H), 8.44 (s, 1H), 10.65 (s, 1H)
10		9	512.4		¹ H NMR d (d6-DMSO): 3.41 (s, 2H), 5.17 (s, 4H), 6.90 (s, 1H), 7.29 to 7.54 (m, 12H), 8.03 (d, 1H), 8.13 (d, 1H), 8.70 (s, 1H), 10.50 (s, 1H), 10.85 (s, 1H)
11		9	484.4		¹ H NMR d (d6-DMSO): 4.04 (s, 2H), 5.20 (s, 4H), 6.89 (s, 1H), 7.30 to 7.51 (m, 12H), 8.12 (d, 1H), 8.22 (d, 1H), 8.81 (s, 1H), 10.05 (s, 1H), 11.00 (s, 1H)







Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
12		7	476.36		¹ H NMR d (d6-DMSO): 5.14 (s, 2H), 5.32 (s, 4H), 6.90 (s, 1H), 7.01 (dd, 1H), 7.35 (s, 2H), 7.59 (m, 2H), 7.80 (m, 6H), 7.90 (d, 2H), 10.38 (s, 1H) + 0.1 EtOAc
13		8	604.29	602.3	¹ H NMR d (d6-DMSO): 1.55 (s, 6H), 2.07 (s, 3H), 5.33 (s, 4H), 6.95 (s, 1H), 7.40 (s, 2H), 7.56 (m, 2H), 7.73 (m, 4H), 7.90 (d, 2H), 7.98 (dd, 1H), 8.13 (d, 1H), 8.63 (s, 1H), 9.71 (s, 1H), 10.82 (s, 1H)
14		9	562.28	560.27	¹ H NMR d (d6-DMSO): 1.34 (s, 6H), 5.32 (s, 4H), 6.97 (s, 1H), 7.40 (s, 2H), 7.57 (m, 2H), 7.75 (m, 4H), 7.90 (d, 2H), 8.09 (d, 1H), 8.21 (dd, 1H), 8.82 (s, 1H), 9.90 (s, 1H), 10.99 (s, 1H)
15		11	534.41		¹ H NMR d (d6-DMSO): 3.22 (t, 2H), 3.28 (t, 3H), 3.50 (t, 2H), 5.31 (s, 4H), 6.92 (s, 1H), 7.12 (dd, 1H), 7.34 (s, 2H), 7.57 (m, 2H), 7.75 (m, 5H), 7.82 (d, 1H), 7.91 (d, 2H), 10.49 (br s, 1H)
16		11	547.86		¹ H NMR d (d6-DMSO): 2.20 (s, 6H), 3.12 (m, 2H), 5.32 (s, 4H), 5.51 (br s, 1H), 6.89 (s, 1H), 7.06 (dd, 1H), 7.37 (s, 2H), 7.57 (m, 2H), 7.74 (m, 5H), 7.83 (d, 1H), 7.92 (d, 2H), 10.41 (s, 1H), and 2H under DMSO or water







Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
17		11	504.54		¹ H NMR d (d6-DMSO): 1.15 (t, 3H), 3.06 (quartet, 2H), 5.32 (s, 4H), 6.90 (s, 1H), 7.00 (dd, 1H), 7.35 (s, 2H), 7.57 (m, 2H), 7.73 (m, 5H), 7.85 (d, 1H), 7.92 (d, 2H), 10.41 (s, 1H)
18		12	485.5	483.5	¹ H NMR d (d6-DMSO): 5.13 (s, 2H), 5.18 (s, 2H), 5.31 (s, 1H), 6.88 (s, 1H), 7.00 (dd, 1H), 7.32 (s, 2H), 7.40 (m, 3H), 7.50 (s, 1H), 7.58 (m, 1H), 7.74 (m, 3H), 7.80 (d, 1H), 7.90 (d, 1H), 10.33 (s, 1H)
19		1	493, 495		¹ H NMR d (d6-DMSO): 2.35 (3H, s); 5.31 (4H, s); 6.98 (1H, t); 7.43-7.48 (6H, m); 7.58-7.61 (2H, m); 7.65-7.71 (3H, m); 8.14 (1H, d); 8.29 (1H, s); 10.84 (1H, s)
20		13	525		¹ H NMR d (d6-DMSO): 3.10 (2H, m); 3.30 (6H, s); 3.60 (2H, m); 5.19 (4H, s); 6.89 (1H, s); 7.31-7.48 (12H, m); 8.29 (2H, m); 8.92 (1H, s); 11.05 (1H, s)
21		14	509		¹ H NMR d (d6-DMSO): 4.5 (1H, d), 5.25 (s, 4H), 6.9 (s, 1H), 7.40 (m, 6H), 7.5 (m, 2H), 7.6 (m, 2H), 7.75 (dd, 1H), 8.10 (d, 1H), 8.3 (s, 1H), 10.8 (br s, 1H);
22		1	494/496		¹ H NMR d (d6-DMSO): 5.25 (4H, s); 5.65 (2H, s); 6.23 (1H, d); 6.85 (1H, s); 7.05-7.15 (3H, m); 7.18-7.22 (5H, m); 7.45-7.55 (2H, m); 7.58-7.62 (2H, m); 10.16 (1H, br s).


Example	Structure	Route	(M+H)+	(M-H)-	NMR
23		1	476		¹ H NMR d (d ₆ -DMSO): 5.25 (4H, s); 5.75 (2H, s); 6.22 (1H, d); 6.90 (1H, s); 7.25-7.41 (4H, m); 7.50-7.60 (2H, m); 7.70-7.80 (4H, m); 7.90 (2H, d); 10.19 (1H, br s).
24		15	536/538		¹ H NMR d (d ₆ -DMSO): 3.25 (3H, s); 5.20 (4H, s); 6.9 (1H, t); 7.25 (2H, d); 7.35-7.40 (4H, m); 7.4 - 7.55 (2H, m); 7.58-7.63 (2H, m); 7.68-7.72 (1H, m); 7.75-7.80 (2H, d); 10.14 (1H, br s); 10.36 (1H, br s).
25		16	479	477	¹ H NMR d (d ₆ -DMSO): 5.19 (4H, s); 6.88 (1H, s); 7.26-7.48 (12H, m); 8.40 (1H, d); 8.46 (1H, dd); 9.04 (1H, s); 11.13 (1H, br s).
26		17	495	493	¹ H NMR d (d ₆ -DMSO): 5.19 (4H, s); 6.87 (1H, s); 7.28-7.46 (12H, m); 8.21 (1H, dd); 8.38 (1H, d); 8.79 (1H, s); 11.14 (1H, br s).
27		18	498		¹ H NMR d (d ₆ -DMSO): 5.18 (4H, s); 6.88 (1H, s); 7.30-7.50 (12H, m); 8.17 (2H, s); 8.79 (1H, s); 10.79 (1H, s); 10.93 (1H, br s).
28		1	460		¹ H NMR δ (d ₆ -DMSO): 2.32 (s, 3H), 2.36 (s, 3H), 3.23 (t, 2H), 4.22 (t, 2H), 5.13 (s, 2H), 6.78 (m, 1H), 7.11-7.24 (brm, 5H), 7.30 (m, 1H), 7.41 (d, 1H), 7.83 (m, 1H), 8.14 (d, 1H), 8.37 (m, 1H), 8.82 (s, 1H), 10.74 (brs, 1H).
29		7	475		¹ H NMR δ (d ₆ -DMSO): 2.32 (s, 3H), 2.36 (s, 3H), 3.22 (t, 2H), 4.20 (t, 2H), 5.11 (s, 4H), 6.72 (m, 1H), 7.00 (m, 1H), 7.15-7.28 (brm, 5H), 7.41 (d, 1H), 7.73 (m, 2H), 8.82 (s, 1H), 10.29 (brs, 1H).

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
30		1			¹ H NMR δ (d ₆ -DMSO): 2.66 (s, 3H), 5.15 (s, 4H), 6.88 (m, 1H), 7.14 (m, 1H), 7.39-7.47 (brm, 7H), 7.52 (s, 1H), 7.83 (m, 1H), 8.15 (d, 1H), 8.38 (m, 1H), 10.72 (brs, 1H).
31		1b	395		
32		1			¹ H NMR δ (d ₆ -DMSO): 1.28 (d, 6H), 2.39 (s, 3H), 4.72 (m, 1H), 5.20 (s, 2H), 6.33 (s, 1H), 6.72 (s, 1H), 7.14 (m, 1H), 7.20 (s, 1H), 7.27 (s, 1H), 7.82 (m, 1H), 8.13 (d, 1H), 8.36 (d, 1H), 10.72 (brs, 1H).
33		1			¹ H NMR δ (d ₆ -DMSO): 1.27 (d, 6H), 4.71 (m, 1H), 5.21 (s, 2H), 6.73 (t, 1H), 7.12 - 7.29 (brm, 5H), 7.22 (m, 1H), 7.56 (m, 1H), 7.83 (m, 1H), 8.14 (d, 1H), 8.35 (m, 1H), 10.72 (brs, 1H).
34		1b	311		
35		1b	451		
36		1b	398		
37		1	374	372	¹ H NMR δ (d ₆ -DMSO): 0.98 (d, 6H), 1.27 (d, 6H), 2.01 (m, 1H), 3.60 (d, 2H), 4.71 (m, 1H), 6.67 (ap t, 1H), 7.17 (ap d, 2H), 8.39 (d, 1H), 8.63 (dd, 1H), 9.20 (d, 1H), 11.43 (bs, 1H)
38		7b	344		¹ H NMR δ (d ₆ -DMSO): 0.97 (d, 6H), 1.26 (d, 6H), 2.00 (m, 1H), 3.78 (d, 2H), 4.69 (m, 1H), 5.12 (s, 2H), 6.58 (t, 1H), 6.99 (dd, 1H), 7.1 (ap d, 2H), 7.73-7.78 (m, 2H), 10.24 (bs, 1H)

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
39		15	386		¹ H NMR δ (d ₆ -DMSO): 0.98 (d, 6H), 1.26 (d, 6H), 2.01 (m, 1H), 2.05 (s, 3H), 3.79 (d, 2H), 4.70 (m, 1H), 6.61 (ap t, 1H), 7.14 (ap d, 2H), 7.95 (dd, 1H), 8.08 (d, 1H), 7.59 (ap d, 1H), 10.07 (bs, 1H)
40		15	422	420	¹ H NMR δ (d ₆ -DMSO): 0.97 (d, 6H), 1.26 (d, 6H), 2.03 (m, 1H), 3.01 (s, 3H), 3.79 (d, 2H), 4.70 (m, 1H), 6.63 (ap t, 1H), 7.14 (ap d, 2H), 7.70 (dd, 1H), 8.12 (d, 1H), 8.34 (ap d, 1H), (9.83, s, 1H), 10.81 (bs, 1H)
41		9	M+H 430 M-H 428		¹ H NMR δ (d ₆ -DMSO): 0.98 (d, 6H), 1.27 (d, 6H), 1.35 (s, 6H), 2.01 (m, 1H), 3.79 (d, 2H), 4.70 (m, 1H), 5.71 (s, 1H), 6.61 (s, 1H), 7.15 (s, 2H), 8.06-8.15 (m, 2H), 8.76 (ap d, 1H), 9.78 (s, 1H), 10.65 (bs, 1H)
42		15	412 (M+HCO OH) ⁺ 456		¹ H NMR δ (d ₆ -DMSO): 0.79-0.82 (m, 4H), 0.98 (d, 6H), 1.26 (d, 6H), 1.77 (m, 1H), 2.01 (m, 1H), 4.70 (h, 1H), 6.11 (ap t, 1H), 7.14 (ap d, 2H), 7.95 (dd, 1H), 8.08 (d, 1H), 8.62 (ap d, 1H), 10.33 (bs, 1H), 10.64 (bs, 1H)
43		27	M+H 450 M-H 448		¹ H NMR δ (d ₆ -DMSO): 0.98 (d, 6H), 1.27 (d, 6H), 2.01 (m, 1H), 3.37 (s, 3H), 3.80 (d, 2H), 4.71 (m, 1H), 6.65 (ap t, 1H), 7.17 (s, 2H), 8.27-8.35 (m, 2H), 8.91 (m, 1H), 11.13 (bs, 1H)
44		1c	352		δ _H (300MHz, DMSO-d ₆) 0.94-1.02 (6H,d); 1.24-1.34 (6H,d); 1.095-2.10 (1H,m); 3.76-3.84 (2H,d); 4.64-4.77 (1H,m); 6.64-6.70 (1H,m); 7.14-7.17 (2H,m); 8.25-8.36 (2H,m); 8.85 (1H,m); 11.21 (1H,s)
45		8 (a) 7c			δ _H (300MHz, DMSO-d ₆) 0.94-1.03 (6H,d); 1.26-1.30 (6H,d); 1.95-2.08 (1H,m); 2.90 (3H,s); 3.75-3.84 (2H,d); 4.04-4.26 (2H,d + H ₂ O); 4.65-4.77 (1H,m); 6.64 (1H,m); 7.15 (2H,m); 7.50-7.62 (1H,broad t); 7.80-7.90 (1H,d of m); 8.08-8.16 (1H,app d); 8.35 (1H, m); 10.84 (1H,m)

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
46		8 (a) 7c			δ_H (300MHz, DMSO- d_6) 0.94-1.02 (6H, d); 1.24-1.30 (6H, d); 1.84 (3H, s); 1.95-2.07 (1H, m); 3.75-3.83 (2H, d), 4.18-4.27 (2H, d); 4.64-4.76 (1H, m); 6.62 (1H, m); 7.15 (2H, m); 7.63-7.73 (1H, app d of m); 8.05-8.13 (1H, app d); 8.27 (1H, s); 8.30-8.38 (1H, app broad t); 10.69 (1H, s)
46a		1a	408	406	δ_H (300MHz, DMSO- d_6) 1.26 (d, 6H), 3.05 (t, 2H), 4.25 (t, 2H), 4.72 (sept, 1H), 6.68 (s, 1H), 7.12 (d, 1H), 7.16 (s, 1H), 7.19 (s, 1H), 7.33 (s, 1H), 7.47 (dd, 1H), 8.30 (m, 2H), 8.83 (s, 1H), 11.23 (bs, 1H)
47		27	504		δ_H (300MHz, DMSO- d_6) 1.27 (d, 6H), 3.06 (t, 2H), 3.38 (s, 3H), 4.25 (t, 2H), 4.71 (sept, 1H), 6.68 (t, 1H), 7.11 (dd, 1H), 7.12 (s, 1H), 7.17 (s, 1H), 7.31 (d, 1H), 7.46 (dd, 1H), 8.29 (d, 1H), 8.34 (dd, 1H), 8.92 (d, 1H), 11.14 (bs, 1H)
48		27	584	582	δ_H (300MHz, DMSO- d_6) 1.25 (d, 6H), 3.04 (t, 2H), 4.23 (t, 2H), 4.69 (sept, 1H), 6.67 (s, 1H), 7.11 (d, 1H), 7.15 (s, 1H), 7.20 (s, 1H), 7.31 (d, 1H), 7.46 (m, 3H), 8.07 (dd, 2H), 8.26 (s, 2H), 8.86 (s, 1H), 11.13 (bs, 1H)
49		27	556		δ_H (300MHz, DMSO- d_6) 1.27 (d, 6H), 3.04 (t, 2H), 4.23 (t, 2H), 4.71 (sept, 1H), 6.64 (s, 1H), 7.11 (d, 1H), 7.18 (s, 1H), 7.22 (s, 1H), 7.32 (s, 1H), 7.46 (dd, 1H), 8.19 (m, 2H), 8.82 (d, 1H), 10.93 (bs, 1H)
50		27	567		δ_H (300MHz, DMSO- d_6) 1.26 (d, 6H), 3.04 (t, 2H), 4.24 (t, 2H), 4.70 (sept, 1H), 6.64 (t, 1H), 7.11 (dd, 1H), 7.16 (s, 1H), 7.21 (s, 1H), 7.30 (m, 1H), 7.46 (m, 1H), 8.16 (m, 3H), 8.62 (d, 1H), 8.83 (s, 1H), 8.98 (s, 1H), 10.90 (bs, 1H)

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
51		27	585	583	δ_H (300MHz, DMSO- d_6) 1.27 (d, 6H), 2.39 (s, 3H), 2.68 (s, 3H), 3.06 (t, 2H), 4.26 (t, 2H), 4.73 (sept, 1H), 6.69 (t, 1H), 7.12 (d, 1H), 7.17 (s, 1H), 7.22 (s, 1H), 7.33 (m, 1H), 7.49 (m, 1H), 8.28 (m, 2H), 8.89 (s, 1H), 11.10 (bs, 1H)
52		27	618/620 (1xCl)	616/618 (1xCl)	δ_H (300MHz, DMSO- d_6) 1.28 (d, 6H), 2.40 (s, 3H), 3.08 (t, 2H), 3.79 (s, 3H), 4.25 (t, 2H), 4.71 (sept, 1H), 6.68 (s, 1H), 7.12 (d, 1H), 7.18 (s, 1H), 7.22 (s, 1H), 7.34 (m, 1H), 7.39 (s, 1H), 7.48 (dd, 1H), 8.30 (m, 2H), 8.92 (s, 1H), 11.15 (bs, 1H)
53		27	584		δ_H (300MHz, DMSO- d_6) 1.26 (d, 6H), 3.04 (t, 2H), 4.24 (t, 2H), 4.70 (sept, 1H), 6.66 (t, 1H), 7.12 (dd, 1H), 7.18 (s, 1H), 7.22 (s, 1H), 7.30 (m, 1H), 7.37 (m, 1H), 7.45 (dd, 1H), 7.67 (m, 1H), 7.78 (dt, 1H), 7.96 (dt, 1H), 8.822 (s, 2H), 8.86 (s, 1H), 11.08 (bs, 1H)
54		27	606	604	δ_H (300MHz, DMSO- d_6) 1.26 (d, 6H), 3.04 (t, 2H), 4.25 (t, 2H), 4.71 (sept, 1H), 6.64 (t, 1H), 7.00 (d, 1H), 7.12 (dd, 1H), 7.16 (s, 1H), 7.22 (s, 1H), 7.32 (m, 2H), 7.46 (dd, 1H), 8.14 (d, 1H), 8.22 (dd, 1H), 8.83 (t, 1H), 10.87 (bs, 1H)
55		16	451	449	δ_H (300MHz, DMSO- d_6) 1.28 (d, 6H), 3.06 (t, 2H), 4.26 (t, 2H), 4.72 (sept, 1H), 6.65 (s, 1H), 7.12 (d, 1H), 7.18 (s, 1H), 7.23 (s, 1H), 7.32 (s, 1H), 7.47 (m, 1H), 8.23 (d, 1H), 8.32 (dd, 1H), 8.95 (s, 1H), 10.81 (bs, 1H)
56		1a	329.48	327.46	

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
57		1a	354.46	352.43	

Notes:

* Final products prepared by hydrolysis method 2a; requisite starting materials prepared according to generic alkylation methodology followed by coupling (Route 1).



- 5 ** Final products prepared by reductive amination method 6 method ; requisite starting materials prepared according to generic alkylation methodology followed by coupling (Route 1) and hydrolysis (Route 2a).






*** Final products prepared by hydrolysis (Route 2a) or acid chloride coupling (Route 1); requisite starting materials prepared according to generic alkylation methodology followed by
10 coupling (Route 1).

EXAMPLE KK:


By analogous methods to those described above the following compounds, Example numbers KK₁ to KK₇, were also made.

15


Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
1		2b *	522	520	¹ H NMR d (d6-DMSO): 5.20 (4H, s); 6.95 (1H, s); 7.25 (2H, s); 7.30-7.5 (4H, m); 7.5 (2H, m); 7.6 (2H, m); 7.8 - 8.0 (4H, s).
2		1	494		No data

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
3		1		556/558	NMR not right
4		2b	522		¹ H NMR d (d6-DMSO): 5.25 (4H, s); 6.95 (1H, s); 7.25 (2H, s); 7.35-7.55 (7H, m); 7.6 - 7.7 (3H, m); 8.05 (1H, d); 8.4 (1H, s); 10.3 (1H, br s); 12.9 (1H, br s).
5		2b *	536	534	¹ H NMR d (d6-DMSO): 3.4 (2H, s); 5.2 (4H, s); 6.95 (1H, s); 7.2 (4H, m); 7.4 (4H, m); 7.5 (2H, m); 7.6 - 7.7 (4H, m); 10.1 (1H, br s).
6		1		519	¹ H NMR d (d6-DMSO): 5.2 (4H, s); 6.95 (1H, m); 7.25 (2H, m); 7.4 (5H, m); 7.5 (2H, m); 7.55 - 7.65 (4H, m); 7.9 (2H, m); 8.2 (1H, s); 10.3 (1H, br s).
7		1		577	V. poor spectrum

* For Examples KK₁ and KK₅, the ester intermediates were prepared by route 1:



^1H NMR δ (d_6 -DMSO): 3.8 (3H, s); 5.25 (4H, s); 6.95 (1H, t); 7.25 (2H, d); 7.4 (4H, m); 7.5 (2H, m); 7.6 (2H, m); 8.0 (4H, q); 10.6 (1H, br s).



5 ^1H NMR δ (d_6 -DMSO): 1.2 (3H, t); 3.6 (2H, s); 4.1 (2H, q); 5.25 (4H, s); 6.95 (1H, t); 7.2 (4H, m); 7.4 (4H, m); 7.5 (2H, m); 7.6 (2H, m); 7.7 (2H, m); 10.15 (1H, br s).



EXAMPLE LL:

10 By analogous methods to those described above the following compounds, Example numbers LL₁ to LL₃, were also made.

Example	Structure	Route	(M+H) ⁺	(M-H) ⁻	NMR
1	<p>Chemical structure of compound 1: 4-isopropoxy-N-(2,4-bis(isopropoxy)phenyl)benzamide with a 1,2,4-triazole group.</p>	1a	360		
2	<p>Chemical structure of compound 2: 4-isopropoxy-N-(2,4-bis(isopropoxy)phenyl)benzamide with a 1,2,4-triazole group and a fluorophenyl group.</p>	1a	382		
3	<p>Chemical structure of compound 3: 4-isopropoxy-N-(2,4-bis(isopropoxy)phenyl)benzamide with a 1,2,4-triazole group and a fluorophenyl group.</p>	1a	412		

EXAMPLE MM:

By analogous methods to those described above the following compounds, Example numbers MM₁ to MM₂, were also made.

Exempl e	Structure	Route	(M+H) ⁺	(M+H)	NMR
1		1a	385		
2		1a	371		

5

BIOLOGICALTests:

The biological effects of the compounds of formula (I) or (IA) or (IB) may be tested in the
10 following way:

(1) Enzymatic activity of GLK may be measured by incubating GLK, ATP and glucose. The rate of product formation may be determined by coupling the assay to a G-6-P dehydrogenase, NADP/NADPH system and measuring the increase in optical density at
15 340nm (Matschinsky et al 1993).

(2) A GLK/GLKRP binding assay for measuring the binding interactions between GLK and GLKRP. The method may be used to identify compounds which modulate GLK by modulating the interaction between GLK and GLKRP. GLKRP and GLK are incubated with
20 an inhibitory concentration of F-6-P, optionally in the presence of test compound, and the extent of interaction between GLK and GLKRP is measured. Compounds which either displace F-6-P or in some other way reduce the GLK/GLKRP interaction will be detected by a decrease in the amount of GLK/GLKRP complex formed. Compounds which promote F-6-P binding or in some other way enhance the GLK/GLKRP interaction will be detected by an

increase in the amount of GLK/GLKRP complex formed. A specific example of such a binding assay is described below

GLK/GLKRP scintillation proximity assay

- 5 The compounds A to S (described in Examples A to S) and 1 to 118 (described in Examples T to Y) were found to have an activity of at least 40% activity at 10 μ m when tested in the GLK/GLKRP scintillation proximity assay described below.

Recombinant human GLK and GLKRP were used to develop a "mix and measure" 96
10 well SPA (scintillation proximity assay) as described in WO01/20327 (the contents of which are incorporated herein by reference). GLK (Biotinylated) and GLKRP are incubated with streptavidin linked SPA beads (Amersham) in the presence of an inhibitory concentration of radiolabelled [3H]F-6-P (Amersham Custom Synthesis TRQ8689), giving a signal. Compounds which either displace the F-6-P or in some other way disrupt the GLK / GLKRP
15 binding interaction will cause this signal to be lost.

Binding assays were performed at room temperature for 2 hours. The reaction mixtures contained 50mM Tris-HCl (pH 7.5), 2mM ATP, 5mM MgCl₂, 0.5mM DTT, recombinant biotinylated GLK (0.1 mg), recombinant GLKRP (0.1 mg), 0.05mCi [3H] F-6-P (Amersham) to give a final volume of 100ml. Following incubation, the extent of
20 GLK/GLKRP complex formation was determined by addition of 0.1mg/well avidin linked SPA beads (Amersham) and scintillation counting on a Packard TopCount NXT.

(3) A F-6-P / GLKRP binding assay for measuring the binding interaction between GLKRP and F-6-P. This method may be used to provide further information on the
25 mechanism of action of the compounds. Compounds identified in the GLK/GLKRP binding assay may modulate the interaction of GLK and GLKRP either by displacing F-6-P or by modifying the GLK/GLKRP interaction in some other way. For example, protein-protein interactions are generally known to occur by interactions through multiple binding sites. It is thus possible that a compound which modifies the interaction between GLK and GLKRP
30 could act by binding to one or more of several different binding sites.

The F-6-P / GLKRP binding assay identifies only those compounds which modulate the interaction of GLK and GLKRP by displacing F-6-P from its binding site on GLKRP.

- 134 -

GLKRP is incubated with test compound and an inhibitory concentration of F-6-P, in the absence of GLK, and the extent of interaction between F-6-P and GLKRP is measured. Compounds which displace the binding of F-6-P to GLKRP may be detected by a change in the amount of GLKRP/F-6-P complex formed. A specific example of such a binding assay is
5 described below

F-6-P / GLKRP scintillation proximity assay

Recombinant human GLKRP was used to develop a "mix and measure" 96 well scintillation proximity assay) as described in WO01/20327 (the contents of which are
10 incorporated herein by reference). FLAG-tagged GLKRP is incubated with protein A coated SPA beads (Amersham) and an anti-FLAG antibody in the presence of an inhibitory concentration of radiolabelled [3H]F-6-P. A signal is generated. Compounds which displace the F-6-P will cause this signal to be lost. A combination of this assay and the GLK/GLKRP binding assay will allow the observer to identify compounds which disrupt the GLK/GLKRP
15 binding interaction by displacing F-6-P.

Binding assays were performed at room temperature for 2 hours. The reaction mixtures contained 50mM Tris-HCl (pH 7.5), 2mM ATP, 5mM MgCl₂, 0.5mM DTT, recombinant FLAG tagged GLKRP (0.1 mg), Anti-Flag M2 Antibody (0.2mg) (IBI Kodak), 0.05mCi [3H] F-6-P (Amersham) to give a final volume of 100µl. Following incubation, the
20 extent of F-6-P/GLKRP complex formation was determined by addition of 0.1mg/well protein A linked SPA beads (Amersham) and scintillation counting on a Packard TopCount NXT.

Production of recombinant GLK and GLKRP:

25 *Preparation of mRNA*

Human liver total mRNA was prepared by polytron homogenisation in 4M guanidine isothiocyanate, 2.5mM citrate, 0.5% Sarkosyl, 100mM β-mercaptoethanol, followed by centrifugation through 5.7M CsCl, 25mM sodium acetate at 135,000g (max) as described in Sambrook J, Fritsch EF & Maniatis T, 1989.

30 Poly A⁺ mRNA was prepared directly using a FastTrack™ mRNA isolation kit (Invitrogen).

PCR amplification of GLK and GLKRP cDNA sequences

Human GLK and GLKRP cDNA was obtained by PCR from human hepatic mRNA using established techniques described in Sambrook, Fritsch & Maniatis, 1989. PCR primers were designed according to the GLK and GLKRP cDNA sequences shown in Tanizawa et al 5 1991 and Bonthron, D.T. *et al* 1994 (later corrected in Warner, J.P. 1995).

Cloning in Bluescript II vectors

GLK and GLKRP cDNA was cloned in E. coli using pBluescript II, (Short et al 1998) a recombinant cloning vector system similar to that employed by Yanisch-Perron C *et al* 10 (1985), comprising a colEI-based replicon bearing a polylinker DNA fragment containing multiple unique restriction sites, flanked by bacteriophage T3 and T7 promoter sequences; a filamentous phage origin of replication and an ampicillin drug resistance marker gene.

Transformations

15 E. Coli transformations were generally carried out by electroporation. 400 ml cultures of strains DH5a or BL21(DE3) were grown in L-broth to an OD 600 of 0.5 and harvested by centrifugation at 2,000g. The cells were washed twice in ice-cold deionised water, resuspended in 1ml 10% glycerol and stored in aliquots at -70°C. Ligation mixes were desalted using Millipore V series™ membranes (0.0025mm) pore size). 40ml of cells were 20 incubated with 1ml of ligation mix or plasmid DNA on ice for 10 minutes in 0.2cm electroporation cuvettes, and then pulsed using a Gene Pulser™ apparatus (BioRad) at 0.5kVcm⁻¹, 250mF, 250 Ω . Transformants were selected on L-agar supplemented with tetracycline at 10mg/ml or ampicillin at 100mg/ml.

25 Expression

GLK was expressed from the vector pTB375NBSE in E.coli BL21 cells,, producing a recombinant protein containing a 6-His tag immediately adjacent to the N-terminal methionine. Alternatively, another suitable vector is pET21(+)-DNA, Novagen, Cat number 697703. The 6-His tag was used to allow purification of the recombinant protein on a column 30 packed with nickel-nitrilotriacetic acid agarose purchased from Qiagen (cat no 30250).

GLKRP was expressed from the vector pFLAG CTC (IBI Kodak) in E.coli BL21 cells, producing a recombinant protein containing a C-terminal FLAG tag. The protein was purified

- 136 -

initially by DEAE Sepharose ion exchange followed by utilisation of the FLAG tag for final purification on an M2 anti-FLAG immunoaffinity column purchased from Sigma-Aldrich (cat no. A1205).

5 Biotinylation of GLK:

GLK was biotinylated by reaction with biotinamidocaproate N-hydroxysuccinimide ester (biotin-NHS) purchased from Sigma-Aldrich (cat no. B2643). Briefly, free amino groups of the target protein (GLK) are reacted with biotin-NHS at a defined molar ratio forming stable amide bonds resulting in a product containing covalently bound biotin. Excess, non-
 10 conjugated biotin-NHS is removed from the product by dialysis. Specifically, 7.5mg of GLK was added to 0.31mg of biotin-NHS in 4mL of 25mM HEPES pH7.3, 0.15M KCl, 1mM dithiothreitol, 1mM EDTA, 1mM MgCl₂ (buffer A). This reaction mixture was dialysed against 100mL of buffer A containing a further 22mg of biotin-NHS. After 4hours excess biotin-NHS was removed by extensive dialysis against buffer A.

15

PHARMACEUTICAL COMPOSITIONS

The following illustrate representative pharmaceutical dosage forms of the invention as
 20 defined herein (the active ingredient being termed "Compound X"), for therapeutic or prophylactic use in humans:

(a)	<u>Tablet I</u>	<u>mg/tablet</u>
	Compound X.....	100
25	Lactose Ph.Eur.....	182.75
	Croscarmellose sodium.....	12.0
	Maize starch paste (5% w/v paste).....	2.25
	Magnesium stearate.....	3.0
30	(b)	
	<u>Tablet II</u>	<u>mg/tablet</u>
	Compound X.....	50
	Lactose Ph.Eur.....	223.75
	Croscarmellose sodium.....	6.0

- 137 -

	Maize starch.....	15.0
	Polyvinylpyrrolidone (5% w/v paste).....	2.25
	Magnesium stearate.....	3.0
5 (c)	<u>Tablet III</u>	<u>mg/tablet</u>
	Compound X.....	1.0
	Lactose Ph.Eur.....	93.25
	Croscarmellose sodium.....	4.0
	Maize starch paste (5% w/v paste).....	0.75
10	Magnesium stearate.....	1.0
(d)	<u>Capsule</u>	<u>mg/capsule</u>
	Compound X.....	10
	Lactose Ph.Eur.....	488.5
15	Magnesium.....	1.5
(e)	<u>Injection I</u>	<u>(50 mg/ml)</u>
	Compound X.....	5.0% w/v
	1M Sodium hydroxide solution.....	15.0% v/v
20	0.1M Hydrochloric acid (to adjust pH to 7.6)	
	Polyethylene glycol 400.....	4.5% w/v
	Water for injection to 100%	
(f)	<u>Injection II</u>	<u>(10 mg/ml)</u>
25	Compound X.....	1.0% w/v
	Sodium phosphate BP.....	3.6% w/v
	0.1M Sodium hydroxide solution.....	15.0% v/v
	Water for injection to 100%	
30 (g)	<u>Injection III</u>	<u>(1mg/ml, buffered to pH6)</u>
	Compound X.....	0.1% w/v
	Sodium phosphate BP.....	2.26% w/v
	Citric acid.....	0.38% w/v

- 138 -

Polyethylene glycol 400..... 3.5% w/v
 Water for injection to 100%

5	(h)	<u>Aerosol I</u>	<u>mg/ml</u>
		Compound X.....	10.0
		Sorbitan trioleate.....	13.5
		Trichlorofluoromethane.....	910.0
		Dichlorodifluoromethane.....	490.0
10	(i)	<u>Aerosol II</u>	<u>mg/ml</u>
		Compound X.....	0.2
		Sorbitan trioleate.....	0.27
		Trichlorofluoromethane.....	70.0
		Dichlorodifluoromethane.....	280.0
15		Dichlorotetrafluoroethane.....	1094.0
20	(j)	<u>Aerosol III</u>	<u>mg/ml</u>
		Compound X.....	2.5
		Sorbitan trioleate.....	3.38
		Trichlorofluoromethane.....	67.5
		Dichlorodifluoromethane.....	1086.0
		Dichlorotetrafluoroethane.....	191.6
25	(k)	<u>Aerosol IV</u>	<u>mg/ml</u>
		Compound X.....	2.5
		Soya lecithin.....	2.7
		Trichlorofluoromethane.....	67.5
		Dichlorodifluoromethane.....	1086.0
		Dichlorotetrafluoroethane.....	191.6
30	(l)	<u>Ointment</u>	<u>ml</u>
		Compound X.....	40 mg
		Ethanol.....	300 µl

- 139 -

Water.....	300 µl
1-Dodecylazacycloheptan-2-one.....	50 µl
Propylene glycol.....	to 1 ml

5 Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate. The aerosol formulations (h)-(k) may be used in conjunction with standard, metered dose aerosol dispensers, and the suspending agents sorbitan trioleate and soya lecithin may be replaced by an alternative suspending agent such as sorbitan monooleate, sorbitan sesquioleate, polysorbate 80, polyglycerol oleate or oleic acid.

15 REFERENCES


- 1 Printz, R. L., Magnuson, M. A. and Granner, D. K. (1993) Annual Review of Nutrition 13, 463-96
- 2 DeFronzo, R. A. (1988) Diabetes 37, 667-87
- 3 Froguel, P., Zouali, H., Vionnet, N., Velho, G., Vaxillaire, M., Sun, F., Lesage, S., Stoffel, M., Takeda, J. and Passa, P. (1993) New England Journal of Medicine 328, 697-702
- 4 Bell, G. I., Pilkis, S. J., Weber, I. T. and Polonsky, K. S. (1996) Annual Review of Physiology 58, 171-86
- 5 Velho, G., Petersen, K. F., Perseghin, G., Hwang, J. H., Rothman, D. L., Pueyo, M. E., Cline, G. W., Froguel, P. and Shulman, G. I. (1996) Journal of Clinical Investigation 98, 1755-61
- 6 Christesen, H. B., Jacobsen, B. B., Odili, S., Buettger, C., Cuesta-Munoz, A., Hansen, T., Brusgaard, K., Massa, O., Magnuson, M. A., Shiota, C., Matschinsky, F. M. and Barbetti, F. (2002) Diabetes 51, 1240-6
- 7 Glaser, B., Kesavan, P., Heyman, M., Davis, E., Cuesta, A., Buchs, A., Stanley, C. A., Thornton, P. S., Permutt, M. A., Matschinsky, F. M. and Herold, K. C. (1998) New England Journal of Medicine 338, 226-30

- 8 Caro, J. F., Triester, S., Patel, V. K., Tapscott, E. B., Frazier, N. L. and Dohm, G. L. (1995) *Hormone & Metabolic Research* **27**, 19-22
- 9 Desai, U. J., Slosberg, E. D., Boettcher, B. R., Caplan, S. L., Fanelli, B., Stephan, Z., Gunther, V. J., Kaleko, M. and Connelly, S. (2001) *Diabetes* **50**, 2287-95
- 5 10 Shiota, M., Postic, C., Fujimoto, Y., Jetton, T. L., Dixon, K., Pan, D., Grimsby, J., Grippo, J. F., Magnuson, M. A. and Cherrington, A. D. (2001) *Diabetes* **50**, 622-9
- 11 Ferre, T., Pujol, A., Riu, E., Bosch, F. and Valera, A. (1996) *Proceedings of the National Academy of Sciences of the United States of America* **93**, 7225-30
- 12 Seoane, J., Barbera, A., Telemaque-Potts, S., Newgard, C. B. and Guinovart, J. J. (1999) *Journal of Biological Chemistry* **274**, 31833-8
- 10 13 Moore, M. C., Davis, S. N., Mann, S. L. and Cherrington, A. D. (2001) *Diabetes Care* **24**, 1882-7
- 14 Alvarez, E., Roncero, I., Chowen, J. A., Vazquez, P. and Blazquez, E. (2002) *Journal of Neurochemistry* **80**, 45-53
- 15 15 Lynch, R. M., Tompkins, L. S., Brooks, H. L., Dunn-Meynell, A. A. and Levin, B. E. (2000) *Diabetes* **49**, 693-700
- 16 Roncero, I., Alvarez, E., Vazquez, P. and Blazquez, E. (2000) *Journal of Neurochemistry* **74**, 1848-57
- 17 Yang, X. J., Kow, L. M., Funabashi, T. and Mobbs, C. V. (1999) *Diabetes* **48**, 1763-1772
- 20 18 Schuit, F. C., Huypens, P., Heimberg, H. and Pipeleers, D. G. (2001) *Diabetes* **50**, 1-11
- 19 Levin, B. E. (2001) *International Journal of Obesity* **25**
- 20 Alvarez, E., Roncero, I., Chowen, J. A., Thorens, B. and Blazquez, E. (1996) *Journal of Neurochemistry* **66**, 920-7
- 25 21 Mobbs, C. V., Kow, L. M. and Yang, X. J. (2001) *American Journal of Physiology - Endocrinology & Metabolism* **281**, E649-54
- 22 Levin, B. E., Dunn-Meynell, A. A. and Routh, V. H. (1999) *American Journal of Physiology* **276**, R1223-31
- 23 Spanswick, D., Smith, M. A., Groppi, V. E., Logan, S. D. and Ashford, M. L. (1997) *Nature* **390**, 521-5
- 30 24 Spanswick, D., Smith, M. A., Mirshamsi, S., Routh, V. H. and Ashford, M. L. (2000) *Nature Neuroscience* **3**, 757-8

- 25 Levin, B. E. and Dunn-Meynell, A. A. (1997) *Brain Research* **776**, 146-53
- 26 Levin, B. E., Govek, E. K. and Dunn-Meynell, A. A. (1998) *Brain Research* **808**, 317-9
- 27 Levin, B. E., Brown, K. L. and Dunn-Meynell, A. A. (1996) *Brain Research* **739**, 293-300
- 5 28 Rowe, I. C., Boden, P. R. and Ashford, M. L. (1996) *Journal of Physiology* **497**, 365-77
- 29 Fujimoto, K., Sakata, T., Arase, K., Kurata, K., Okabe, Y. and Shiraishi, T. (1985) *Life Sciences* **37**, 2475-82
- 30 Kurata, K., Fujimoto, K. and Sakata, T. (1989) *Metabolism: Clinical & Experimental* **38**, 46-51
- 10 31 Kurata, K., Fujimoto, K., Sakata, T., Etou, H. and Fukagawa, K. (1986) *Physiology & Behavior* **37**, 615-20

CLAIMS:

1. The use of a compound of Formula (I) or a salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the treatment or prevention of a disease or
 5 medical condition mediated through GLK:



Formula (I)

wherein

- m is 0, 1 or 2;
 10 n is 0, 1, 2, 3 or 4;
 and $n + m > 0$;
 each R^1 is independently selected from OH, $-(CH_2)_{1-4}OH$, $-CH_3-aF_a$, $-(CH_2)_{1-4}CH_3-aF_a$,
 $-OCH_3-aF_a$, halo, $C_{1-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, NH_2 , $-NH-C_{1-4}alkyl$,
 $-N-di-(C_{1-4}alkyl)$, CN, formyl, phenyl or heterocyclyl optionally substituted by
 15 $C_{1-6}alkyl$;
 each R^2 is the group $Y-X-$
 wherein each X is a linker independently selected from:
 $-O-Z-$, $-O-Z-O-Z-$, $-C(O)O-Z-$, $-OC(O)-Z-$, $-S-Z-$, $-SO-Z-$, $-SO_2-Z-$, $-N(R^6)-Z-$,
 $-N(R^6)SO_2-Z-$, $-SO_2N(R^6)-Z-$, $-(CH_2)_{1-4}-$, $-CH=CH-Z-$, $-C\equiv C-Z-$, $-N(R^6)CO-Z-$,
 20 $-CON(R^6)-Z-$, $-C(O)N(R^6)S(O)_2-Z-$, $-S(O)_2N(R^6)C(O)-Z-$, $-C(O)-Z-$, $-Z-$,
 $-C(O)-Z-O-Z-$, $-N(R^6)-C(O)-Z-O-Z-$, $-O-Z-N(R^6)-Z-$, $-O-C(O)-Z-O-Z-$ or a
 direct bond;
 each Z is independently a direct bond, $C_{2-6}alkenylene$ or a group of the formula
 $-(CH_2)_p-C(R^{6a})_2-(CH_2)_q-$;
 25 each Y is independently selected from aryl- Z^1- , heterocyclyl- Z^1- ,
 $C_{3-7}cycloalkyl-Z^1-$, $C_{1-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, $-(CH_2)_{1-4}CH_3-aF_a$ or
 $-CH(OH)CH_3-aF_a$; wherein each Y is independently optionally substituted by up
 to 3 R^4 groups;

- 143 -

- each R^4 is independently selected from halo, $-CH_3$, $-F$, $-CN$, $-NH_2$, C_{1-6} alkyl, $-OC_{1-6}$ alkyl, $-COOH$, $-C(O)OC_{1-6}$ alkyl, $-OH$ or phenyl optionally substituted by C_{1-6} alkyl or $-C(O)OC_{1-6}$ alkyl, or R^5-X^1 , where X^1 is independently as defined in X above and R^5 is selected from hydrogen, C_{1-6} alkyl, $-CH_3$, $-F$, phenyl, naphthyl, heterocyclyl or C_{3-7} cycloalkyl; and R^5 is optionally substituted by one or more substituents independently selected from: halo, C_{1-6} alkyl, $-OC_{1-6}$ alkyl, $-CH_3$, $-F$, $-CN$, $-OH$, $-NH_2$, $-COOH$, or $-C(O)OC_{1-6}$ alkyl, each Z^1 is independently a direct bond, C_{2-6} alkenylene or a group of the formula $-(CH_2)_p-C(R^{6a})_2-(CH_2)_q$;
- R^3 is selected from phenyl or a heterocyclyl, and R^3 is optionally substituted by one or more R^7 groups;
- R^6 is independently selected from hydrogen, C_{1-6} alkyl or $-C_{2-4}$ alkyl-O- C_{1-4} alkyl;
- R^{6a} is independently selected from hydrogen, halo, C_{1-6} alkyl or $-C_{2-4}$ alkyl-O- C_{1-4} alkyl;
- each R^7 is independently selected from:
- C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, $(CH_2)_{0-3}$ aryl, $(CH_2)_{0-3}$ heterocyclyl, $(CH_2)_{0-3}C_{3-7}$ cycloalkyl, $-OH$, C_{1-6} alkyl-OH, halo, C_{1-6} alkyl-halo, OC_{1-6} alkyl, $(CH_2)_{0-3}S(O)_{0-2}R^8$, $-SH$, $-SO_3$, thiooxo, $-NH_2$, $-CN$, $(CH_2)_{0-3}NHSO_2R^8$, $(CH_2)_{0-3}COOH$, $(CH_2)_{0-3}O-(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}C(O)OR^8$, $(CH_2)_{0-3}C(O)NH_2$, $(CH_2)_{0-3}C(O)NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NH(CH_2)_{0-3}R^8$, $(CH_2)_{0-3}NHC(O)(CH_2)_{0-3}R^8$; $(CH_2)_{0-3}C(O)NHSO_2-R^8$ and $(CH_2)_{0-3}SO_2NHC(O)-R^8$ wherein an alkyl chain, cycloalkyl ring or heterocyclyl ring within R^7 is optionally substituted by one of more substituents independently selected from: C_{1-4} alkyl, $-OH$, halo, $-CN$, $-NH_2$, $N-C_{1-4}$ alkylamino, N,N -di- C_{1-4} alkylamino and OC_{1-4} alkyl;
- R^8 is selected from hydrogen, C_{1-6} alkyl, aryl, heterocyclyl, C_{3-7} cycloalkyl, $-OH$, C_{1-6} alkyl-OH, $-COOH$, $-C(O)OC_{1-6}$ alkyl, $N(R^6)C_{1-6}$ alkyl, OC_{1-6} alkyl, C_{0-6} alkylOC(O) C_{1-6} alkyl, $C(OH)(C_{1-6}alkyl)C_{1-6}alkyl$; wherein an alkyl chain or aryl, heterocyclyl or cycloalkyl ring within R^8 is optionally substituted by one of more substituents independently selected from: C_{1-4} alkyl, $-OH$, halo, $-CN$, $-NH_2$, $-NH-C_{1-4}$ alkyl, $-N$ -di- $(C_{1-4}alkyl)$ and $OC_{1-4}alkyl$;

- 144 -

each a is independently 1, 2 or 3;

p is an integer between 0 and 3;


q is an integer between 0 and 3;

and $p + q < 4$.

- 5 provided that when R^3 is 2-pyridyl and X is other than $-Z-$, $-C(O)-Z-O-Z-$, $-N((R^6)-C(O)-Z-O-Z-$ or $-O-Z-N(R^6)-Z-$, then R^3 cannot be mono-substituted at the 5-position with an R^7 group selected from $COOH$ or $C(O)OC_{1-6}alkyl$.

2. A pharmaceutical composition comprising a compound of Formula (I) as claimed in
10 claim 1, or a salt, solvate or pro-drug thereof, together with a pharmaceutically-acceptable diluent or carrier for use in the preparation of a medicament for use in the treatment or prevention of a disease or medical condition mediated through GLK.

3. A compound of Formula (Ib) or a salt, solvate or pro-drug thereof



Formula (Ib)

wherein

m is 0, 1 or 2;

n is 1, 2 or 3;

20 and $n + m$ is 2 or 3;

each R^1 is independently selected from OH , $-(CH_2)_{1-4}OH$, $-CH_2F$, $-(CH_2)_{1-4}CH_2F$, $-OCH_2F$, halo, OCH_3 , C_2H_5O , $CH_3C(O)O-$, $C_{1-6}alkyl$, $C_{2-6}alkenyl$, $C_{2-6}alkynyl$, $-NH-C_{1-4}alkyl$, $-N-di-(C_{1-4}alkyl)$, CN , formyl, phenyl or heterocyclyl optionally substituted by $C_{1-6}alkyl$;

25 each R^2 is the group $Y-X-$

with the proviso that $Y-X-$ cannot be CH_3O , C_2H_5O or $CH_3C(O)O-$;

wherein each X is a linker independently selected from:

$-O-Z-$, $-O-Z-O-Z-$, $-C(O)O-Z-$, $-OC(O)-Z-$, $-S-Z-$, $-SO-Z-$, $-SO_2-Z-$, $-N(R^6)-Z-$, $-N(R^6)SO_2-Z-$, $-SO_2N(R^6)-Z-$, $-CH=CH-Z-$, $-C\equiv C-Z-$, $-N(R^6)CO-Z-$,

- 145 -

-CON(R⁶)-Z-, -C(O)N(R⁶)S(O)₂-Z-, -S(O)₂N(R⁶)C(O)-Z-, -C(O)-Z-, -Z-,
 -C(O)-Z-O-Z-, -N(R⁶)-C(O)-Z-O-Z-, -O-Z-N(R⁶)-Z-, -O-C(O)-Z-O-Z- or a
 direct bond except where Z is C₁₋₆alkyl;

each Z is independently a direct bond, C₂₋₆alkenylene or a group of the formula

5 $-(CH_2)_p-C(R^{6a})_2-(CH_2)_q-$;

each Y is independently selected from aryl-Z¹-, heterocyclyl-Z¹-,

C₃₋₇cycloalkyl-Z¹-, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, -(CH₂)₁₋₄CH₃₋₄F₁ or

-CH(OH)CH₃₋₄F₁; wherein each Y is independently optionally substituted by
 up to 3 R⁴ groups;

10 each R⁴ is independently selected from halo, -CH₃₋₄F₁, CN, NH₂, C₁₋₄alkyl,

-OC₁₋₆alkyl, -COOH, -C(O)OC₁₋₆alkyl, OH or phenyl optionally
 substituted by C₁₋₆alkyl or -C(O)OC₁₋₆alkyl,

or R⁵-X¹-, where X¹ is independently as defined in X above and R⁵ is
 selected from hydrogen, C₁₋₆alkyl, -CH₃₋₄F₁, phenyl, naphthyl,

15 heterocyclyl or C₃₋₇cycloalkyl; and R⁵ is optionally substituted by one

or more substituents independently selected from: halo, C₁₋₆alkyl,
 -OC₁₋₆alkyl, -CH₃₋₄F₁, CN, OH, NH₂, COOH, or -C(O)OC₁₋₆alkyl,

each Z¹ is independently a direct bond, C₂₋₆alkenylene or a group of
 the formula $-(CH_2)_p-C(R^{6a})_2-(CH_2)_q-$;

20 R³ is heterocyclyl, wherein the atom at the two position of the heterocyclyl ring
 relative to the amide group, to which R³ is attached, is a heteroatom and when the
 atom at the two position of the heterocyclyl ring relative to the amide group is
 nitrogen, this is an sp² hybridised nitrogen, and R³ is optionally substituted by up
 to 2 R⁷ groups;

25 R⁶ is independently selected from hydrogen, C₁₋₆alkyl or -C₂₋₄alkyl-O-C₁₋₄alkyl;

R^{6a} is independently selected from hydrogen, halo, C₁₋₆alkyl or
 -C₂₋₄alkyl-O-C₁₋₄alkyl;

each R⁷ is independently selected from:

C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, (CH₂)₀₋₃aryl, (CH₂)₀₋₃heterocyclyl,

30 (CH₂)₀₋₃C₃₋₇cycloalkyl, OH, C₁₋₆alkyl-OH, halo, C₁₋₆alkyl-halo, OC₁₋₆alkyl,

(CH₂)₀₋₃S(O)₀₋₂R⁸, SH, SO₃, thioxo, NH₂, CN, (CH₂)₀₋₃NHSO₂R⁸,

(CH₂)₀₋₃COOH, (CH₂)₀₋₃-O-(CH₂)₀₋₃R⁸, (CH₂)₀₋₃C(O)(CH₂)₀₋₃R⁸,

- 146 -

$(\text{CH}_2)_{0-3}\text{C}(\text{O})\text{OR}^8$, $(\text{CH}_2)_{0-3}\text{C}(\text{O})\text{NH}_2$, $(\text{CH}_2)_{0-3}\text{C}(\text{O})\text{NH}(\text{CH}_2)_{0-3}\text{R}^8$,
 $(\text{CH}_2)_{0-3}\text{NH}(\text{CH}_2)_{0-3}\text{R}^8$, $(\text{CH}_2)_{0-3}\text{NHC}(\text{O})(\text{CH}_2)_{0-3}\text{R}^8$; $(\text{CH}_2)_{0-3}\text{C}(\text{O})\text{NHSO}_2\text{-R}^8$ and
 $(\text{CH}_2)_{0-3}\text{SO}_2\text{NHC}(\text{O})\text{-R}^8$ wherein an alkyl chain, cycloalkyl ring or heterocyclyl
 ring within R^7 is optionally substituted by one of more substituents independently
 5 selected from: $\text{C}_{1-4}\text{alkyl}$, OH, halo, CN, NH_2 , $\text{N-C}_{1-4}\text{alkylamino}$,
 $\text{N,N-di-C}_{1-4}\text{alkylamino}$ and $\text{OC}_{1-4}\text{alkyl}$;

R^8 is selected from hydrogen, $\text{C}_{1-6}\text{alkyl}$, aryl, heterocyclyl, $\text{C}_{3-7}\text{cycloalkyl}$, OH,
 $\text{C}_{1-6}\text{alkyl-OH}$, COOH, $\text{C}(\text{O})\text{OC}_{1-6}\text{alkyl}$, $\text{N}(\text{R}^6)\text{C}_{1-6}\text{alkyl}$, $\text{OC}_{1-6}\text{alkyl}$,
 $\text{C}_{0-6}\text{alkylOC}(\text{O})\text{C}_{1-6}\text{alkyl}$, $\text{C}(\text{OH})(\text{C}_{1-6}\text{alkyl})\text{C}_{1-6}\text{alkyl}$; wherein an alkyl chain or
 10 aryl, heterocyclyl or cycloalkyl ring within R^8 is optionally substituted by one of
 more substituents independently selected from: $\text{C}_{1-4}\text{alkyl}$, OH, halo, CN, NH_2 ,
 $-\text{NH-C}_{1-4}\text{alkyl}$, $-\text{N-di-(C}_{1-4}\text{alkyl)}$ and $\text{OC}_{1-4}\text{alkyl}$;

each a is independently 1, 2 or 3;

p is an integer between 0 and 3;

15 q is an integer between 0 and 3;

and $p + q < 4$.

provided that

- (i) when R^3 is 2-pyridyl and X is other than $-\text{Z}-$, $-\text{C}(\text{O})\text{-Z-O-Z-}$, $-\text{N}(\text{R}^6)\text{-C}(\text{O})\text{-Z-O-Z-}$
 or $-\text{O-Z-N}(\text{R}^6)\text{-Z-}$, then R^3 cannot be mono-substituted at the 5-position with an
 20 R^7 group selected from COOH or $\text{C}(\text{O})\text{OC}_{1-6}\text{alkyl}$;
- (ii) positions 3,5 on the phenyl ring (to which R^1 and R^2 are attached) relative to the
 amide bond are substituted and at least one of the groups at position 3 and 5 is an
 R^2 group;
- (iii) an unbranched, unsubstituted $\text{C}_{1-6}\text{alkyl}$ chain cannot exceed C_6alkyl in length;
- 25 (iv) when n is 2 or 3 then only one X group can be $-\text{NHC}(\text{O})-$;
- (v) when R^3 is pyridyl and R^7 is halo or methyl then the phenyl ring to which R^2 is
 attached cannot be substituted by an R^2 group at the 2-position relative to the
 amide bond wherein X is $-\text{C}(\text{O})\text{NH-}$ and Y is optionally substituted phenyl,
 optionally substituted thienyl or optionally substituted pyridyl;
- 30 (vi) when $n+m$ is 2, m is 0 or m is 1 and R^1 is OH, n is 1 and X is $-\text{NHC}(\text{O})-$ or n is 2
 and X is independently selected from $-\text{C}(\text{O})\text{NH-}$, $-\text{NHC}(\text{O})-$, $-\text{O-}$, $-\text{S}(\text{O}_2)\text{NH-}$ or
 a direct bond wherein one X group is $-\text{NHC}(\text{O})-$, Y is selected from phenyl,

- 147 -

- cyclohexyl, 4,5-dihydro-5-oxo-pyrazolyl, thienyl,
 1,3-dihydro-1,3-dioxo-isoindolyl, 2-oxo-1-benzopyran or pyridyl and Y is
 optionally substituted by R^4 then R^3 cannot be unsubstituted thiazole,
 4,5-dihydro-5-oxo-pyrazolyl substituted by trichlorophenyl,
 5 4,5,6,7-tetrahydro-benzo[b]thiophene substituted by ethoxycarbonyl or pyridyl
 optionally independently mono or di-substituted by methyl, ethoxy or
 propylcarbonylamino; and
 (vii) when $n+m$ is 3, m is 0 or 2, R^1 is independently selected from methyl, methoxy or
 hydroxy, n is 1, 2 or 3, X is independently selected from $-O-$, $-S(O_2)NH-$,
 10 $-C(O)-$, $-S(O_2)-$, $-CH_2-$ or a direct bond, Y is selected from pyrrolidinyl,
 morpholino, phenyl, tetrazolyl or propyl wherein Y is optionally substituted by
 R^4 and R^4 is selected from di-hydroxy, methoxy, C_{1-4} alkyl then R^3 cannot be
 unsubstituted tetrazolyl, unsubstituted thiazolyl or thiazolyl substituted by
 ethoxycarbonylmethyl.
 15
4. A compound according to claim 3, or salt, solvate or pro-drug thereof, wherein each
 R^1 is independently selected from OH, formyl, CH_3-aF_a , OCH_3-aF_a , halo, C_{1-6} alkyl,
 NH_2 , CN, $(CH_2)_{1-4}OH$ or a heterocyclyl optionally substituted by C_{1-6} alkyl.
- 20 5. A compound according to claim 3 or claim 4, or salt, solvate or pro-drug thereof,
 wherein each R^2 is the group $Y-X-$, each X is independently selected from $-Z-$,
 $-CH=CH-Z-$, $-O-Z-$, $-C(O)-Z-$, $-C(O)O-Z-$, $-OC(O)-Z-$, $-C(O)-Z-O-Z-$,
 $-O-C(O)-Z-O-Z-$, $-S-Z-$, $-SO-Z-$, $-SO_2-Z-$, $-N(R^6)-Z-$, $-N(R^6)CO-Z-$, $-CON(R^6)-Z-$,
 $-N(R^6)-C(O)-Z-O-Z-$, $-SO_2N(R^6)-Z-$, $-N(R^6)SO_2-Z-$ or $-O-Z-N(R^6)-Z-$, each Y is
 25 independently selected from C_{1-6} alkyl, C_{2-6} alkenyl, aryl- Z^1- , heterocyclyl- Z^1- ,
 C_{3-7} cycloalkyl $(CH_2)_{0-2}$, $-(CH_2)_{1-4}CH_3-aF_a$ and each Y is independently optionally
 substituted by R^4 .
6. A compound according to any one of claims 3 to 5, or salt, solvate or pro-drug thereof,
 30 wherein each R^4 is independently selected from halo, CH_3-aF_a , OCH_3-aF_a , CN, C_{1-6} alkyl,
 OC_{1-6} alkyl, COOH, $C(O)OC_{1-6}$ alkyl, $(CH_2)_{0-3}COOH$, $O(CH_2)_{0-3}COOH$, CO-phenyl,
 $CONH_2$, CONH-phenyl, SO_2NH_2 , SO_2C_{1-6} alkyl, OH, or phenyl optionally substituted by


- 148 -

one or more R^5 groups where R^5 is selected from hydrogen, C_{1-6} alkyl or $C(O)OC_{1-6}$ alkyl.

7. A compound according to any one of claims 3 to 6, or salt, solvate or pro-drug thereof,
5 wherein R^3 is a nitrogen-containing heterocyclyl, optionally substituted by one or more R^7 groups.
8. A compound according to claim 7, or a salt, solvate or prodrug thereof, wherein R^3 is
selected from thiazole, benzothiazole, thiadiazole, pyridine, pyrazine, pyridazine,
10 pyrazole, imidazole, pyrimidine, oxazole and indole.
9. A compound according to any one of claims 3 to 8, or salt, solvate or pro-drug thereof,
wherein R^3 is unsubstituted or is substituted by one R^7 group.
- 15 10. A compound according to any one of claims 3 to 9, or salt, solvate or pro-drug thereof,
wherein each R^7 is independently selected from OH, CN, NH_2 , SO_3 , thioxo, halo, C_{1-4} alkyl, C_{1-4} alkyl-OH, O- C_{1-4} alkyl, C_{1-4} alkyl-halo, $(CH_2)_{0-1}COOH$, $(CH_2)_{0-1}C(O)OR^8$,
 $(CH_2)_{0-1}NH(CH_2)_{0-2}R^8$, $(CH_2)_{0-1}NHC(O)(CH_2)_{0-2}R^8$, $(CH_2)_{0-1}C(O)NH(CH_2)_{0-2}R^8$,
- $(CH_2)_{0-2}S(O)_{0-2}R^8$, - $(CH_2)_{0-1}N(R^6)SO_2R^8$, $(CH_2)_{0-1}C(O)N(R^6)S(O)_2R^8$ or
20 $(CH_2)_{0-1}$ heterocyclyl.
11. A compound according to any one of claims 3 to 10, or salt, solvate or pro-drug thereof,
wherein Y is phenyl- Z^1 - optionally substituted by halo or C_{1-6} alkyl.
- 25 12. A compound according to any one of claims 3 to 11, or salt, solvate or pro-drug thereof,
wherein each R^2 is the group Y-X-, Z within the definition of X is a direct bond and Z^1
within the definition of Y is a group of the formula $-(CH_2)_p-C(R^{6a})_2-(CH_2)_q-$.
13. A pharmaceutical composition comprising a compound according to any one of claims 3
30 to 12, or a salt, solvate or prodrug thereof, together with a pharmaceutically-acceptable
diluent or carrier.

- 149 -

14. The use of a compounds of Formula (I), as defined in claim 1, or a salt, solvate or pro-drug thereof, in the preparation of a medicament for use in the combined treatment or prevention of diabetes and obesity.
15. The use of a compound of Formula (Ib) or a salt, pro-drug or solvate thereof, as defined in claim 3, as a medicament.
16. A process for the preparation of a compound of Formula (I), or salt, solate or pro-drug thereof, as defined in claim 1, which comprises:
- (a) reaction of a compound of Formula (IIIa) with a compound of Formula (IIIb),




Formula (IIIa)



Formula (IIIb); or

wherein X^1 is a leaving group

- (b) for compounds of Formula (I) wherein R^3 is hydrogen, de-protection of a compound of Formula (IIIc),




Formula (IIIc)

wherein P^1 is a protecting group;

- (c) for compounds of Formula (I) wherein n is 1, 2, 3 or 4, reaction of a compound of Formula (IIId) with a compound of Formula (IIIe),



Formula (IIId)




Formula (IIIe)

wherein X' and X'' comprises groups which when reacted together form the group X ;

- 150 -

- (d) for a compound of Formula (I) wherein n is 1, 2, 3 or 4 and X or X¹ is -SO-Z- or -SO₂-Z-, oxidation of the corresponding compound of Formula (I) wherein X or X¹ respectively is -S-Z-;
- (e) reaction of a compound of Formula (III_f) with a compound of Formula (III_g),

Formula (III_f)Formula (III_g)

wherein X² is a leaving group;

and thereafter, if necessary:

i) converting a compound of Formula (I) into another compound of Formula (I);

ii) removing any protecting groups;

iii) forming a salt, pro-drug or solvate thereof.

INTERNATIONAL SEARCH REPORT

 Int'l. Application No
 PCT/GB 02/03745

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 A61K31/425 C07D417/12 C07D233/48 C07D285/12 C07D213/75
 C07D241/20 C07D231/40 C07D237/20 C07D239/42 C07D307/66
 A61P03/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

CHEM ABS Data, PAJ, EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FORD, ROGER E. ET AL: "Synthesis and quantitative structure-activity relationships of antiallergic 2-hydroxy-N-(1H-tetrazol-5-yl)benzamides and N-(2-hydroxyphenyl)-1H-tetrazole-5-carboxamides" J. MED. CHEM. (1986), 29(4), 538-49 , XP001026086 formula 2 table 1 --- -/--	3

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

10 October 2002

Date of mailing of the international search report

23/10/2002

Name and mailing address of the ISA

 European Patent Office, P.B. 5816 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Seelmann, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 02/03745

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ZHANG, SAN-QI ET AL: "Synthesis based on affinity separation (SAS): separation of products having barbituric acid tag from untagged compounds by using hydrogen bond interaction" SYNLETT (2001), (5), 590-596 , XP001106577 page 591	3
X	PATENT ABSTRACTS OF JAPAN vol. 1999, no. 06, 31 March 1999 (1999-03-31) -& JP 08 173525 A (LADD LELAND L), 9 July 1996 (1996-07-09) abstract	3
X	US 4 146 631 A (MARSHALL STUART M ET AL) 27 March 1979 (1979-03-27) column 2, line 17 -column 3, line 18; claim 1	3
X	WO 01 19788 A (COR THERAPEUTICS INC) 22 March 2001 (2001-03-22) claim 1; example 196	3
A	WO 01 44216 A (HOFFMANN LA ROCHE) 21 June 2001 (2001-06-21) cited in the application the whole document	1-16

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 02/03745

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 02 03745

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

The present claims relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. For the present scope of claim 3 the initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claim may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claim is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds of formula (I) with R2 = O-CH2-Phenyl or O-CH2-CH2-Phenyl.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 02/03745

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 08173525	A	09-07-1996	US 5792109 A EP 0699448 A1	11-08-1998 06-03-1996
US 4146631	A	27-03-1979	GB 1561350 A AU 514372 B2 AU 3031277 A BE 860513 A1 CA 1094565 A1 DE 2749518 A1 DK 489777 A FR 2370041 A1 HU 176842 B IE 45723 B1 JP 1357474 C JP 53059669 A JP 61021233 B LU 78459 A1 NL 7711901 A ZA 7706555 A	20-02-1980 05-02-1981 10-05-1979 05-05-1978 27-01-1981 18-05-1978 06-05-1978 02-06-1978 28-05-1981 03-11-1982 13-01-1987 29-05-1978 26-05-1986 14-07-1978 09-05-1978 30-08-1978
WO 0119788	A	22-03-2001	AU 7486600 A AU 7486700 A CZ 20020959 A3 EP 1216231 A2 EP 1216228 A2 NO 20021229 A NO 20021230 A WO 0119798 A2 WO 0119788 A2 US 2002091116 A1 AU 4535301 A AU 5078301 A WO 0164642 A2 WO 0164643 A2 US 2002002183 A1	17-04-2001 17-04-2001 17-07-2002 26-06-2002 26-06-2002 21-05-2002 21-05-2002 22-03-2001 22-03-2001 11-07-2002 12-09-2001 12-09-2001 07-09-2001 07-09-2001 03-01-2002
WO 0144216	A	21-06-2001	AU 2365201 A WO 0144216 A1 EP 1242397 A1 NO 20022863 A US 6353111 B1	25-06-2001 21-06-2001 25-09-2002 14-06-2002 05-03-2002

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)